

Colour perception in ADHD

Tobias Banaschewski,¹ Sinje Ruppert,¹ Rosemary Tannock,² Björn Albrecht,¹ Andreas Becker,¹ Henrik Uebel,¹ Joseph A. Sergeant,³ and Aribert Rothenberger¹

¹Child and Adolescent Psychiatry, University of Göttingen, Germany; ²The Hospital for Sick Children, Toronto, Canada; ³Department of Clinical Neuropsychology, Faculteit der Psychologie, Vrije Universiteit, Amsterdam, The Netherlands

Attention-deficit/hyperactivity disorder (ADHD) is associated with unexplained impairments on speeded naming of coloured stimuli. These deficits may reflect hypofunctioning retinal dopaminergic mechanisms impairing particularly blue–yellow colour discrimination. Colour perception and rapid colour naming ability were investigated in 14 children with ADHD and 13 healthy peers matched for age, gender, and IQ, using the Farnsworth–Munsell 100 Hue Test (FMT) and the Stroop–Colour–Word test. Children with ADHD committed more errors on the FMT, particularly on discrimination of colours along the blue–yellow axis, and were slower on Stroop subtests involving colour naming. However, the latter deficit was accounted for similarly by blue–yellow and red–green discrimination abilities. Blue–yellow colour perception problems in ADHD contribute to but do not fully explain the observed slowed colour naming. **Keywords:** Colour vision, attention-deficit hyperactivity disorder, dopamine, Stroop–Colour–Word test. **Abbreviations:** CBCL: Child Behavior Checklist; FBB-HKS: Fremdbeurteilungsbogen Hyperkinetische Störung; FMT: Farnsworth–Munsell 100 Hue Test; LPS: Leistungsprüfsystem.

Attention-deficit/hyperactivity disorder (ADHD) is a common childhood-onset disorder characterised by inattention, overactivity, and impulsiveness. The dopaminergic neurotransmitter system is believed to play a central role in its pathophysiology (e.g., Banaschewski et al., 2005; Castellanos & Tannock, 2002; Sagvolden, Johansen, Aase, & Russell, 2004; Taylor et al., 2004). By contrast to well-documented deficits in higher cortical functions, individuals with ADHD also exhibit unexplained problems in the speeded processing of coloured stimuli. For example, they exhibit slower naming speed for colours on various rapid naming tasks (e.g., the Rapid Automated Naming Test, the Stroop–Colour–Word Test), but tend not to exhibit slower naming of letters, words, or digits (e.g., Rucklidge & Tannock, 2002; Tannock, Martinussen, & Frijters, 2000; van Mourik, Oosterlaan, & Sergeant, 2005). A recent study of selective attention to colour, using event-related potentials (ERP), found that boys with ADHD exhibit a *perceptual deficit* in selection between visual stimuli on the basis of *colour* (red versus blue) as well as in later (semantic) stages of visual selective attention (van der Stelt, van der Molen, Boudewijn Gunning, & Kok, 2001).

To date, there is no adequate explanation to account for these observed difficulties in rapid processing of coloured stimuli in individuals with ADHD. However, poor performance on tasks involving speeded colour processing may reflect subtle visual impairments as well as (or rather than) impairments in higher-order cognitive function (van Boxtel, ten Tusscher, Metsemakers, Willems, & Jolles, 2001). Colour perception is based on three cone photoreceptor types, maximally sensitive to long, middle and short wavelengths, that constitute two function-

ally and anatomically distinct systems at the retina; a 'red–green' system and a 'blue–yellow' pathway. Colour perception is controlled, at least partly, by retinal dopaminergic neurons which are involved in controlling the coupling of horizontal and amacrine cell lateral systems, the organisation of the ganglion cell and the bipolar cell receptive fields and modulation of the physiological activity of photoreceptors (for reviews see: Djamgoz, Hankins, Hirano, & Archer, 1997; Hart, 1987; Masson, Mestre, & Blin, 1993; Witkovsky, 2004). Thus, retinal and central dopaminergic alterations may occur together and abnormalities of the dopaminergic system associated with ADHD may result in impaired colour perception which then may be expressed as slower colour naming (Tannock, Banaschewski, & Gold, under review).

A retinal deficit of dopamine is reflected particularly in alterations along the blue–yellow axis (Djamgoz et al., 1997; Hart, 1987; Masson et al., 1993). Moreover, a selective impairment of the blue–yellow vision system suggests a retinal location of the disturbance rather than a central one (Hart, 1987). Specific blue–yellow colour vision disturbances are found in various other disorders involving altered dopaminergic neurotransmission, such as Tourette syndrome (e.g., Melun, Morin, Muise, & DesRosiers, 2001), Parkinson's disease (e.g., Haug, Kolle, Trenkwalder, Oertel, & Paulus, 1995), and Huntington's disease (e.g., Büttner, Schulz, Kuhn, Blumenschein, & Przuntek, 1994). Converging changes of retinal dopamine levels associated with cocaine-withdrawal (e.g., Roy, Roy, Berman, & Gonzalez, 2003) and normal aging (e.g., Djamgoz et al., 1997) have also been associated with blue–yellow colour vision losses. Finally, acquired dyschromatopsias due to exposure to environmental pollutants (Gobba &

Cavalleri, 2003) usually impair blue-yellow colour discrimination, while congenital dyschromatopsias more frequently result in red-green deficits (Hart, 1987).

Methods

The 14 children with ADHD-combined type were selected from sequential referrals to the outpatient clinic of the department of Child and Adolescent Psychiatry of the University of Göttingen. Diagnosis was based on information ascertained from clinical interview with the parents, teacher reports, behavior rating scales, and medical reports. Rating scales included the parent-rated Child Behavior Checklist (CBCL; Achenbach, 1991), and a German version of the ADHD symptom list (FBB-HKS; Bruehl, Doepfner, & Lehmkuhl, 2000). Those children using methylphenidate were free of medication for at least 48 hr before being tested.

All children underwent standardised IQ testing, as well as testing of spelling abilities and word fluency (LPS; Horn, 1983). All children were free of ophthalmologic disorders or congenital colour blindness and had a full-scale IQ above 85. Also, only 3 children in the ADHD group and 2 in the control group met diagnostic criteria for dyslexia.

Colour discrimination ability was investigated individually using the Farnsworth–Munsell 100 Hue Test (FMT; Farnsworth, 1943), which is a widely used clinical instrument for measuring chromatic discrimination (Kinnear & Sahraie, 2002). It consists of four trays containing a total of 85 removable colour reference caps (incremental hue variation) spanning the visible spectrum. Colour vision abnormalities and aptitude were determined by the ability of the child to place the colour caps in order of hue. Error scores are measures of accuracy in arranging the caps so as to form a gradual transition in chroma between two anchor caps reflecting the number of misplacements. Blue–yellow and red–green partial error scores and the total error score were computed. All testing was performed under standard light conditions in the same room and at the same place. The illuminance was maintained at 325 lux, measured with a LT Lutron LX-101 Lux-meter.

The groups were compared for differences in Stroop test and colour perception performance as indicated by the FMT using analyses of variance (ANOVAs) with 'group' as between-subjects factor and discrimination along 'colour axis' (BY: blue-yellow; RG: red-green) as within-subject factor. In case of significant interaction effects, subsequent *t*-tests were computed. Furthermore, subsequent ANCOVAs with the covariates BY, respectively RG, and Pearson correlations between Stroop Colour naming and BY, respectively RG, were calculated. All analyses were performed with SPSS 11.5.2.1.

Table 1 Sample description and results

Measure	Controls (<i>N</i> = 13 (2; 2) ^a		ADHD <i>N</i> = 14 (1; 3) ^a		ANOVA	
	Mean	SD	Mean	SD	<i>F</i>	<i>p</i> <
Full-scale IQ	108.9	10.0	104.4	7.6	1.8	ns
Age (years)	10.9	.7	110.5	1.0	1.2	ns
Spelling abilities (T-score)	49.9	13.1	46.9	9.0	.5	ns
Word fluency (n/3*min)	27.4	6.9	26.8	11.6	.0	ns
Farnsworth–Munsell 100 Hue						
Total error score	54.5	31.1	86.8	33.6	6.7	.016
Partial error score blue–yellow	28.6	19.1	50.4	22.0	7.5	.011
Partial error score red–green	25.9	12.9	36.4	15.1	3.7	.065
Stroop						
Word (sec/72 items)	44.5	10.4	56.9	26.6	2.5	ns
Colour (sec/72 items)	66.2	12.3	83.7	27.9	4.4	.047
Colour/Word (sec/72 items)	116.8	30.9	174.7	56.2	10.8	.003
Nomination ^b (<i>t</i> score)	52.0	7.5	51.6	12.5	.01	ns
Selectivity ^{bc} (<i>t</i> score)	53.8	8.4	47.4	13.7	2.1	ns

^aTotal number (number of girls; number of probands with dyslexia). ^bNote: higher *t* scores = >better performance. ^cAbility to resist interference.

higher Total Error Score than did controls ($df = 1$, $F = 6.7$, $p < .02$; partial $\eta^2 = .15$). However, these main effects were modified by a significant interaction effect between the factors 'colour axis' and 'group' ($df = 25$, $F = 4.5$, $p < .05$; partial $\eta^2 = .28$). Post hoc analyses revealed that the colour perception impairment of children with ADHD was particularly pronounced along the blue–yellow axis as opposed to the red–green axis (see Table 1), indicating that children with ADHD make more blue/yellow errors ($F = 7.5$, $p < .011$; partial $\eta^2 = .23$) but not more red/green errors compared to normal controls; furthermore, children with ADHD make more blue/yellow errors compared to red/green errors ($df = 13$; $T = 3.1$, $p = .009$; partial $\eta^2 = .42$), whereas the controls didn't differ in number of errors on both axes ($df = 12$; $T = 1.0$, $p = .331$).

Analysis of the Stroop data indicated that the two groups did not differ in the speed of naming words in the Word condition. However, the children with ADHD were significantly slower naming colours in both the Colour and Colour/Word conditions. The factor 'group' accounted for 14.8% of the variance of Stroop-Colour condition, corresponding to a large effect according to Cohen.

To determine whether colour perception abilities contribute to variance in the Stroop colour naming condition, Pearson correlations between Stroop-Colour naming and 'blue–yellow colour discrimination' (BY), respectively 'red–green colour discrimination' (RG), (for the total group and separately for ADHD and controls), as well as ANCOVAs with the covariates BY, respectively RG, were calculated. The correlations between Stroop Colour naming and colour discrimination (BY, RG) were all insignificant, probably because of small sample sizes. Interestingly, correlation coefficients were consistently larger for RG with Stroop-Colour naming than for BY with Stroop-Colour naming (ADHD: $r_{RG} = .27$, r_{BY} :

$r = .19$; controls: $r_{RG} = .28$, $r_{BY} = .1$; total group: $r_{RG} = .36$, $r_{BY} = .31$). Correspondingly, ANCOVAs revealed partial eta squared values of .024 for the covariate BY, respectively .066 for the covariate RG, which can be regarded as a small effect size of the covariate BY according to Cohen ($f = .16$), respectively a medium ($f = .27$) effect size of the covariate RG.

Discussion

Children with ADHD exhibited more errors than their typically developing peers on a colour perception test (FMT), particularly with blue–yellow stimuli. Our finding is consistent with previous studies in other disorders involving altered dopaminergic mechanisms, which have found that discrimination along the blue–yellow axis (compared to the red–green axis) is particularly impaired (e.g., Büttner et al., 1994; Gobba & Cavalleri, 2003; Haug et al., 1995; Melun et al., 2001; Roy et al., 2003), indicating subtle problems in the blue–yellow mechanism and changes in retinal dopaminergic mechanisms.

The fundamental mechanisms causing a specific retinal impairment of colour discrimination along the blue–yellow axis in dopaminergic disorders remain unclear. Short wavelength sensitive cones may be more fragile than long and medium wavelength sensitive cones or their relative scarcity and anatomical distribution may be responsible for the greater vulnerability of the blue–yellow perception by alterations of the dopaminergic system (Djamgoz et al., 1997; Hart, 1987; Masson et al., 1993; Witkovsky, 2004).

A second and predicted finding was that children with ADHD were slower in naming the colours of stimuli on both the Colour and Colour-Word conditions of the Stroop, but did not differ in naming

speed on the Word condition, nor in Stroop interference. This result is consistent with findings from numerous studies reporting colour naming difficulties in children, adolescents and adults with ADHD (e.g., Banaschewski et al., 2005; Rucklidge & Tannock, 2002; van Mourik et al., 2005).

However, while the Stroop colour naming condition is dependent in part upon an intact visual function, we did not find robust evidence of a specific relationship between colour perception ability along the blue–yellow axis and colour naming on the Stroop. Rather, an equal amount of variance of the Stroop–Colour condition could be accounted by discrimination ability along the blue–yellow as along the red–green axis. Thus our findings mitigate the argument that colour perception has a substantial influence on Stroop performance.

Several methodological limitations must be considered when interpreting the findings from the present study. First, the children with ADHD were obtained through referrals to a paediatric psychiatry clinic and thus it is unknown whether findings are generalizable to non-clinical samples. Second, we used a 48 hr washout period prior to testing for those children receiving stimulant treatment, which might not have been of sufficient duration to ensure that there were not any residual effects of the stimulant drug on dopamine neurotransmission in children with ADHD. However, if this were the case, then the effect would be to attenuate rather than accentuate the group differences in colour vision. Third, we relied on a single clinical measure of colour perception (FMT), which requires the participant to sequence coloured caps on the basis of hue. FMT is often the clinical test of choice for assessing acquired colour vision defects (which typically affect the blue–yellow mechanism). However, it requires movement execution as well as sustained attention: motor impairments have been associated with FMT errors scores in patients with Parkinson's disease (e.g., Haug et al., 1995). But problems in attention and/or motor execution, which are associated with ADHD, would not account for the children's differential performance with blue–yellow versus red–green stimuli. Nonetheless, future studies should include additional measure of colour perception.

Conclusion

To our knowledge, this is the first study of colour perception in ADHD. Although distorted colour discrimination abilities may reflect alterations of more central mechanisms of the vision system as well, the present findings of specific blue–yellow colour perception problems suggest colour vision impairment that is related to retinal dopaminergic function in children with ADHD. However, performance deficits on Stroop colour naming seem not to be fully

attributable to colour vision impairment. Further research on colour perception in ADHD is warranted, using more precise approaches to measurement that permit an evaluation of neuronal integrity, such as colour visual evoked potentials.

Acknowledgements

This article is part of the international and interdisciplinary project 'ADHD: From genes to therapy' (Project leader: Terje Sagvolden) conducted at the Centre for Advanced Study (CAS) in Oslo, Norway, 2004–2005, in which several authors (TB, JS, AR, RT) participated.

Correspondence to

T. Banaschewski, Child and Adolescent Psychiatry, University of Göttingen, von-Siebold-Str. 5, D - 37075 Göttingen, Germany; Tel: 0049-551-396727; Fax: 0049-551-398120; Email: tbanasc@gwdg.de

References

- Achenbach, T.M. (1991). *Manual for Child Behavior Checklist/4-18 and 1991 Profile*. Burlington, VT: University of Vermont Department of Psychiatry.
- Banaschewski, T., Hollis, C., Oosterlaan, J., Roeyers, H., Rubia, K., Willcutt, E., et al. (2005). Towards an understanding of unique and shared pathways in the psychopathophysiology of ADHD. *Developmental Science*, 8, 132–140.
- Bäumler, G. (1985). *Farbe-Wort-Interferenztest (FWIT) nach JR Stroop*. Göttingen: Hogrefe.
- Bruehl, B., Doepfner, M., & Lehmkuhl, G. (2000). Der Fremdbeurteilungsbogen für hyperkinetische Störungen (FBB-HKS) Prävalenz hyperkinetischer Störungen im Elternurteil und psychometrische Kriterien. *Kindheit und Entwicklung*, 9, 116–126.
- Büttner, T., Schulz, S., Kuhn, W., Blumenschein, A., & Przuntek, H. (1994). Impaired colour discrimination in Huntington's disease. *European Journal of Neurology*, 1, 153–157.
- Castellanos, F.X., & Tannock, R. (2002). Neuroscience of attention-deficit/hyperactivity disorder: The search for endophenotypes. *Nature Reviews Neuroscience*, 3, 617–628.
- Djamgoz, M., Hankins, M., Hirano, J., & Archer, S. (1997). Neurobiology of retinal dopamine in relation to degenerative states of the tissue. *Vision Research*, 37, 3509–3529.
- Farnsworth, D. (1943). The Farnsworth Munsell 100-Hue dichotomous tests for colour vision. *Journal of the Optical Society of America*, 33, 568–576.
- Gobba, F., & Cavalleri, A. (2003). Color vision impairment in workers exposed to neurotoxic chemicals. *NeuroToxicology*, 24, 693–702.
- Hart, W.M. (1987). Acquired dyschromatopsias. *Surveys in Ophthalmology*, 32, 10–31.
- Haug, B.A., Kolle, R.U., Trenkwalder, C., Oertel, W.H., & Paulus, W. (1995). Predominant affection of the

- blue cone pathway in Parkinson's disease. *Brain*, 118(Pt 3), 771–778.
- Horn, W. (1983). *Das Leistungsprüfsystem (LPS)* (vol. 2. Ausgabe). Bern: Hans Huber Verlag.
- Kinnear, P., & Sahraie, A. (2002). New Farnsworth–Munsell 100 hue test norms of normal observers for each year of age 5–22 and for age decades 30–70. *British Journal of Ophthalmology*, 86, 1408–1411.
- Masson, G., Mestre, D., & Blin, O. (1993). Dopaminergic modulation of visual sensitivity in man. *Fundamental and Clinical Pharmacology*, 7, 449–463.
- Melun, J.P., Morin, L.M., Muise, J.G., & DesRosiers, M. (2001). Color vision deficiencies in Gilles de la Tourette syndrome. *Journal of Neurological Science*, 186, 107–110.
- Roy, A., Roy, M., Berman, J., & Gonzalez, B. (2003). Blue cone electroretinogram amplitudes are related to dopamine function in cocaine-dependent patients. *Psychiatric Research*, 117, 191–195.
- Rucklidge, J.J., & Tannock, R. (2002). Neuropsychological profiles of adolescents with ADHD: Effects of reading difficulties and gender. *Journal of Child Psychology and Psychiatry*, 43, 988–1003.
- Sagvolden, T., Johansen, E.B., Aase, H., & Russell, V.A. (2004). A dynamic developmental theory of attention-deficit/hyperactivity disorder (ADHD) predominantly hyperactive/impulsive and combined subtypes. *Behavioral and Brain Science*.
- Tannock, R., Martinussen, R., & Frijters, J. (2000). Naming speed performance and stimulant effects indicate effortful, semantic processing deficits in attention-deficit/hyperactivity disorder. *Journal of Abnormal Child Psychology*, 28, 237–252.
- Tannock, R., Banaschewski, T., & Gold, D.. Color naming deficits and attention-deficit/hyperactivity disorder: A retinal dopaminergic hypothesis. *Behavioral and Brain Functions*. Manuscript under review.
- Taylor, E., Dopfner, M., Sergeant, J., Asherson, P., Banaschewski, T., Buitelaar, J., et al. (2004). European clinical guidelines for hyperkinetic disorder – first upgrade. *European Child and Adolescent Psychiatry*, 13(Suppl. 1), I7–I30.
- van Boxtel, M.P., ten Tusscher, M.P., Metsemakers, J.F., Willems, B., & Jolles, J. (2001). Visual determinants of reduced performance on the Stroop color-word test in normal aging individuals. *Journal of Clinical and Experimental Neuropsychology*, 23, 620–627.
- van der Stelt, O., van der Molen, M., Boudewijn Gunning, W., & Kok, A. (2001). Neuroelectrical signs of selective attention to color in boys with attention-deficit hyperactivity disorder. *Brain Research. Cognitive Brain Research*, 12, 245–264.
- van Mourik, R., Oosterlaan, J., & Sergeant, J.A. (2005). The Stroop revisited: A meta-analysis of interference control in AD/AD. *Journal of Child Psychology and Psychiatry*, 46, 150–165.
- Witkovsky, P. (2004). Dopamine and retinal function. *Documenta Ophthalmologica*, 108, 17–40.

Manuscript accepted 25 May 2005