LIFELOOP

AI-Assisted Continuous Care for Diabetes and Hypertension

A Bangla-first, patient-centred care companion that connects daily health tracking with better clinical follow-up.

Document type	Detailed concept paper
Competition track	HealthSphere AI / ChronicCare AI
Prepared by	Team Sic Parvis Magna
Version	1.0 | 12 October 2026

THE CORE IDEA

Help people with diabetes and high blood pressure understand changes in their daily health, and help their doctors follow those changes between visits. LifeLoop supports care; it does not replace a doctor.

Status: Concept-stage proposal. The risk model, clinical benefit, and patient outcomes described here are research goals, not proven results. No clinical validation or regulatory approval is claimed.

Executive Summary

People living with Type 2 diabetes and high blood pressure often need to manage their health every day, not only when they visit a clinic. A doctor can give advice based on a visit and the information available at that time. However, it can be difficult for a doctor to see what happens during every day between appointments: what the patient eats, how active they are, how well they sleep, whether their blood glucose or blood pressure is changing, and whether their daily routine is becoming harder to maintain.

LifeLoop is a proposed digital care companion for this gap. It combines simple home measurements, optional activity and sleep data, easier lifestyle logging, personal health trends, and a secure way to share useful summaries with a chosen doctor or care team. It is designed first for people in Bangladesh, with Bangla-first interaction and support for familiar local meals and daily routines.

The central research idea is to test whether changes across a person's own health history and several available signals can help identify concerning deterioration better than relying only on separate fixed thresholds. The system will begin with clear rules and simple, testable machine-learning models. It will show uncertainty when data are missing or weak. An AI language model may help explain already-computed information in simple language, but it will not diagnose a patient, choose medication doses, or control urgent safety alerts.

LifeLoop follows a B2C + B2B2C model. Patients use the mobile app, while doctors and clinics can use a separate dashboard with patient consent. The first version will focus on a small number of features that can be built and tested: reliable entry of glucose and blood pressure readings, easy daily logs, trend views, simple lifestyle support, and a patient-approved doctor report. Advanced wearable integration, personalised learning, and new blood pressure hardware are later research directions, not requirements for the first prototype.

> **WHAT LIFELOOP WILL NOT CLAIM**
>
> The first version will not claim to predict heart attacks, diagnose emergencies, replace clinical judgment, or measure blood pressure accurately without a suitable validated device. These would require separate evidence and clinical validation.

Concept at a Glance

Item	Proposal
Primary users	Adults with Type 2 diabetes and hypertension who need regular home tracking and follow-up.
Main problem	Important changes in health and daily habits can occur between clinic visits, while tracking is often split across devices, notebooks, and memory.
Core product	Patient mobile app + secure doctor/clinic dashboard + shared backend for data, analysis, and alerts.
Main R&D question	Can personal baselines and multiple health trends improve the identification of pre-defined deterioration compared with simple threshold rules, without creating too many false alerts?
First technical approach	Measurement-quality checks, personal trend features, transparent rule-based safety, and baseline machine-learning models where data support them.
Business model	Free/basic patient access with optional paid features; clinic or care-program subscriptions; optional consultation booking kept separate from monitoring.
Evidence status	Proposal only. Dataset access, clinical partnership, real-world model performance, and commercial demand must be confirmed.

1. The Problem

1.1 Daily care continues after the appointment

A clinic visit provides an important but limited view of a person's health. The patient may have a measurement taken in the clinic, receive advice, and then return to normal life. During the following days, the patient's glucose, blood pressure, eating, movement, sleep, and medication routine may change. The next visit may be weeks or months later. A doctor cannot personally observe every part of every patient's day, and many patients find it hard to record everything accurately by hand.

This is not a problem caused by doctors doing a poor job. It is a gap between clinic-based care and daily life. LifeLoop is meant to help both sides: patients get a simpler way to follow their routine, and clinicians receive a clearer summary of changes that may need attention.

1.2 Why focus on diabetes and hypertension?

Type 2 diabetes and high blood pressure are long-term conditions that need ongoing care. They can occur together, and poor control can increase the risk of serious complications over time. International Diabetes Federation estimates indicate that about 13.9 million adults aged 20-79 were living with diabetes in Bangladesh in 2024. This is an estimate of the national disease burden, not a count of people who would use or pay for LifeLoop. [1]

Daily habits such as food choices and physical activity are part of diabetes self-management, but collecting reliable daily information can take time. [2] For this reason, the first target group will be adults managing both Type 2 diabetes and hypertension, rather than every patient or every chronic disease at once.

1.3 The specific gap LifeLoop will study

> **THE SILENT GAP**
>
> A patient can have a reading that does not cross a single fixed alert threshold while their longer-term pattern is changing. LifeLoop will test whether personal trends across available signals help flag such changes early enough for a useful follow-up. This is a research question, not an established result.

The project will not assume that any single change means that a dangerous event will happen. It will study defined signs of deterioration, such as sustained worsening of glucose or blood pressure control, using outcomes selected with qualified clinical partners. Predicting a heart attack or another sudden emergency is outside the first version.

2. The Proposed Solution

LifeLoop will turn scattered daily health information into a simple routine: measure, record, understand the trend, take a small safe action, and share the important changes with a doctor when needed. The product should feel easy to use for a patient and useful rather than noisy for a clinician.

Step	What happens	Why it matters
1. Capture	The patient enters glucose and blood pressure readings from home devices. Optional data may come from supported wearables or phone health services.	Starts with devices people may already use, without requiring an expensive wearable.
2. Check	The app checks whether the entry is complete and helps the patient follow a proper measurement routine. Unusual readings are not silently deleted.	Reduces avoidable recording errors and keeps the original measurement visible.
3. Compare	The system compares recent values with the patient's own history and shows simple trends.	Helps explain what is changing, not just whether one number is high or low.

Step	What happens	Why it matters
4. Support	The patient gets a small, non-medication lifestyle prompt based on approved content and their selected goals.	Makes daily care easier without acting as a doctor.
5. Share or alert	When a rule or validated model indicates a concern, the system follows a reviewed alert path and can prepare a short report for the assigned care team.	Connects daily life with clinical follow-up.

2.1 Example user journey

A patient records blood pressure and glucose over several days. The first reading is not enough to tell a full story. Over time, the app notices that several available measurements are moving away from that person's usual pattern. The app checks data quality and the relevant safety rules. If a pre-defined concern is met, it shows a clear message, explains which measurements drove the concern, and follows the clinic's agreed contact process. The doctor receives a short timeline and can decide what follow-up is appropriate. LifeLoop does not tell the patient to change a prescribed dose.

2.2 Daily lifestyle support

- Meal photo assistance: the patient can take a photo of a meal. AI may suggest food types and a rough portion estimate. The patient can correct the result. The photo cannot reliably reveal every ingredient, exact portion, cooking method, or nutrient value, so the estimate must be labelled as an estimate. [5]
- Walking and activity: supported phone or wearable data can show steps and activity trends. It will not claim to know all movement or all time spent sitting.
- Sleep: supported devices may provide an estimate. If there is no suitable device or data, the app may ask a simple question rather than pretend to measure sleep precisely.
- Weight: the patient may record weight over time. Weight can help show a longer trend, but it cannot prove what the patient ate on a particular day.
- Simple coaching: early versions will use a small set of clinician-reviewed, non-medication prompts. More adaptive coaching will only be explored after safety and usefulness are tested.

3. Users and Their Needs

User	Need	LifeLoop value
Patient	An easy daily routine, simple language, less manual entry, and a clear view of personal trends.	Bangla-first screens, reminders, supported device data, photo-assisted meal logging, trend summaries, and control over sharing.
Doctor or nurse	A quick way to review changes between visits without sorting through long, messy records.	A short patient timeline, data quality notes, alert history, and a one-page visit brief.
Clinic or chronic-care programme	A consistent follow-up process for many enrolled patients.	A dashboard for assigned patients, alert ownership, follow-up status, and simple programme reports.
Family member or community health worker	A patient-approved way to support routine and follow-up.	Limited role-based access, selected reminders, and agreed escalation support when enabled.

Patients should be able to choose whether to share information and with whom. A QR code can make it easier to connect a patient account with a doctor or clinic, but the QR code itself is not a substitute for secure sign-in, identity checks, access control, and consent.

4. What Makes LifeLoop Different?

LifeLoop is not different simply because it uses AI, displays charts, or has a doctor dashboard. These features already exist in many digital health products. Its proposed difference is the way it combines a personal trend view, easier daily logging, transparent warnings, and consent-based clinical follow-up in one simple workflow.

Layer	Proposed distinction	How it will be tested
Personal baseline	Compare recent readings with the person's own history, instead of using only population-wide fixed values.	Compare against a simple threshold-based baseline and report whether performance improves.
Multi-signal trend	Test whether more than one reliable signal adds value when identifying pre-defined deterioration.	Ablation tests: compare glucose/BP alone with additional signals, only where data exist.
Honest uncertainty	Show when data are missing, old, or too limited for a dependable estimate.	Measure missing-data behaviour, calibration, and the frequency of unsupported outputs.
Safety separated from language AI	Clinical safety rules run independently of the language model. The language model explains results; it does not decide emergency thresholds.	Test that critical rules still work when the language model is unavailable or produces an invalid answer.
Local and usable design	Use Bangla-first interaction, local meal labels, low-burden input, and a doctor-linked workflow.	Observe task completion, user effort, retention, and clinician feedback in pilot testing.

IMPORTANT DISTINCTION

A common app feature is not automatically research novelty. LifeLoop's research contribution must come from a clear, testable question and a fair comparison against existing simple methods. Results will be reported even if they show no improvement.

5. Research and Development Plan

5.1 Main research question

PRIMARY RESEARCH QUESTION

For adults with Type 2 diabetes and hypertension, can patient-specific baseline and trend features from repeated glucose and blood pressure readings, with selected additional signals where available, improve the identification of pre-defined health deterioration compared with simple single-threshold rules, without creating an unacceptable number of false alerts?

5.2 Supporting questions

1. Which trend features are useful and stable when readings are incomplete or taken at different times?
2. Does adding activity, sleep, resting heart rate, or meal estimates improve the model enough to justify the extra work and device needs?
3. Can the system show uncertainty clearly when there is not enough good data?
4. Can the app reduce daily logging effort and help patients stay engaged?
5. Can a concise health brief help clinicians review the period between visits more easily?

5.3 Proposed method

The first research phase will use available datasets that fit the question, after checking their access conditions, data fields, licensing, and suitability. The original concept lists CGMacros and AI-READI as candidate sources for data and

signal testing. Their use is not guaranteed, and their data may not represent Bangladeshi patients. Synthetic data can help test software logic, edge cases, and alert routing, but it cannot prove that a model works for real patients.

The project will begin with transparent methods: trend summaries, patient-specific baselines, simple threshold rules, and a small number of tabular machine-learning models such as Random Forest or XGBoost if the available data support them. A complex custom foundation model is not needed for the first research question. A language model may be used for plain-language summaries after the risk features and rules have been computed, with guideline retrieval and strict limits on what it may say.

5.4 Candidate trend design

- Recent window: a 14-day window may be tested to describe shorter-term changes, where enough readings are available.
- Longer baseline: a 90-day history may be used as a reference for slower changes if the patient has enough reliable data.
- Patient-specific features: direction of change, repeated high or low measurements, glucose variability where there are enough data, time-of-day patterns, and the reliability of each input.
- Additional data: steps, sleep estimates, resting heart rate, weight, or meal labels may be tested one at a time to determine whether they add real value.
- Missing-data handling: the system must not treat missing measurements as normal measurements. If data are too limited, it will show low confidence or say that it cannot assess the trend.

These windows and features are starting design choices from the concept, not clinically proven settings. The clinical adviser and research evaluation may require changes. Any label used to train or test a prediction model must be defined independently of the model inputs where possible, to reduce circular reasoning and data leakage.

5.5 Define the outcome before training

The team must agree with a qualified clinical collaborator on what counts as a useful warning. A first study may focus on a clearly defined period of worsening glucose or blood pressure control, or another clinician-reviewed event that can be measured consistently. “Heart attack” must not be used as the outcome unless the team has suitable labelled data, a sound study design, and the required clinical oversight. A system cannot claim that a pattern predicts a critical event just because it can reproduce a rule used to label that same pattern.

5.6 Evaluation approach

The model should be compared with a simple baseline, not only shown on selected examples. The study plan should be written before the final evaluation and follow good reporting practice for prediction models, such as the TRIPOD+AI guideline. [6] The team should keep test data separate from model development, check for data leakage, and report limitations plainly.

Measure	Question it answers
Sensitivity and specificity	How often does the system find the defined event, and how often does it avoid a false warning?
Positive predictive value	When the system raises a warning, how often does the defined event occur?
False alerts per patient-week	Will the system create too many interruptions for patients and care teams?
Calibration and uncertainty	Do estimated risk levels match observed outcomes, and does the system show less confidence when evidence is weak?
Lead time	How early does a useful warning appear before the pre-defined event, if the data support this measure?
Usability and engagement	Can patients complete key tasks, and do they continue using the system?
Clinician usefulness	Does the report save review effort or make follow-up clearer?

No performance number is presented as an achieved result in this concept paper. Any success thresholds will be agreed before evaluation with appropriate clinical and research input. Sample size for a real-patient study should be planned with a qualified research or statistical adviser.

6. Product Scope: What We Will Build First

6.1 Minimum viable product (MVP)

The MVP should prove the main problem and the workflow. It should not attempt to build every feature in the first release.

MVP feature	First-version behaviour
Patient account and consent	A patient can create an account, choose language, see what data are collected, and control approved sharing.
Glucose and BP records	Manual entry first; device integration only where reliable and practical. Source, time, and units are recorded.
Measurement support	Simple guidance for correct use of a suitable upper-arm BP cuff and clear reminders to follow the device instructions. [3]
Health timeline	Easy charts and summaries for repeated readings, with recent history compared to a personal baseline when enough data exist.
Lifestyle log	Optional meal photo, activity, sleep, and weight entries. AI results are estimates and can be corrected.
Simple daily support	A small set of reviewed, non-medication habit prompts. The app does not change treatment.
Doctor connection	Patient-approved doctor/clinic link, a concise summary, and visible access permissions.
Basic safety and escalation	Clinician-reviewed rules, understandable instructions, alert log, and an agreed person responsible for follow-up.

6.2 Later features, only after evidence

- Integration with more wearables, continuous glucose monitoring (CGM), sleep data, or night-time blood pressure, where device support and clinical usefulness are established.
- A more personalised machine-learning model, after it has passed suitable offline and clinical evaluation.
- Adaptive coaching that learns from response and adherence, only after the safe action set and monitoring process have been tested.
- A larger clinic network, family co-pilot options, and integration with clinical record systems where partners and standards permit.
- Separate research into cuffless blood pressure estimation or a new sensor. This is not part of the MVP.

6.3 Why not build everything now?

Too many features can make the product hard to explain, hard to test, and hard for patients to use. The first version should focus on a single meaningful result: make daily records easier to maintain and test whether personal trends help a care team spot defined problems sooner. Every extra sensor, AI agent, and dashboard feature should be added only when it supports that purpose.

7. System Design and Technical Plan

LifeLoop will use a shared backend so the mobile app and doctor dashboard use the same user permissions, data rules, and analysis services. The first prototype may use the existing web technology to test screens and workflow. A native Android app can be introduced for deeper device integration and approved background tasks. A Next.js web dashboard is suitable for doctors and clinic staff. Native development improves access to supported Android capabilities, but it does not remove Android permissions, battery, background-processing, or device limits.

PATIENT APP Kotlin Android / Bangla UI	**DATA ENTRY** Glucometer, BP cuff, optional lifestyle data	**DOCTOR WEB** Next.js dashboard / clinic workflow
IDENTITY & CONSENT Sign-in, permissions, audit trail	**SHARED BACKEND** Secure API, database, data-quality checks	**NOTIFICATION SERVICE** Approved contact route and delivery status
TREND ENGINE Baseline, features, missing-data handling	**SAFETY LAYER** Deterministic clinical rules; independent of LLM	**SUMMARY ENGINE** LLM may explain approved computed results

Concept architecture; final choices depend on prototype findings, device support, security review, and clinical needs.

7.1 The two safety loops

Loop	Role	Rule
Safety loop	Checks clinician-approved alert conditions and the required next step.	It must not wait for an LLM response. It must continue to work if the language model is offline.
Explanation loop	Turns computed trends into short, plain-language explanations and cites approved guidance when available.	It cannot create a diagnosis, change medication, hide a safety alert, or invent a measurement.

7.2 Use of AI agents during development

AI coding agents may help the team create separate modules in parallel, such as the patient app, backend, model pipeline, test suite, and doctor dashboard. Architecture decisions and clinical rules should be documented in one shared repository. Each module must still pass code review, automated tests, privacy checks, and integration tests. AI agents are a development method, not the product's research novelty, and they must not be allowed to invent clinical rules.

8. Blood Pressure Accuracy and Device Limits

LifeLoop will initially rely on measurements from suitable existing devices rather than claim to measure blood pressure on its own. Even a digital cuff reading can be affected by incorrect cuff size, body position, talking, recent activity, or poor measurement technique. CDC guidance advises resting quietly, supporting the back and arm, keeping feet flat, placing the cuff on bare skin, and not talking during measurement. [3] The app can make these steps easier to follow and record whether the patient used the suggested process.

Cuffless blood pressure devices, including some wearables, estimate blood pressure indirectly. The American Heart Association's 2025 scientific statement notes important questions about real-world accuracy, calibration, posture, movement, and clinical usefulness. [4] LifeLoop will not present an unvalidated watch or phone estimate as a medical-grade blood pressure reading.

> **FUTURE R&D OPTION**
>
> A separate project could test whether a new sensor or cuffless method can estimate blood pressure accurately across different users and daily conditions. It would need a suitable reference device, a careful protocol, diverse participants, repeated calibration checks, and clinical/statistical oversight. It is not required to prove the first LifeLoop concept.

9. Safety, Clinical Oversight, and Privacy

9.1 Clinical boundaries

- LifeLoop supports a care team; it does not replace a doctor, nurse, or emergency service.
- The app will not prescribe medication, advise a patient to change a dose, or tell a patient to stop a prescribed treatment.
- The first version will not claim to diagnose a heart attack or predict sudden emergencies.
- Any threshold, escalation rule, and lifestyle content used for care must be reviewed by a qualified clinical collaborator and aligned with relevant current guidance.
- A user-facing warning must state what triggered it, what the system can and cannot tell, and what safe next step the reviewed protocol requires.

9.2 Alert ownership

An alert is useful only if somebody is responsible for the next step. For a pilot, the team and partner clinic must agree who receives each type of alert, how receipt is recorded, how long a response may take, and what happens if the first person does not respond. LifeLoop must not promise 24-hour monitoring unless a real, staffed service has agreed to provide it. Critical safety rules will operate independently of the language model, and the app must give clear instructions for urgent symptoms rather than ask the user to wait for an AI answer.

9.3 Privacy and access

- Collect only the data needed for the agreed product and research aims.
- Use clear consent, role-based access, secure sign-in, encryption in transit and at rest where supported, and logs of access to patient records.
- A QR code can start a connection, but access must still require identity checks and the patient's approval. A shared QR image must not grant open access to the full health record.
- Patients should be able to review and revoke sharing according to the product's policy and applicable law.
- Research datasets should be de-identified where appropriate, stored with controlled access, and used only within approved terms.

9.4 Regulatory and research approval

Before a real-patient pilot or public clinical use, the team will seek advice on which Bangladesh requirements apply to the intended function of the software. The Directorate General of Drug Administration (DGDA) publishes medical-device guidance, including a guideline dated 15 February 2024, and has also announced guidance on AI and medical robotics in healthcare in 2026. The team should confirm the current, applicable requirements directly with DGDA and qualified advisers rather than assume that a prototype is exempt. [8] Research with people or identifiable health data will require appropriate ethics, consent, and institutional review where applicable.

The World Health Organization also cautions that digital health interventions should complement functioning health services, not replace them. Privacy, user needs, health-worker workflows, and local infrastructure must be considered. [7]

10. Feasibility and Implementation Roadmap

LifeLoop should be built in stages. Each stage has a clear question and a stop/go decision. The calendar below is a proposed plan; it is not a claim that partners, data, or approvals have already been secured.

Stage	Main work	Evidence needed before moving on
Stage 1: Problem and clinical review	Talk with patients, clinicians, and potential clinic partners. Agree on the first user group, key workflow, safety boundaries, and a measurable research outcome.	A written problem statement, user needs, clinician feedback, and a defined evaluation plan.
Stage 2: MVP prototype	Build patient data entry, measurement guidance, history charts, Bangla UI, consent-based sharing, a doctor summary, and basic deterministic rules.	Demonstration of the full journey with test accounts and no real clinical claims.
Stage 3: Technical research	Check available datasets and permissions. Compare simple rules with candidate models. Test missing data, false alerts, and language-model failure cases.	A fair offline evaluation with limitations and no leakage between development and test data.
Stage 4: Usability test	Run a small usability study with appropriate consent and oversight; measure effort, confusion, and completion of daily tasks.	Evidence that the main workflow is understandable and practical, plus a list of changes.
Stage 5: Supervised pilot	If approvals, data, and a partner clinic are secured, plan an 8-12 week pilot with clinician-defined monitoring and escalation.	Predefined outcomes, response ownership, safety monitoring, and an agreed plan for analysis.
Stage 6: Business test	Test a clinic subscription or care-program partnership and a basic patient tier. Keep consultation fees separate.	Real feedback on willingness to use/pay, cost to serve, retention, and clinic workload.

10.1 Feasibility principles

- Begin with ordinary home devices and a clear workflow rather than requiring every user to buy a smartwatch or CGM.
- Support low-effort input: remember the last used device, allow correction, use supported data import, and avoid asking the patient to fill out a long form every day.
- Design for Bangla and familiar foods, with plain explanations and large, readable controls.
- Do not treat a lack of data as proof that the patient is well.
- Avoid expensive hardware development until there is evidence that the software workflow solves a real problem and a strong reason to build new hardware.

11. Business Model and Adoption Plan

LifeLoop is designed as a B2C + B2B2C product. Patients need daily value, while doctors and clinics need a useful follow-up workflow. The first business test should find out which group gains enough value to pay, rather than assume that every patient or doctor will subscribe.

Channel	What is offered	Possible revenue
B2C: patient	Basic records, easy trends, daily routine support, and controlled sharing.	Free or low-cost entry; optional paid features such as richer trend reports or more convenient logging. Exact pricing requires user research.
B2B: doctor/clinic	Assigned patient list, patient-approved timeline, short visit reports, alert history, and follow-up status.	Subscription for clinic or clinician workspace, subject to demonstrated value and a sustainable support cost.
B2B2C: care programmes	A clinic, employer, insurer, or health programme offers LifeLoop to enrolled patients.	Contract or per-enrolled-user model; requires an operational partner and clear service responsibilities.

Channel	What is offered	Possible revenue
Optional consultation booking	Patients can book a doctor when they need clinical advice.	Consultation fee is paid separately to the provider; any platform fee must be clear. Monitoring must not force a consultation purchase.
Device partnerships: later	Help users access compatible devices from qualified suppliers.	Possible referral or distribution revenue only after supplier quality, accuracy, and applicable requirements are checked.

11.1 Why doctors might use it

The value for a clinician is not simply having another dashboard. LifeLoop must reduce the work needed to understand a patient's recent history, make the next visit better prepared, and show which alerts have been reviewed. The product should not flood doctors with low-value alarms. A clinic pilot must measure clinician workload and usefulness, not only app downloads.

11.2 First market

The first market will be a small, reachable group of adults with Type 2 diabetes and hypertension who already measure glucose or blood pressure at home and are willing to work with a doctor or clinic. The team should start with a partner clinic or adviser, learn which parts of the workflow are genuinely useful, and expand only after the evidence and service model are clear. The national diabetes estimate gives context for the problem, but it does not prove market demand or willingness to pay. [1]

12. Key Challenges and How We Will Address Them

Challenge	Why it matters	Planned response
Incorrect or inconsistent readings	Bad inputs can create false concern or false reassurance.	Measurement guidance, source/time tracking, data-quality flags, and suitable existing devices.
False alarms	Too many alerts can worry users and burden clinicians.	Compare with a simple baseline; measure false alerts per patient-week; use reviewed rules and a clear alert budget.
Sparse or missing data	The model may not have enough information to assess a trend.	Show uncertainty; avoid filling gaps with invented values; state when no assessment can be made.
Unproven model performance	An AI model may work on one dataset but fail on new patients.	Separate train/test data, evaluate outside the training set where possible, report calibration and limitations, and require clinical validation before use for care.
Food photo errors	A photo may miss portion size, hidden ingredients, or how food was cooked.	Label results as estimates, allow patient correction, and avoid using meal estimates as the only basis for a critical alert.
Low daily engagement	Patients may stop logging if it takes too long or feels boring.	Minimise required fields, use Bangla, offer optional automation, and test whether daily actions are realistic.
Alert not acted on	A notification does not help if nobody sees or owns it.	Define the named receiver, response process, backup path, and operating hours before a pilot.
Privacy and trust	Health information is sensitive and could be exposed or shared without permission.	Consent, least-privilege access, encryption, audit logs, revocation, and careful research data handling.
Limited clinical data	Public or synthetic datasets may not represent local patients and outcomes.	Treat them as early technical tests; secure ethical clinical collaboration and real-world evaluation before making clinical claims.

13. Success Measures

LifeLoop will measure both product usefulness and research quality. A good demo is not the same as proof of clinical benefit. Success measures will be agreed before testing and reported honestly, including negative or inconclusive findings.

Area	Measures to collect
Product usability	Time to enter a reading, task completion, user-reported ease, correction rate for AI meal labels, and common points of confusion.
Engagement	Daily/weekly logging, retention over time, reminders acted on, and reasons for dropping out.
Technical reliability	Missing-data handling, error rate, alert delivery status, system uptime, and whether safety rules work without the LLM.
Model quality	Sensitivity, specificity, positive predictive value, false alerts per patient-week, calibration, and lead time where measurable.
Clinician value	Time needed to review a patient, usefulness of the summary, number of actionable versus non-actionable alerts, and feedback on workload.
Business feasibility	Patient and clinic willingness to use/pay, onboarding cost, support needs, and renewal or continued-use intent.

14. Limits and Non-Goals

- LifeLoop is not an emergency service and does not promise continuous human monitoring unless a partner explicitly provides it.
- The first version does not diagnose, treat, or prescribe, and does not replace medical advice.
- The first version does not claim to predict heart attacks or sudden emergencies.
- The first version does not claim to measure blood pressure using only a phone camera or an unvalidated wearable.
- AI meal estimates, activity imports, and sleep estimates may be incomplete or wrong; users must be able to review them.
- Simulated data can test code and workflows, but cannot establish clinical effectiveness.
- A model trained on one population may not work equally well for Bangladeshi patients, older adults, rural users, or users with different devices. Local validation is needed.

15. Summary and Next Steps

LifeLoop aims to solve a clear and important problem: making daily diabetes and hypertension care easier for patients and making the time between clinic visits more visible to clinicians. Its strongest product idea is the connection between simple daily tracking and useful doctor follow-up. Its strongest proposed R&D question is whether personal baselines and multiple reliable signals can help identify defined deterioration better than simple fixed thresholds without causing too many false alerts.

The next step is not to add more technologies. It is to confirm the problem with patients and clinicians, agree on one measurable outcome, build a focused MVP, and test it against a clear baseline. The team should also identify a clinical adviser, confirm data access, and seek advice on the applicable approval and ethics process before testing with real patient data.

> **LIFELOOP IN ONE SENTENCE:** LifeLoop helps people with diabetes and high blood pressure manage their daily health and helps their doctors follow meaningful changes between visits - without pretending to replace clinical judgment.

References

The following sources support the problem context, product design choices, and proposed research methods. They do not validate LifeLoop itself.

[1] International Diabetes Federation. “Diabetes in Bangladesh.” IDF Diabetes Atlas, country report (2024 estimates). The page reports an estimated 13.9 million adults aged 20-79 living with diabetes in Bangladesh in 2024 and identifies Bangladesh Demographic and Health Survey 2017-18 among the data sources. Accessed 12 October 2026. Open source

[2] American Diabetes Association Professional Practice Committee. “Facilitating Positive Health Behaviors and Well-being to Improve Health Outcomes: Standards of Care in Diabetes—2026.” Diabetes Care, 2026. DOI: 10.2337/dc26-S005. Open source

[3] Centers for Disease Control and Prevention. “Measuring Your Blood Pressure.” Published 15 May 2024; updated 4 September 2026. Accessed 12 October 2026. Open source

[4] Cohen JB, Byfield RL, Hardy ST, et al. “Cuffless Devices for the Measurement of Blood Pressure: A Scientific Statement From the American Heart Association.” Hypertension. Published online 11 December 2025. DOI: 10.1161/HYP.0000000000000254. Open source

[5] Chotwanvirat P, Prachansuwan A, Sridonpai P, Kriengsinyos W. “Advancements in Using AI for Dietary Assessment Based on Food Images: Scoping Review.” Journal of Medical Internet Research. 2024. Open source

[6] Collins GS, Moons KGM, Dhiman P, et al. “TRIPOD+AI statement: updated reporting guidance for clinical prediction models that use regression or machine learning.” BMJ. 2024;385:e078378. DOI: 10.1136/bmj-2023-078378. Open source

[7] World Health Organization. “Recommendations on Digital Interventions for Health System Strengthening.” WHO Guideline. 2019. ISBN 978-92-4-155050-5. Open source

[8] Directorate General of Drug Administration, Bangladesh. Medical-device registration guidance (guideline dated 15 February 2024) and notice confirming approval of the “Guideline on the Regulation of Artificial Intelligence and Medical Robotics in Healthcare, Bangladesh” on 22 April 2026. The current requirements for this product must be confirmed with DGDA before deployment. Medical-device guideline | 2026 AI-in-health notice

Appendix: Immediate Team Checklist

Before making clinical claims or submitting the final pitch, the team should close these basic tasks:

- Confirm the first target user and the exact problem with at least a small set of patients and clinicians.
- Choose one measurable outcome for the first R&D comparison and define it with a qualified clinical adviser.
- Confirm dataset access, licensing, relevant fields, and whether each dataset fits the research question.
- Build one complete demo journey: patient measurement entry, trend view, safe message, and doctor summary.
- Show what happens with missing data, an incorrect meal estimate, an unusual reading, or an unavailable LLM.
- Name the proposed alert owner and backup process for any supervised pilot; do not promise 24/7 care without staffing.
- Write down which features are working, which are mock-ups, and which are future research.
- Review every health claim, reference, privacy rule, and regulatory assumption before the presentation.

Document Status

This document is a concept proposal prepared on 12 October 2026. The product scope, research questions, and roadmap are proposed plans. The team has not claimed that a validated predictive model, clinical pilot, regulatory approval, partnership, or commercial traction already exists. Those items should be updated only when evidence is available.