

Thesis Proposal:

# Advances in Algorithms for Matrix Approximation

via **Sampling** and **Sketching**

Taisuke (Tai) Yasuda



**Carnegie Mellon University**

Computer Science Department

April 10, 2023

Thesis Committee:

- David P. Woodruff (Carnegie Mellon University, Chair)
- Anupam Gupta (Carnegie Mellon University)
- Richard Peng (Carnegie Mellon University)
- Cameron Musco (University of Massachusetts Amherst)

# Matrix Approximation

**“Turning big data into tiny data”**

# Matrix Approximation

**“Turning big data into tiny data”**

- **Matrix approximation** is the problem of approximating large matrices by smaller matrices

# Matrix Approximation

**“Turning big data into tiny data”**

- **Matrix approximation** is the problem of approximating large matrices by smaller matrices
- Modern large-scale machine learning problems deal with huge matrices!

# Matrix Approximation

**“Turning big data into tiny data”**

- **Matrix approximation** is the problem of approximating large matrices by smaller matrices
- Modern large-scale machine learning problems deal with huge matrices!
  - Billions of training examples and labels

# Matrix Approximation

**“Turning big data into tiny data”**

- **Matrix approximation** is the problem of approximating large matrices by smaller matrices
- Modern large-scale machine learning problems deal with huge matrices!
  - Billions of training examples and labels
  - Thousands of features

# Matrix Approximation

**“Turning big data into tiny data”**

- **Matrix approximation** is the problem of approximating large matrices by smaller matrices
- Modern large-scale machine learning problems deal with huge matrices!
  - Billions of training examples and labels
  - Thousands of features
- Goal: replace a large dataset with a smaller dataset to **improve efficiency of data analytic tasks**

# Matrix Approximation

## **Randomized Numerical Linear Algebra**



# Matrix Approximation

## Randomized Numerical Linear Algebra

- **Numerical linear algebra** [pre 2000's]: **deterministic** algorithms for solving linear algebra to **machine precision**

# Matrix Approximation

## Randomized Numerical Linear Algebra

- **Numerical linear algebra** [pre 2000's]: **deterministic** algorithms for solving linear algebra to **machine precision**
- **Randomized numerical linear algebra** [2000's]: **randomized approximation** algorithms for numerical linear algebra

# Matrix Approximation

## Randomized Numerical Linear Algebra

- **Numerical linear algebra** [pre 2000's]: **deterministic** algorithms for solving linear algebra to **machine precision**
- **Randomized numerical linear algebra** [2000's]: **randomized approximation** algorithms for numerical linear algebra
  - Randomized: succeed every time → succeed with 99% probability

# Matrix Approximation

## Randomized Numerical Linear Algebra

- **Numerical linear algebra** [pre 2000's]: **deterministic** algorithms for solving linear algebra to **machine precision**
- **Randomized numerical linear algebra** [2000's]: **randomized approximation** algorithms for numerical linear algebra
  - Randomized: succeed every time → succeed with 99% probability
  - Approximation: solve exactly → solve up to (small) error

# Matrix Approximation

## Randomized Numerical Linear Algebra

- **Numerical linear algebra** [pre 2000's]: **deterministic** algorithms for solving linear algebra to **machine precision**
- **Randomized numerical linear algebra** [2000's]: **randomized approximation** algorithms for numerical linear algebra
  - Randomized: succeed every time → succeed with 99% probability
  - Approximation: solve exactly → solve up to (small) error
  - **This flexibility leads to extraordinary improvements in efficiency!**

# Matrix Approximation

## Randomized Numerical Linear Algebra

- **Numerical linear algebra** [pre 2000's]: **deterministic** algorithms for solving linear algebra to **machine precision**
- **Randomized numerical linear algebra** [2000's]: **randomized approximation** algorithms for numerical linear algebra
  - Randomized: succeed every time → succeed with 99% probability
  - Approximation: solve exactly → solve up to (small) error
  - **This flexibility leads to extraordinary improvements in efficiency!**
  - Key techniques: **sampling**, **sketching**, and **optimization**

# Matrix Approximation

## **New Challenges**

# Matrix Approximation

## New Challenges

1. **Tight bounds:** what are the fundamental possibilities and limitations for matrix approximation?



# Matrix Approximation

## New Challenges

1. **Tight bounds:** what are the fundamental possibilities and limitations for matrix approximation?
2. **Uncertain/resource-limited settings:** are there algorithms for matrix approximation using limited information or computational resources?

# Matrix Approximation

## New Challenges

1. **Tight bounds:** what are the fundamental possibilities and limitations for matrix approximation?
2. **Uncertain/resource-limited settings:** are there algorithms for matrix approximation using limited information or computational resources?
3. **Generalized loss functions:** are there algorithms for matrix approximation for generalized objectives and loss functions?

# Matrix Approximation

## New Challenges

1. **Tight bounds:** what are the fundamental possibilities and limitations for matrix approximation?
2. **Uncertain/resource-limited settings:** are there algorithms for matrix approximation using limited information or computational resources?
3. **Generalized loss functions:** are there algorithms for matrix approximation for generalized objectives and loss functions?
4. **Applications:** can techniques for matrix approximation be applied to solve problems in adjacent areas of computer science?

# Subspace Embeddings and Linear Regression

# Subspace Embeddings and Linear Regression

- Oblivious  $\ell_p$  subspace embeddings: high distortion and low distortion

# Subspace Embeddings and Linear Regression

- Oblivious  $\ell_p$  subspace embeddings: high distortion and low distortion
- Non-oblivious subspace embeddings:  $\ell_p$  Lewis weight sampling, general losses

# Subspace Embeddings and Linear Regression

- Oblivious  $\ell_p$  subspace embeddings: high distortion and low distortion
- Non-oblivious subspace embeddings:  $\ell_p$  Lewis weight sampling, general losses
- Applications: active learning, streaming computational geometry, low rank approximation

# Subspace Embeddings and Linear Regression

Oblivious  $\ell_p$  subspace embeddings: high distortion and low distortion

- Non-oblivious subspace embeddings:  $\ell_p$  Lewis weight sampling, general losses
- Applications: active learning, streaming computational geometry, low rank approximation



# Matrix Approximation

## **Linear Regression**

# Matrix Approximation

## Linear Regression

- **Linear regression:** given matrix  $\mathbf{A} \in \mathbb{R}^{n \times d}$  and vector  $\mathbf{b} \in \mathbb{R}^n$ , solve

# Matrix Approximation

## Linear Regression

- **Linear regression:** given matrix  $\mathbf{A} \in \mathbb{R}^{n \times d}$  and vector  $\mathbf{b} \in \mathbb{R}^n$ , solve

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_2^2$$

# Matrix Approximation

## Linear Regression

- **Linear regression:** given matrix  $\mathbf{A} \in \mathbb{R}^{n \times d}$  and vector  $\mathbf{b} \in \mathbb{R}^n$ , solve

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_2^2$$

- Simple model for supervised learning

# Matrix Approximation

## Linear Regression

- **Linear regression:** given matrix  $\mathbf{A} \in \mathbb{R}^{n \times d}$  and vector  $\mathbf{b} \in \mathbb{R}^n$ , solve

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_2^2$$

- Simple model for supervised learning
- Building block for complex models and algorithms

# Matrix Approximation

## Linear Regression

- **Linear regression:** given matrix  $\mathbf{A} \in \mathbb{R}^{n \times d}$  and vector  $\mathbf{b} \in \mathbb{R}^n$ , solve


$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_2^2$$

- Simple model for supervised learning
- Building block for complex models and algorithms
- Can we design efficient approximation algorithms for linear regression?

# Matrix Approximation

Approximating the Linear Regression Loss Function


$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_2^2$$



# Matrix Approximation

## Approximating the Linear Regression Loss Function

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_2^2$$




- One approach: approximate the original instance by a smaller instance



# Matrix Approximation

## Approximating the Linear Regression Loss Function



$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_2^2$$


- One approach: approximate the original instance by a smaller instance

**Goal:** Replace  $\mathbf{A}$  and  $\mathbf{b}$  by a smaller  $\tilde{\mathbf{A}}$  and  $\tilde{\mathbf{b}}$

$$\text{s.t. } \|\mathbf{Ax} - \mathbf{b}\|_2 \approx \|\tilde{\mathbf{A}}\mathbf{x} - \tilde{\mathbf{b}}\|_2$$

# Matrix Approximation

## Approximating the Linear Regression Loss Function



$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_2^2$$

- One approach: approximate the original instance by a smaller instance


**Goal:** Replace  $\mathbf{A}$  and  $\mathbf{b}$  by a smaller  $\tilde{\mathbf{A}}$  and  $\tilde{\mathbf{b}}$

$$\text{s.t. } \|\mathbf{Ax} - \mathbf{b}\|_2 \approx \|\tilde{\mathbf{A}}\mathbf{x} - \tilde{\mathbf{b}}\|_2$$

- There are many possible ways to choose  $\tilde{\mathbf{A}}$  and  $\tilde{\mathbf{b}}$ !

# Matrix Approximation

## Approximating the Linear Regression Loss Function



$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_2^2$$

- One approach: approximate the original instance by a smaller instance

**Goal:** Replace  $\mathbf{A}$  and  $\mathbf{b}$  by a smaller  $\tilde{\mathbf{A}}$  and  $\tilde{\mathbf{b}}$


$$\text{s.t. } \|\mathbf{Ax} - \mathbf{b}\|_2 \approx \|\tilde{\mathbf{A}}\mathbf{x} - \tilde{\mathbf{b}}\|_2$$

- There are many possible ways to choose  $\tilde{\mathbf{A}}$  and  $\tilde{\mathbf{b}}$ !

**Idea.**  $\tilde{\mathbf{A}} = \mathbf{SA}$  and  $\tilde{\mathbf{b}} = \mathbf{Sb}$  for some  $\mathbf{S} \in \mathbb{R}^{r \times n}$ ,  $r \ll n$

# Matrix Approximation

## Approximating the Linear Regression Loss Function



$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_2^2$$

- One approach: approximate the original instance by a smaller instance

**Goal:** Replace  $\mathbf{A}$  and  $\mathbf{b}$  by a smaller  $\tilde{\mathbf{A}}$  and  $\tilde{\mathbf{b}}$

$$\text{s.t. } \|\mathbf{Ax} - \mathbf{b}\|_2 \approx \|\tilde{\mathbf{A}}\mathbf{x} - \tilde{\mathbf{b}}\|_2$$

- There are many possible ways to choose  $\tilde{\mathbf{A}}$  and  $\tilde{\mathbf{b}}$ !

**Idea.**  $\tilde{\mathbf{A}} = \mathbf{SA}$  and  $\tilde{\mathbf{b}} = \mathbf{Sb}$  for some  $\mathbf{S} \in \mathbb{R}^{r \times n}$ ,  $r \ll n$

“Sketch”

# Matrix Approximation

## Approximating the Linear Regression Loss Function

**Goal:** Replace  $\mathbf{A}$  and  $\mathbf{b}$  by a smaller  $\tilde{\mathbf{A}}$  and  $\tilde{\mathbf{b}}$

$$\text{s.t. } \|\mathbf{Ax} - \mathbf{b}\|_2 \approx \|\tilde{\mathbf{A}}\mathbf{x} - \tilde{\mathbf{b}}\|_2$$

**Idea.**  $\tilde{\mathbf{A}} = \mathbf{SA}$  and  $\tilde{\mathbf{b}} = \mathbf{Sb}$


for some  $\mathbf{S} \in \mathbb{R}^{r \times n}$ ,  $r \ll n$

# Matrix Approximation

## Approximating the Linear Regression Loss Function

**Goal:** Replace  $\mathbf{A}$  and  $\mathbf{b}$  by a smaller  $\tilde{\mathbf{A}}$  and  $\tilde{\mathbf{b}}$   
s.t.  $\| \mathbf{Ax} - \mathbf{b} \|_2 \approx \| \tilde{\mathbf{A}}\mathbf{x} - \tilde{\mathbf{b}} \|_2$

**Idea.**  $\tilde{\mathbf{A}} = \mathbf{SA}$  and  $\tilde{\mathbf{b}} = \mathbf{Sb}$   
for some  $\mathbf{S} \in \mathbb{R}^{r \times n}$ ,  $r \ll n$



$$\mathbf{A} \mathbf{x} - \mathbf{b}$$

# Matrix Approximation

## Approximating the Linear Regression Loss Function

**Goal:** Replace  $\mathbf{A}$  and  $\mathbf{b}$  by a smaller  $\tilde{\mathbf{A}}$  and  $\tilde{\mathbf{b}}$   
s.t.  $\| \mathbf{Ax} - \mathbf{b} \|_2 \approx \| \tilde{\mathbf{A}}\mathbf{x} - \tilde{\mathbf{b}} \|_2$

**Idea.**  $\tilde{\mathbf{A}} = \mathbf{SA}$  and  $\tilde{\mathbf{b}} = \mathbf{Sb}$   
for some  $\mathbf{S} \in \mathbb{R}^{r \times n}$ ,  $r \ll n$




Original instance ( $n$  rows)

# Matrix Approximation

## Approximating the Linear Regression Loss Function

**Goal:** Replace  $\mathbf{A}$  and  $\mathbf{b}$  by a smaller  $\tilde{\mathbf{A}}$  and  $\tilde{\mathbf{b}}$   
s.t.  $\| \mathbf{Ax} - \mathbf{b} \|_2 \approx \| \tilde{\mathbf{A}}\mathbf{x} - \tilde{\mathbf{b}} \|_2$

**Idea.**  $\tilde{\mathbf{A}} = \mathbf{SA}$  and  $\tilde{\mathbf{b}} = \mathbf{Sb}$   
for some  $\mathbf{S} \in \mathbb{R}^{r \times n}$ ,  $r \ll n$






# Matrix Approximation

## Approximating the Linear Regression Loss Function

**Goal:** Replace  $\mathbf{A}$  and  $\mathbf{b}$  by a smaller  $\tilde{\mathbf{A}}$  and  $\tilde{\mathbf{b}}$   
s.t.  $\| \mathbf{Ax} - \mathbf{b} \|_2 \approx \| \tilde{\mathbf{A}}\mathbf{x} - \tilde{\mathbf{b}} \|_2$

**Idea.**  $\tilde{\mathbf{A}} = \mathbf{SA}$  and  $\tilde{\mathbf{b}} = \mathbf{Sb}$   
for some  $\mathbf{S} \in \mathbb{R}^{r \times n}$ ,  $r \ll n$




# Matrix Approximation

## Approximating the Linear Regression Loss Function

**Goal:** Replace  $\mathbf{A}$  and  $\mathbf{b}$  by a smaller  $\tilde{\mathbf{A}}$  and  $\tilde{\mathbf{b}}$   
s.t.  $\| \mathbf{Ax} - \mathbf{b} \|_2 \approx \| \tilde{\mathbf{A}}\mathbf{x} - \tilde{\mathbf{b}} \|_2$

**Idea.**  $\tilde{\mathbf{A}} = \mathbf{SA}$  and  $\tilde{\mathbf{b}} = \mathbf{Sb}$   
for some  $\mathbf{S} \in \mathbb{R}^{r \times n}$ ,  $r \ll n$



# Matrix Approximation

## Approximating the Linear Regression Loss Function

**Definition (Sarlos 2006).**  $S \in \mathbb{R}^{r \times n}$  is a **subspace embedding** of  $A \in \mathbb{R}^{n \times d}$  if

$$\|A\mathbf{x}\|_2 \leq \|S A \mathbf{x}\|_2 \leq \kappa \|A\mathbf{x}\|_2$$

for every  $\mathbf{x} \in \mathbb{R}^d$ .

# Matrix Approximation

## Approximating the Linear Regression Loss Function

**Definition (Sarlos 2006).**  $S \in \mathbb{R}^{r \times n}$  is a **subspace embedding** of  $A \in \mathbb{R}^{n \times d}$  if

$$\|A\mathbf{x}\|_2 \leq \|S A \mathbf{x}\|_2 \leq \kappa \|A\mathbf{x}\|_2$$

for every  $\mathbf{x} \in \mathbb{R}^d$ .

Distortion/error

# Matrix Approximation

## Approximating the Linear Regression Loss Function

**Definition (Sarlos 2006).**  $S \in \mathbb{R}^{r \times n}$  is a **subspace embedding** of  $A \in \mathbb{R}^{n \times d}$  if

$$\|A\mathbf{x}\|_2 \leq \|S A \mathbf{x}\|_2 \leq \kappa \|A\mathbf{x}\|_2$$

for every  $\mathbf{x} \in \mathbb{R}^d$ .

Distortion/error

Why is this useful?

# Matrix Approximation

## Approximating the Linear Regression Loss Function

**Definition (Sarlos 2006).**  $S \in \mathbb{R}^{r \times n}$  is a **subspace embedding** of  $A \in \mathbb{R}^{n \times d}$  if

$$\|A\mathbf{x}\|_2 \leq \|S A \mathbf{x}\|_2 \leq \kappa \|A\mathbf{x}\|_2$$

for every  $\mathbf{x} \in \mathbb{R}^d$ .

Distortion/error

Why is this useful?

- Let  $A' = [A \ \mathbf{b}] \in \mathbb{R}^{n \times (d+1)}$  and let  $S \in \mathbb{R}^{r \times n}$  be a subspace embedding of  $A'$

# Matrix Approximation

## Approximating the Linear Regression Loss Function

**Definition (Sarlos 2006).**  $S \in \mathbb{R}^{r \times n}$  is a **subspace embedding** of  $A \in \mathbb{R}^{n \times d}$  if

$$\|A\mathbf{x}\|_2 \leq \|S A \mathbf{x}\|_2 \leq \kappa \|A\mathbf{x}\|_2$$

for every  $\mathbf{x} \in \mathbb{R}^d$ .

Distortion/error

Why is this useful?

- Let  $A' = [A \ \mathbf{b}] \in \mathbb{R}^{n \times (d+1)}$  and let  $S \in \mathbb{R}^{r \times n}$  be a subspace embedding of  $A'$
- $A\mathbf{x} - \mathbf{b} = A'\mathbf{x}'$  for  $\mathbf{x}' = [\mathbf{x}; -1] \in \mathbb{R}^{d+1}$

# Matrix Approximation

## Approximating the Linear Regression Loss Function

**Definition (Sarlos 2006).**  $S \in \mathbb{R}^{r \times n}$  is a **subspace embedding** of  $A \in \mathbb{R}^{n \times d}$  if

$$\|A\mathbf{x}\|_2 \leq \|S A \mathbf{x}\|_2 \leq \kappa \|A\mathbf{x}\|_2$$

for every  $\mathbf{x} \in \mathbb{R}^d$ .

Distortion/error

Why is this useful?

- Let  $A' = [A \ \mathbf{b}] \in \mathbb{R}^{n \times (d+1)}$  and let  $S \in \mathbb{R}^{r \times n}$  be a subspace embedding of  $A'$
- $A\mathbf{x} - \mathbf{b} = A'\mathbf{x}'$  for  $\mathbf{x}' = [\mathbf{x}; -1] \in \mathbb{R}^{d+1}$
- $\|A\mathbf{x} - \mathbf{b}\|_2 \leq \|S A \mathbf{x} - S \mathbf{b}\|_2 \leq \kappa \|A\mathbf{x} - \mathbf{b}\|_2$



# Matrix Approximation

## Approximating the Linear Regression Loss Function

**Definition (Sarlos 2006).**  $S \in \mathbb{R}^{r \times n}$  is a **subspace embedding** of  $A \in \mathbb{R}^{n \times d}$  if

$$\|A\mathbf{x}\|_2 \leq \|S A \mathbf{x}\|_2 \leq \kappa \|A\mathbf{x}\|_2$$

for every  $\mathbf{x} \in \mathbb{R}^d$ .

Distortion/error

Why is this useful?

- Let  $A' = [A \ \mathbf{b}] \in \mathbb{R}^{n \times (d+1)}$  and let  $S \in \mathbb{R}^{r \times n}$  be a subspace embedding of  $A'$
- $A\mathbf{x} - \mathbf{b} = A'\mathbf{x}'$  for  $\mathbf{x}' = [\mathbf{x}; -1] \in \mathbb{R}^{d+1}$
- $\|A\mathbf{x} - \mathbf{b}\|_2 \leq \|S A \mathbf{x} - S \mathbf{b}\|_2 \leq \kappa \|A\mathbf{x} - \mathbf{b}\|_2$

Cost of  $SA$  and  $S\mathbf{b}$  approximates cost of  $A$  and  $\mathbf{b}$

# Matrix Approximation

## Approximating the Linear Regression Loss Function

**Definition (Sarlos 2006).**  $S \in \mathbb{R}^{r \times n}$  is a **subspace embedding** of  $A \in \mathbb{R}^{n \times d}$  if

$$\|A\mathbf{x}\|_2 \leq \|S A \mathbf{x}\|_2 \leq \kappa \|A\mathbf{x}\|_2$$

for every  $\mathbf{x} \in \mathbb{R}^d$ .

Distortion/error


Why is this useful?

- Let  $A' = [A \ \mathbf{b}] \in \mathbb{R}^{n \times (d+1)}$  and let  $S \in \mathbb{R}^{r \times n}$  be a subspace embedding of  $A'$
- $A\mathbf{x} - \mathbf{b} = A'\mathbf{x}'$  for  $\mathbf{x}' = [\mathbf{x}; -1] \in \mathbb{R}^{d+1}$
- $\|A\mathbf{x} - \mathbf{b}\|_2 \leq \|S A \mathbf{x} - S \mathbf{b}\|_2 \leq \kappa \|A\mathbf{x} - \mathbf{b}\|_2$  Cost of  $SA$  and  $S\mathbf{b}$  approximates cost of  $A$  and  $\mathbf{b}$
- Solve linear regression on  $r \times d$  matrix  $SA$  instead of  $n \times d$  matrix  $A$ !

# Matrix Approximation

## Oblivious Subspace Embeddings


**Theorem (Sarlos 2006).** Let  $\kappa = (1 + \varepsilon)$ . Let  $r = \tilde{O}(\varepsilon^{-2}d)$ . If  $\mathbf{S}$  is an  $r \times n$  Gaussian matrix, then  $\mathbf{S}$  is a subspace embedding for any  $\mathbf{A}$  with distortion  $\kappa$ , with probability 99%.



# Matrix Approximation

## Oblivious Subspace Embeddings


**Theorem (Sarlos 2006).** Let  $\kappa = (1 + \varepsilon)$ . Let  $r = \tilde{O}(\varepsilon^{-2}d)$ . If  $\mathbf{S}$  is an  $r \times n$  Gaussian matrix, then  $\mathbf{S}$  is a subspace embedding for any  $\mathbf{A}$  with distortion  $\kappa$ , with probability 99%.



# Matrix Approximation

## Oblivious Subspace Embeddings


**Theorem (Sarlos 2006).** Let  $\kappa = (1 + \varepsilon)$ . Let  $r = \tilde{O}(\varepsilon^{-2}d)$ . If  $\mathbf{S}$  is an  $r \times n$  Gaussian matrix, then  $\mathbf{S}$  is a subspace embedding for any  $\mathbf{A}$  with distortion  $\kappa$ , with probability 99%.



# Matrix Approximation

## Oblivious Subspace Embeddings

**Theorem (Sarlos 2006).** Let  $\kappa = (1 + \varepsilon)$ . Let  $r = \tilde{O}(\varepsilon^{-2}d)$ . If  $\mathbf{S}$  is an  $r \times n$  Gaussian matrix, then  $\mathbf{S}$  is a subspace embedding for any  $\mathbf{A}$  with distortion  $\kappa$ , with probability 99%.



So  $\ell_2$  regression is resolved.  
What's next?

# Matrix Approximation

$\ell_p$  Linear Regression

# Matrix Approximation

$\ell_p$  Linear Regression

$\ell_2$  linear regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_2^2$$

- Minimize the **sum of squares** of errors



# Matrix Approximation

## $\ell_p$ Linear Regression

$\ell_1$  linear regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_1$$

- Minimize the **average** error
- **Robust** loss function

$\ell_2$  linear regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_2^2$$

- Minimize the **sum of squares** of errors

# Matrix Approximation

## $\ell_p$ Linear Regression

$\ell_1$  linear regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_1$$

- Minimize the **average** error
- **Robust** loss function

$\ell_2$  linear regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_2^2$$

- Minimize the **sum of squares** of errors

$\ell_\infty$  linear regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_\infty$$

- Minimize the **worst-case** error
- **Sensitive** loss function

# Matrix Approximation

## $\ell_p$ Linear Regression

$\ell_1$  linear regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_1$$

- Minimize the **average** error
- **Robust** loss function

$\ell_2$  linear regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_2^2$$

- Minimize the **sum of squares** of errors



$\ell_\infty$  linear regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_\infty$$

- Minimize the **worst-case** error
- **Sensitive** loss function

$\ell_p$  linear regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_p^p$$



# Matrix Approximation

## $\ell_p$ Linear Regression

$\ell_1$  linear regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_1$$

- Minimize the **average** error
- **Robust** loss function

$\ell_2$  linear regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_2^2$$

- Minimize the **sum of squares** of errors

$\ell_\infty$  linear regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_\infty$$

- Minimize the **worst-case** error
- **Sensitive** loss function

$\ell_p$  linear regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_p^p$$


**Question.** What trade-offs are possible for oblivious subspace embeddings under the  $\ell_p$  loss?

# Matrix Approximation

**Oblivious  $\ell_p$  Subspace Embeddings**


# Matrix Approximation

Oblivious  $\ell_p$  Subspace Embeddings



# Matrix Approximation


## Oblivious $\ell_p$ Subspace Embeddings



Oblivious = does not depend on  $A$

# Matrix Approximation


## Oblivious $\ell_p$ Subspace Embeddings





# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings




$$\|\mathbf{Ax}\|_p \leq \|\mathbf{SAx}\|_p \leq \kappa \|\mathbf{Ax}\|_p$$

for every  $\mathbf{x} \in \mathbb{R}^d$

# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

Two regimes for  $1 \leq p < 2$



Oblivious = does not depend on  $\mathbf{A}$


$$\|\mathbf{Ax}\|_p \leq \|\mathbf{SAx}\|_p \leq \kappa \|\mathbf{Ax}\|_p$$

for every  $\mathbf{x} \in \mathbb{R}^d$

# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

Two regimes for  $1 \leq p < 2$




$$\|\mathbf{Ax}\|_p \leq \|\mathbf{SAx}\|_p \leq \kappa \|\mathbf{Ax}\|_p$$

for every  $\mathbf{x} \in \mathbb{R}^d$

# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

Two regimes for  $1 \leq p < 2$



$$\|\mathbf{Ax}\|_p \leq \|\mathbf{SAx}\|_p \leq \kappa \|\mathbf{Ax}\|_p$$


for every  $\mathbf{x} \in \mathbb{R}^d$

# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

Two regimes for  $1 \leq p < 2$

- **High distortion:** sketch  $\mathbf{S}$  has...



Oblivious = does not depend on  $\mathbf{A}$

$$\|\mathbf{Ax}\|_p \leq \|\mathbf{SAx}\|_p \leq \kappa \|\mathbf{Ax}\|_p$$

for every  $\mathbf{x} \in \mathbb{R}^d$


# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

Two regimes for  $1 \leq p < 2$

- **High distortion:** sketch  $\mathbf{S}$  has...

- **Low distortion:** sketch  $\mathbf{S}$  has...



Oblivious = does not depend on  $\mathbf{A}$

$$\|\mathbf{Ax}\|_p \leq \|\mathbf{SAx}\|_p \leq \kappa \|\mathbf{Ax}\|_p$$

for every  $\mathbf{x} \in \mathbb{R}^d$

# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings


Two regimes for  $1 \leq p < 2$

- **High distortion:** sketch  $\mathbf{S}$  has...

- $r = \text{poly}(d)$  rows

- $\kappa = \text{poly}(d)$  distortion

- **Low distortion:** sketch  $\mathbf{S}$  has...



$$\|\mathbf{Ax}\|_p \leq \|\mathbf{SAx}\|_p \leq \kappa \|\mathbf{Ax}\|_p$$

for every  $\mathbf{x} \in \mathbb{R}^d$

# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

Two regimes for  $1 \leq p < 2$

- **High distortion:** sketch  $\mathbf{S}$  has...


- $r = \text{poly}(d)$  rows

- $\kappa = \text{poly}(d)$  distortion

- **Low distortion:** sketch  $\mathbf{S}$  has...

- $r \gg \text{poly}(d)$  rows

- $\kappa = (1 + \varepsilon)$  distortion



$$\|\mathbf{Ax}\|_p \leq \|\mathbf{SAx}\|_p \leq \kappa \|\mathbf{Ax}\|_p$$

for every  $\mathbf{x} \in \mathbb{R}^d$



# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

Two regimes for  $1 \leq p < 2$

- **High distortion:** sketch  $\mathbf{S}$  has...

- $r = \text{poly}(d)$  rows

- $\kappa = \text{poly}(d)$  distortion

- **Low distortion:** sketch  $\mathbf{S}$  has...

- $r \gg \text{poly}(d)$  rows

- $\kappa = (1 + \varepsilon)$  distortion

# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

Two regimes for  $1 \leq p < 2$

- **High distortion:** sketch  $\mathbf{S}$  has...

- $r = \text{poly}(d)$  rows

- $\kappa = \text{poly}(d)$  distortion

- **Low distortion:** sketch  $\mathbf{S}$  has...

- $r \gg \text{poly}(d)$  rows

- $\kappa = (1 + \varepsilon)$  distortion

**Best of both worlds is not possible! (Wang—Woodruff 2019)**

# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

Two regimes for  $1 \leq p < 2$

- **High distortion:** sketch  $\mathbf{S}$  has...

- $r = \text{poly}(d)$  rows

- $\kappa = \text{poly}(d)$  distortion

- **Low distortion:** sketch  $\mathbf{S}$  has...

- $r \gg \text{poly}(d)$  rows

- $\kappa = (1 + \varepsilon)$  distortion

**Best of both worlds is not possible! (Wang—Woodruff 2019)**

- **High distortion:** sketch  $\mathbf{S}$  has...

- $r = \text{poly}(d) \Rightarrow \kappa \gtrsim d^{1/p}$

# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

Two regimes for  $1 \leq p < 2$

- **High distortion:** sketch  $\mathbf{S}$  has...
  - $r = \text{poly}(d)$  rows
  - $\kappa = \text{poly}(d)$  distortion

- **Low distortion:** sketch  $\mathbf{S}$  has...
  - $r \gg \text{poly}(d)$  rows
  - $\kappa = (1 + \varepsilon)$  distortion

**Best of both worlds is not possible! (Wang—Woodruff 2019)**

- **High distortion:** sketch  $\mathbf{S}$  has...
  - $r = \text{poly}(d) \Rightarrow \kappa \gtrsim d^{1/p}$

- **Low distortion:** sketch  $\mathbf{S}$  has...
  - $\kappa = O(1)$  distortion  $\Rightarrow r \gtrsim \exp(\sqrt{d})$

# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

**Best of both worlds is not possible! (Wang—Woodruff 2019)**

- **High distortion:** sketch  $\mathbf{S}$  has...

- $r = \text{poly}(d) \Rightarrow \kappa \gtrsim d^{1/p}$

- **Low distortion:** sketch  $\mathbf{S}$  has...

- $\kappa = O(1) \text{ distortion} \Rightarrow r \gtrsim \exp(\sqrt{d})$

# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

**Best of both worlds is not possible! (Wang—Woodruff 2019)**

- |   |   |
|---|---|
| <ul style="list-style-type: none"><li>• <b>High distortion:</b> sketch <math>\mathbf{S}</math> has...<ul style="list-style-type: none"><li>– <math>r = \text{poly}(d) \Rightarrow \kappa \gtrsim d^{1/p}</math></li></ul></li></ul> | <ul style="list-style-type: none"><li>• <b>Low distortion:</b> sketch <math>\mathbf{S}</math> has...<ul style="list-style-type: none"><li>– <math>\kappa = O(1)</math> distortion <math>\Rightarrow r \gtrsim \exp(\sqrt{d})</math></li></ul></li></ul> |
|---|---|

**Question.** Are the lower bounds of Wang—Woodruff 2019 tight?

# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

Best of both worlds is not possible! (Wang—Woodruff 2019)

- |   |   |
|---|---|
| <ul style="list-style-type: none"><li>• <b>High distortion:</b> sketch <math>\mathbf{S}</math> has...<ul style="list-style-type: none"><li>– <math>r = \text{poly}(d) \Rightarrow \kappa \gtrsim d^{1/p}</math></li></ul></li></ul> | <ul style="list-style-type: none"><li>• <b>Low distortion:</b> sketch <math>\mathbf{S}</math> has...<ul style="list-style-type: none"><li>– <math>\kappa = O(1)</math> distortion <math>\Rightarrow r \gtrsim \exp(\sqrt{d})</math></li></ul></li></ul> |
|---|---|

**Question.** Are the lower bounds of Wang—Woodruff 2019 tight?

Yes!

- **Woodruff—Y 2023**
  - $r = \tilde{O}(d), \kappa = \tilde{O}(d^{1/p})$

# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

**Best of both worlds is not possible! (Wang—Woodruff 2019)**

- |   |   |
|---|---|
| <ul style="list-style-type: none"><li>• <b>High distortion:</b> sketch <math>\mathbf{S}</math> has...<ul style="list-style-type: none"><li>– <math>r = \text{poly}(d) \Rightarrow \kappa \gtrsim d^{1/p}</math></li></ul></li></ul> | <ul style="list-style-type: none"><li>• <b>Low distortion:</b> sketch <math>\mathbf{S}</math> has...<ul style="list-style-type: none"><li>– <math>\kappa = O(1)</math> distortion <math>\Rightarrow r \gtrsim \exp(\sqrt{d})</math></li></ul></li></ul> |
|---|---|

**Question.** Are the lower bounds of Wang—Woodruff 2019 tight?

**Yes!**

- **Woodruff—Y 2023**
  - $r = \tilde{O}(d), \kappa = \tilde{O}(d^{1/p})$
- Prior best: Wang—Woodruff 2019
  - $r = \tilde{O}(d), \kappa = \tilde{O}(d)$



# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

**Best of both worlds is not possible! (Wang—Woodruff 2019)**

- |   |   |
|---|---|
| <ul style="list-style-type: none"><li>• <b>High distortion:</b> sketch <math>\mathbf{S}</math> has...<ul style="list-style-type: none"><li>– <math>r = \text{poly}(d) \Rightarrow \kappa \gtrsim d^{1/p}</math></li></ul></li></ul> | <ul style="list-style-type: none"><li>• <b>Low distortion:</b> sketch <math>\mathbf{S}</math> has...<ul style="list-style-type: none"><li>– <math>\kappa = O(1)</math> distortion <math>\Rightarrow r \gtrsim \exp(\sqrt{d})</math></li></ul></li></ul> |
|---|---|

**Question.** Are the lower bounds of Wang—Woodruff 2019 tight?

**Yes!**

- **Woodruff—Y 2023**
  - $r = \tilde{O}(d), \kappa = \tilde{O}(d^{1/p})$
- Prior best: Wang—Woodruff 2019
  - $r = \tilde{O}(d), \kappa = \tilde{O}(d)$

**Almost!**

# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

**Best of both worlds is not possible! (Wang—Woodruff 2019)**

- |   |   |
|---|---|
| <ul style="list-style-type: none"><li>• <b>High distortion:</b> sketch <math>\mathbf{S}</math> has...<ul style="list-style-type: none"><li>– <math>r = \text{poly}(d) \Rightarrow \kappa \gtrsim d^{1/p}</math></li></ul></li></ul> | <ul style="list-style-type: none"><li>• <b>Low distortion:</b> sketch <math>\mathbf{S}</math> has...<ul style="list-style-type: none"><li>– <math>\kappa = O(1)</math> distortion <math>\Rightarrow r \gtrsim \exp(\sqrt{d})</math></li></ul></li></ul> |
|---|---|

**Question.** Are the lower bounds of Wang—Woodruff 2019 tight?

**Yes!**

- **Woodruff—Y 2023**
  - $r = \tilde{O}(d), \kappa = \tilde{O}(d^{1/p})$
- Prior best: Wang—Woodruff 2019
  - $r = \tilde{O}(d), \kappa = \tilde{O}(d)$

**Almost!**

- **Li—Woodruff—Y 2021**
  - $r = \exp(\varepsilon^{-1}d), \kappa = (1 + \varepsilon)$

# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

**Best of both worlds is not possible! (Wang—Woodruff 2019)**

- |   |   |
|---|---|
| <ul style="list-style-type: none"><li>• <b>High distortion:</b> sketch <math>\mathbf{S}</math> has...<ul style="list-style-type: none"><li>– <math>r = \text{poly}(d) \Rightarrow \kappa \gtrsim d^{1/p}</math></li></ul></li></ul> | <ul style="list-style-type: none"><li>• <b>Low distortion:</b> sketch <math>\mathbf{S}</math> has...<ul style="list-style-type: none"><li>– <math>\kappa = O(1)</math> distortion <math>\Rightarrow r \gtrsim \exp(\sqrt{d})</math></li></ul></li></ul> |
|---|---|

**Question.** Are the lower bounds of Wang—Woodruff 2019 tight?

**Yes!**

- **Woodruff—Y 2023**
  - $r = \tilde{O}(d), \kappa = \tilde{O}(d^{1/p})$
- Prior best: Wang—Woodruff 2019
  - $r = \tilde{O}(d), \kappa = \tilde{O}(d)$

**Almost!**

- **Li—Woodruff—Y 2021**
  - $r = \exp(\varepsilon^{-1}d), \kappa = (1 + \varepsilon)$
- Prior best: Wang—Woodruff 2019
  - $r = \exp(\exp(\varepsilon^{-2}d)), \kappa = (1 + \varepsilon)$

# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

**Best of both worlds is not possible! (Wang—Woodruff 2019)**

- |   |   |
|---|---|
| <ul style="list-style-type: none"><li>• <b>High distortion:</b> sketch <math>\mathbf{S}</math> has...<ul style="list-style-type: none"><li>– <math>r = \text{poly}(d) \Rightarrow \kappa \gtrsim d^{1/p}</math></li></ul></li></ul> | <ul style="list-style-type: none"><li>• <b>Low distortion:</b> sketch <math>\mathbf{S}</math> has...<ul style="list-style-type: none"><li>– <math>\kappa = O(1)</math> distortion <math>\Rightarrow r \gtrsim \exp(\sqrt{d})</math></li></ul></li></ul> |
|---|---|

**Question.** Are the lower bounds of Wang—Woodruff 2019 tight?

**Yes!**

- **Woodruff—Y 2023**

- $r = \tilde{O}(d), \kappa = \tilde{O}(d^{1/p})$

- Prior best: Wang—Woodruff 2019

- $r = \tilde{O}(d), \kappa = \tilde{O}(d)$

**Almost!**

- **Li—Woodruff—Y 2021**


- $r = \exp(\varepsilon^{-1}d), \kappa = (1 + \varepsilon)$

- Prior best: Wang—Woodruff 2019

- $r = \exp(\exp(\varepsilon^{-2}d)), \kappa = (1 + \varepsilon)$

# Matrix Approximation


Oblivious  $\ell_p$  Subspace Embeddings



# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

**Fact.** Oblivious  $\ell_p$  subspace embeddings reduce to constructing **well-conditioned bases** for subspaces




# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

**Fact.** Oblivious  $\ell_p$  subspace embeddings reduce to constructing **well-conditioned bases** for subspaces

$\approx$  orthonormal bases for  $\ell_p$  norms




# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

**Fact.** Oblivious  $\ell_p$  subspace embeddings reduce to constructing **well-conditioned bases** for subspaces

$\approx$  orthonormal bases for  $\ell_p$  norms

- Let  $\mathbf{U}$  be an orthonormal basis for  $\mathbf{A}$






# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

**Fact.** Oblivious  $\ell_p$  subspace embeddings reduce to constructing **well-conditioned bases** for subspaces

$\approx$  orthonormal bases for  $\ell_p$  norms

- Let  $\mathbf{U}$  be an orthonormal basis for  $\mathbf{A}$ 
  - $\|\mathbf{U}\|_F \leq d^{1/2}$  (with equality)




# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

**Fact.** Oblivious  $\ell_p$  subspace embeddings reduce to constructing **well-conditioned bases** for subspaces

$\approx$  orthonormal bases for  $\ell_p$  norms

- Let  $\mathbf{U}$  be an orthonormal basis for  $\mathbf{A}$ 
  - $\|\mathbf{U}\|_F \leq d^{1/2}$  (with equality)
  - $\|\mathbf{x}\|_2 \leq \|\mathbf{U}\mathbf{x}\|_2$  for every  $\mathbf{x} \in \mathbb{R}^d$  (with equality)




# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

**Fact.** Oblivious  $\ell_p$  subspace embeddings reduce to constructing **well-conditioned bases** for subspaces

$\approx$  orthonormal bases for  $\ell_p$  norms

- Let  $\mathbf{U}$  be an ~~orthonormal basis~~ for  $\mathbf{A}$   
**well-conditioned basis**
  - $\|\mathbf{U}\|_F \leq d^{1/2}$  (with equality)
  - $\|\mathbf{x}\|_2 \leq \|\mathbf{U}\mathbf{x}\|_2$  for every  $\mathbf{x} \in \mathbb{R}^d$  (with equality)




# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

**Fact.** Oblivious  $\ell_p$  subspace embeddings reduce to constructing **well-conditioned bases** for subspaces

$\approx$  orthonormal bases for  $\ell_p$  norms

- Let  $\mathbf{U}$  be an ~~orthonormal basis~~ for  $\mathbf{A}$   
**well-conditioned basis**
  - ~~$\|\mathbf{U}\|_F \leq d^{1/2}$  (with equality)~~
  - $\|\mathbf{U}\|_{p,p} \leq \alpha$  entrywise  $\ell_p$  norm
  - $\|\mathbf{x}\|_2 \leq \|\mathbf{U}\mathbf{x}\|_2$  for every  $\mathbf{x} \in \mathbb{R}^d$  (with equality)




# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

**Fact.** Oblivious  $\ell_p$  subspace embeddings reduce to constructing **well-conditioned bases** for subspaces

$\approx$  orthonormal bases for  $\ell_p$  norms

- Let  $\mathbf{U}$  be an ~~orthonormal basis~~ for  $\mathbf{A}$   
**well-conditioned basis**
  - ~~$\|\mathbf{U}\|_F \leq d^{1/2}$  (with equality)~~
  - $\|\mathbf{U}\|_{p,p} \leq \alpha$  entrywise  $\ell_p$  norm
  - ~~$\|\mathbf{x}\|_2 \leq \|\mathbf{U}\mathbf{x}\|_2$  for every  $\mathbf{x} \in \mathbb{R}^d$  (with equality)~~
  - $\|\mathbf{x}\|_q \leq \|\mathbf{U}\mathbf{x}\|_p$  for every  $\mathbf{x} \in \mathbb{R}^d$



# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings


**Fact.** Oblivious  $\ell_p$  subspace embeddings reduce to constructing **well-conditioned bases** for subspaces

$\approx$  orthonormal bases for  $\ell_p$  norms

- Let  $\mathbf{U}$  be an ~~orthonormal basis~~ for  $\mathbf{A}$   
**well-conditioned basis**
  - ~~$\|\mathbf{U}\|_F \leq d^{1/2}$  (with equality)~~
  - $\|\mathbf{U}\|_{p,p} \leq \alpha$  entrywise  $\ell_p$  norm
  - ~~$\|\mathbf{x}\|_2 \leq \|\mathbf{U}\mathbf{x}\|_2$  for every  $\mathbf{x} \in \mathbb{R}^d$  (with equality)~~

$$\|\mathbf{x}\|_q \leq \|\mathbf{U}\mathbf{x}\|_p \text{ for every } \mathbf{x} \in \mathbb{R}^d$$

Hölder conjugate,  $\frac{1}{p} + \frac{1}{q} = 1$




# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

**Fact.** Oblivious  $\ell_p$  subspace embeddings reduce to constructing **well-conditioned bases** for subspaces

$\approx$  orthonormal bases for  $\ell_p$  norms



- Let  $\mathbf{U}$  be an ~~orthonormal basis~~ for  $\mathbf{A}$   
~~well-conditioned basis~~
  - ~~$\|\mathbf{U}\|_F \leq d^{1/2}$  (with equality)~~
  - $\|\mathbf{U}\|_{p,p} \leq \alpha$  entrywise  $\ell_p$  norm
  - ~~$\|\mathbf{x}\|_2 \leq \|\mathbf{U}\mathbf{x}\|_2$  for every  $\mathbf{x} \in \mathbb{R}^d$  (with equality)~~

$$\|\mathbf{x}\|_q \leq \|\mathbf{U}\mathbf{x}\|_p \text{ for every } \mathbf{x} \in \mathbb{R}^d$$

Hölder conjugate,  $\frac{1}{p} + \frac{1}{q} = 1$

**Theorem (Auerbach 1930).** For any  $\mathbf{A}$ , there is  $\mathbf{U}$  with  $\alpha = d$ .




# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

**Fact.** Oblivious  $\ell_p$  subspace embeddings reduce to constructing **well-conditioned bases** for subspaces

$\approx$  orthonormal bases for  $\ell_p$  norms



- Let  $\mathbf{U}$  be an ~~orthonormal basis~~ for  $\mathbf{A}$   
~~well-conditioned basis~~
  - ~~$\|\mathbf{U}\|_F \leq d^{1/2}$  (with equality)~~
  - $\|\mathbf{U}\|_{p,p} \leq \alpha$  entrywise  $\ell_p$  norm
  - ~~$\|\mathbf{x}\|_2 \leq \|\mathbf{U}\mathbf{x}\|_2$  for every  $\mathbf{x} \in \mathbb{R}^d$  (with equality)~~

$$\|\mathbf{x}\|_q \leq \|\mathbf{U}\mathbf{x}\|_p \text{ for every } \mathbf{x} \in \mathbb{R}^d$$

Hölder conjugate,  $\frac{1}{p} + \frac{1}{q} = 1$

**Theorem (Auerbach 1930).** For any  $\mathbf{A}$ , there is  $\mathbf{U}$  with  $\alpha = d$ .

**Conjecture.** For any  $\mathbf{A}$ , there is  $\mathbf{U}$  with  $\alpha = d^{1/p}$ .




# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

**Fact.** Oblivious  $\ell_p$  subspace embeddings reduce to constructing **well-conditioned bases** for subspaces

$\approx$  orthonormal bases for  $\ell_p$  norms



- Let  $\mathbf{U}$  be an ~~orthonormal basis~~ for  $\mathbf{A}$   
~~well-conditioned basis~~
  - ~~$\|\mathbf{U}\|_F \leq d^{1/2}$  (with equality)~~
  - $\|\mathbf{U}\|_{p,p} \leq \alpha$  entrywise  $\ell_p$  norm
  - ~~$\|\mathbf{x}\|_2 \leq \|\mathbf{U}\mathbf{x}\|_2$  for every  $\mathbf{x} \in \mathbb{R}^d$  (with equality)~~

$$\|\mathbf{x}\|_q \leq \|\mathbf{U}\mathbf{x}\|_p \text{ for every } \mathbf{x} \in \mathbb{R}^d$$

Hölder conjugate,  $\frac{1}{p} + \frac{1}{q} = 1$

**Theorem (Auerbach 1930).** For any  $\mathbf{A}$ , there is  $\mathbf{U}$  with  $\alpha = d$ .

**Conjecture.** For any  $\mathbf{A}$ , there is  $\mathbf{U}$  with  $\alpha = d^{1/p}$ .


No success in showing this conjecture! 😞

# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

**Fact.** Oblivious  $\ell_p$  subspace embeddings reduce to constructing **well-conditioned bases** for subspaces

$\approx$  orthonormal bases for  $\ell_p$  norms



**Theorem (Auerbach 1930).** For any  $A$ , there is  $U$  with  $\alpha = d$ .

**Conjecture.** For any  $A$ , there is  $U$  with  $\alpha = d^{1/p}$ .

No success in showing this conjecture! 😞


# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

**Fact.** Oblivious  $\ell_p$  subspace embeddings reduce to constructing **well-conditioned bases** for subspaces

$\approx$  orthonormal bases for  $\ell_p$  norms

- **Idea.** What if we relax well-conditioned **bases** to well-conditioned **spanning sets**?



**Theorem (Auerbach 1930).** For any  $A$ , there is  $U$  with  $\alpha = d$ .

**Conjecture.** For any  $A$ , there is  $U$  with  $\alpha = d^{1/p}$ .

No success in showing this conjecture! 😞

# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings


**Fact.** Oblivious  $\ell_p$  subspace embeddings reduce to constructing **well-conditioned bases** for subspaces

$\approx$  orthonormal bases for  $\ell_p$  norms

- **Idea.** What if we relax well-conditioned **bases** to well-conditioned **spanning sets**?

**Theorem (Woodruff—Y 2023).** For any  $\mathbf{A}$ , there is  $\mathbf{U} \in \mathbb{R}^{n \times s}$  for  $s = O(d \log \log d)$  such that

- $\|\mathbf{U}\|_{p,p} \leq s^{1/p}$
- for every  $\mathbf{z} \in \mathbb{R}^d$ , there is  $\mathbf{x} \in \mathbb{R}^s$  such that  $\mathbf{Az} = \mathbf{Ux}$  and  $\|\mathbf{x}\|_2 \leq \|\mathbf{Ux}\|_p$



**Theorem (Auerbach 1930).** For any  $\mathbf{A}$ , there is  $\mathbf{U}$  with  $\alpha = d$ .

**Conjecture.** For any  $\mathbf{A}$ , there is  $\mathbf{U}$  with  $\alpha = d^{1/p}$ .

No success in showing this conjecture! 😞

# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings


**Fact.** Oblivious  $\ell_p$  subspace embeddings reduce to constructing **well-conditioned bases** for subspaces

$\approx$  orthonormal bases for  $\ell_p$  norms

- **Idea.** What if we relax well-conditioned **bases** to well-conditioned **spanning sets**?

**Theorem (Woodruff—Y 2023).** For any  $\mathbf{A}$ , there is  $\mathbf{U} \in \mathbb{R}^{n \times s}$  for  $s = O(d \log \log d)$  such that

- $\|\mathbf{U}\|_{p,p} \leq s^{1/p}$
- for every  $\mathbf{z} \in \mathbb{R}^d$ , there is  $\mathbf{x} \in \mathbb{R}^s$  such that  $\mathbf{Az} = \mathbf{Ux}$  and  $\|\mathbf{x}\|_2 \leq \|\mathbf{Ux}\|_p$



**Theorem (Auerbach 1930).** For any  $\mathbf{A}$ , there is  $\mathbf{U}$  with  $\alpha = d$ .

**Conjecture.** For any  $\mathbf{A}$ , there is  $\mathbf{U}$  with  $\alpha = d^{1/p}$ .

No success in showing this conjecture! 😞



# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings


**Fact.** Oblivious  $\ell_p$  subspace embeddings reduce to constructing **well-conditioned bases** for subspaces

$\approx$  orthonormal bases for  $\ell_p$  norms

- **Idea.** What if we relax well-conditioned **bases** to well-conditioned **spanning sets**?

**Theorem (Woodruff—Y 2023).** For any  $\mathbf{A}$ , there is  $\mathbf{U} \in \mathbb{R}^{n \times s}$  for  $s = O(d \log \log d)$  such that

- $\|\mathbf{U}\|_{p,p} \leq s^{1/p}$
- for every  $\mathbf{z} \in \mathbb{R}^d$ , there is  $\mathbf{x} \in \mathbb{R}^s$  such that  $\mathbf{Az} = \mathbf{Ux}$  and  $\|\mathbf{x}\|_2 \leq \|\mathbf{Ux}\|_p$



**Theorem (Auerbach 1930).** For any  $\mathbf{A}$ , there is  $\mathbf{U}$  with  $\alpha = d$ .

**Conjecture.** For any  $\mathbf{A}$ , there is  $\mathbf{U}$  with  $\alpha = d^{1/p}$ .

No success in showing this conjecture! 😞

# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings


**Fact.** Oblivious  $\ell_p$  subspace embeddings reduce to constructing **well-conditioned bases** for subspaces

$\approx$  orthonormal bases for  $\ell_p$  norms

- **Idea.** What if we relax well-conditioned **bases** to well-conditioned **spanning sets**?

**Theorem (Woodruff—Y 2023).** For any  $\mathbf{A}$ , there is  $\mathbf{U} \in \mathbb{R}^{n \times s}$  for  $s = O(d \log \log d)$  such that

- $\|\mathbf{U}\|_{p,p} \leq s^{1/p}$
- for every  $\mathbf{z} \in \mathbb{R}^d$ , there is  $\mathbf{x} \in \mathbb{R}^s$  such that  $\mathbf{Az} = \mathbf{Ux}$  and  $\|\mathbf{x}\|_2 \leq \|\mathbf{Ux}\|_p$



**Theorem (Auerbach 1930).** For any  $\mathbf{A}$ , there is  $\mathbf{U}$  with  $\alpha = d$ .

**Conjecture.** For any  $\mathbf{A}$ , there is  $\mathbf{U}$  with  $\alpha = d^{1/p}$ .

No success in showing this conjecture! 😞

# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings


**Fact.** Oblivious  $\ell_p$  subspace embeddings reduce to constructing **well-conditioned bases** for subspaces

$\approx$  orthonormal bases for  $\ell_p$  norms

- **Idea.** What if we relax well-conditioned **bases** to well-conditioned **spanning sets**?

**Theorem (Woodruff—Y 2023).** For any  $\mathbf{A}$ , there is  $\mathbf{U} \in \mathbb{R}^{n \times s}$  for  $s = O(d \log \log d)$  such that

- $\|\mathbf{U}\|_{p,p} \leq s^{1/p}$
- for every  $\mathbf{z} \in \mathbb{R}^d$ , there is  $\mathbf{x} \in \mathbb{R}^s$  such that  $\mathbf{Az} = \mathbf{Ux}$  and  $\|\mathbf{x}\|_2 \leq \|\mathbf{Ux}\|_p$



**Theorem (Auerbach 1930).** For any  $\mathbf{A}$ , there is  $\mathbf{U}$  with  $\alpha = d$ .

**Conjecture.** For any  $\mathbf{A}$ , there is  $\mathbf{U}$  with  $\alpha = d^{1/p}$ .

No success in showing this conjecture! 😞

←  $\approx$  as good as showing  $\alpha = s^{1/p}$




# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings

**Fact.** Oblivious  $\ell_p$  subspace embeddings reduce to constructing **well-conditioned bases** for subspaces

$\approx$  orthonormal bases for  $\ell_p$  norms



- **Idea.** What if we relax well-conditioned **bases** to well-conditioned **spanning sets**?


**Theorem (Woodruff—Y 2023).** For any  $A$ , there is  $U \in \mathbb{R}^{n \times s}$  for  $s = O(d \log \log d)$  such that

- $\|U\|_{p,p} \leq s^{1/p}$
- for every  $z \in \mathbb{R}^d$ , there is  $x \in \mathbb{R}^s$  such that  $Az = Ux$  and  $\|x\|_2 \leq \|Ux\|_p$

**Theorem (Woodruff—Y 2023).** There are oblivious  $\ell_p$  subspace embeddings with  $r = \tilde{O}(d)$  and  $\kappa = \tilde{O}(d^{1/p})$ , which is **nearly optimal**.

# Matrix Approximation

## Oblivious $\ell_p$ Subspace Embeddings



# Subspace Embeddings and Linear Regression

- Oblivious  $\ell_p$  subspace embeddings: high distortion and low distortion




Non-oblivious subspace embeddings:  $\ell_p$  Lewis weight sampling, general losses

- Applications: active learning, streaming computational geometry, low rank approximation

# Matrix Approximation

## Non-oblivious Subspace Embeddings

Oblivious






Oblivious = does not depend on  $A$

# Matrix Approximation

## Non-oblivious Subspace Embeddings

Oblivious






Oblivious = does not depend on  $A$

Non-oblivious/  
Sampling

# Matrix Approximation


## Non-oblivious Subspace Embeddings

Oblivious



Oblivious = does not depend on  $A$


Non-oblivious/  
Sampling



# Matrix Approximation

## Non-oblivious Subspace Embeddings

Oblivious



$S$

$A$

Oblivious = does not depend on  $A$

Non-oblivious/  
Sampling

Step 1. Compute “importance scores”  
for the rows of  $A$

$q_1$

$q_2$

$\vdots$


$q_n$

$A$

# Matrix Approximation

## Non-oblivious Subspace Embeddings

Oblivious



$S$

$A$

Oblivious = does not depend on  $A$

Non-oblivious/  
Sampling

Step 1. Compute “importance scores”  
for the rows of  $A$


$q_1$

$q_2$

$\vdots$

$q_n$

$A$




Step 2. Sample rows proportionally to  
the importance scores



# Matrix Approximation

## Non-oblivious Subspace Embeddings

Oblivious




$S$

$A$

Oblivious = does not depend on  $A$

Non-oblivious/  
Sampling

Step 1. Compute “importance scores”  
for the rows of  $A$



Step 2. Sample rows proportionally to  
the importance scores

$q_1$

$q_2$

$\vdots$

$q_n$

$A$

$SA$


# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Theorem (Leverage score sampling).** For any  $\mathbf{A} \in \mathbb{R}^{n \times d}$ , there are probabilities  $q_1, q_2, \dots, q_n$  that sample  $r = \tilde{O}(\varepsilon^{-2}d)$  rows of  $\mathbf{A}$  that forms an  $\ell_2$  subspace embedding with distortion  $\kappa = (1 + \varepsilon)$ , with probability 99%.

Non-oblivious/  
Sampling

Step 1. Compute “importance scores”  
for the rows of  $\mathbf{A}$




Step 2. Sample rows proportionally to  
the importance scores

$q_1$

$q_2$

$\vdots$

$q_n$



$\mathbf{SA}$


# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Theorem (Leverage score sampling).** For any  $\mathbf{A} \in \mathbb{R}^{n \times d}$ , there are probabilities  $q_1, q_2, \dots, q_n$  that sample  $r = \tilde{O}(\varepsilon^{-2}d)$  rows of  $\mathbf{A}$  that forms an  $\ell_2$  subspace embedding with distortion  $\kappa = (1 + \varepsilon)$ , with probability 99%.

Non-oblivious/  
Sampling

Step 1. Compute “importance scores”  
for the rows of  $\mathbf{A}$




Step 2. Sample rows proportionally to  
the importance scores

$q_1$

$q_2$

$\vdots$

$q_n$



$\mathbf{A}$




$\mathbf{SA}$

# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Theorem (Leverage score sampling).** For any  $\mathbf{A} \in \mathbb{R}^{n \times d}$ , there are probabilities  $q_1, q_2, \dots, q_n$  that sample  $r = \tilde{O}(\varepsilon^{-2}d)$  rows of  $\mathbf{A}$  that forms an  $\ell_2$  subspace embedding with distortion  $\kappa = (1 + \varepsilon)$ , with probability 99%.



# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Theorem (Leverage score sampling).** For any  $\mathbf{A} \in \mathbb{R}^{n \times d}$ , there are probabilities  $q_1, q_2, \dots, q_n$  that sample  $r = \tilde{O}(\varepsilon^{-2}d)$  rows of  $\mathbf{A}$  that forms an  $\ell_2$  subspace embedding with distortion  $\kappa = (1 + \varepsilon)$ , with probability 99%.

- Same row count ( $r$ ) vs distortion ( $\kappa$ ) trade-off as the oblivious case...



# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Theorem (Leverage score sampling).** For any  $\mathbf{A} \in \mathbb{R}^{n \times d}$ , there are probabilities  $q_1, q_2, \dots, q_n$  that sample  $r = \tilde{O}(\varepsilon^{-2}d)$  rows of  $\mathbf{A}$  that forms an  $\ell_2$  subspace embedding with distortion  $\kappa = (1 + \varepsilon)$ , with probability 99%.

- Same row count ( $r$ ) vs distortion ( $\kappa$ ) trade-off as the oblivious case...
- Generalizes to much better trade-offs for  $\ell_p$  norms with  $p \neq 2$ !



# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Theorem (Leverage score sampling).** For any  $\mathbf{A} \in \mathbb{R}^{n \times d}$ , there are probabilities  $q_1, q_2, \dots, q_n$  that sample  $r = \tilde{O}(\varepsilon^{-2}d)$  rows of  $\mathbf{A}$  that forms an  $\ell_2$  subspace embedding with distortion  $\kappa = (1 + \varepsilon)$ , with probability 99%.

- Same row count ( $r$ ) vs distortion ( $\kappa$ ) trade-off as the oblivious case...
- Generalizes to much better trade-offs for  $\ell_p$  norms with  $p \neq 2$ !
  - Oblivious: either  $\kappa = \text{poly}(d)$  or  $r \gg \text{poly}(d)$ , and only for  $p \leq 2$ ...





# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Theorem (Leverage score sampling).** For any  $\mathbf{A} \in \mathbb{R}^{n \times d}$ , there are probabilities  $q_1, q_2, \dots, q_n$  that sample  $r = \tilde{O}(\varepsilon^{-2}d)$  rows of  $\mathbf{A}$  that forms an  $\ell_2$  subspace embedding with distortion  $\kappa = (1 + \varepsilon)$ , with probability 99%.

- Same row count ( $r$ ) vs distortion ( $\kappa$ ) trade-off as the oblivious case...
- Generalizes to much better trade-offs for  $\ell_p$  norms with  $p \neq 2$ !
  - Oblivious: either  $\kappa = \text{poly}(d)$  or  $r \gg \text{poly}(d)$ , and only for  $p \leq 2$ ...
  - Non-oblivious:  $\kappa = (1 + \varepsilon)$  and  $r = \text{poly}(d)$  for any fixed  $p$ !






# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Definition (Leverage scores).** For  $\mathbf{A} \in \mathbb{R}^{n \times d}$  and  $i \in [n]$ , the  $i$ th leverage score is

$$\tau_i(\mathbf{A}) = \sup_{\mathbf{Ax} \neq 0} \frac{\langle \mathbf{a}_i, \mathbf{x} \rangle^2}{\|\mathbf{Ax}\|_2^2}$$




# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Definition (Leverage scores).** For  $\mathbf{A} \in \mathbb{R}^{n \times d}$  and  $i \in [n]$ , the  $i$ th leverage score is

$$\tau_i(\mathbf{A}) = \sup_{\mathbf{Ax} \neq 0} \frac{\langle \mathbf{a}_i, \mathbf{x} \rangle^2}{\|\mathbf{Ax}\|_2^2}$$

$\ell_2$  norm of  $\mathbf{Ax}$



# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Definition (Leverage scores).** For  $\mathbf{A} \in \mathbb{R}^{n \times d}$  and  $i \in [n]$ , the  $i$ th leverage score is

$$\tau_i(\mathbf{A}) = \sup_{\mathbf{Ax} \neq 0} \frac{\langle \mathbf{a}_i, \mathbf{x} \rangle^2}{\|\mathbf{Ax}\|_2^2}$$

Contribution of the  $i$ th coordinate

$\ell_2$  norm of  $\mathbf{Ax}$



# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Definition (Leverage scores).** For  $\mathbf{A} \in \mathbb{R}^{n \times d}$  and  $i \in [n]$ , the  $i$ th leverage score is

$$\tau_i(\mathbf{A}) = \sup_{\mathbf{Ax} \neq 0} \frac{\langle \mathbf{a}_i, \mathbf{x} \rangle^2}{\|\mathbf{Ax}\|_2^2}$$

Contribution of the  $i$ th coordinate

$\ell_2$  norm of  $\mathbf{Ax}$

“Importance score”: largest fraction of  $\ell_2$  norm occupied by the  $i$ th coordinate



# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Definition (Leverage scores).** For  $\mathbf{A} \in \mathbb{R}^{n \times d}$  and  $i \in [n]$ , the  $i$ th leverage score is

$$\tau_i(\mathbf{A}) = \sup_{\mathbf{Ax} \neq 0} \frac{\langle \mathbf{a}_i, \mathbf{x} \rangle^2}{\|\mathbf{Ax}\|_2^2}$$

Contribution of the  $i$ th coordinate

$\ell_2$  norm of  $\mathbf{Ax}$

“Importance score”: largest fraction of  $\ell_2$  norm occupied by the  $i$ th coordinate

- Many generalizations:



# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Definition (Leverage scores).** For  $\mathbf{A} \in \mathbb{R}^{n \times d}$  and  $i \in [n]$ , the  $i$ th leverage score is

$$\tau_i(\mathbf{A}) = \sup_{\mathbf{Ax} \neq 0} \frac{\langle \mathbf{a}_i, \mathbf{x} \rangle^2}{\|\mathbf{Ax}\|_2^2}$$

Contribution of the  $i$ th coordinate

$\ell_2$  norm of  $\mathbf{Ax}$

“Importance score”: largest fraction of  $\ell_2$  norm occupied by the  $i$ th coordinate

- Many generalizations:
  - $\ell_p$  sensitivity scores (Langberg—Schulman 2010)



# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Definition (Leverage scores).** For  $\mathbf{A} \in \mathbb{R}^{n \times d}$  and  $i \in [n]$ , the  $i$ th leverage score is

$$\tau_i(\mathbf{A}) = \sup_{\mathbf{Ax} \neq 0} \frac{\langle \mathbf{a}_i, \mathbf{x} \rangle^2}{\|\mathbf{Ax}\|_2^2}$$

Contribution of the  $i$ th coordinate

$\ell_2$  norm of  $\mathbf{Ax}$

“Importance score”: largest fraction of  $\ell_2$  norm occupied by the  $i$ th coordinate

- Many generalizations:
  - $\ell_p$  sensitivity scores (Langberg—Schulman 2010)
  - $\ell_p$  Lewis weights (Lewis 1978)



# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Definition (Leverage scores).** For  $\mathbf{A} \in \mathbb{R}^{n \times d}$  and  $i \in [n]$ , the  $i$ th leverage score is

$$\tau_i(\mathbf{A}) = \sup_{\mathbf{Ax} \neq 0} \frac{\langle \mathbf{a}_i, \mathbf{x} \rangle^2}{\|\mathbf{Ax}\|_2^2}$$

Contribution of the  $i$ th coordinate

$\ell_2$  norm of  $\mathbf{Ax}$

“Importance score”: largest fraction of  $\ell_2$  norm occupied by the  $i$ th coordinate

- Many generalizations:
  - $\ell_p$  sensitivity scores (Langberg—Schulman 2010)
  - $\ell_p$  Lewis weights (Lewis 1978)
  - Online leverage scores (Cohen—Musco—Pachockski 2016)





# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Definition (Leverage scores).** For  $\mathbf{A} \in \mathbb{R}^{n \times d}$  and  $i \in [n]$ , the  $i$ th leverage score is

$$\tau_i(\mathbf{A}) = \sup_{\mathbf{Ax} \neq 0} \frac{\langle \mathbf{a}_i, \mathbf{x} \rangle^2}{\|\mathbf{Ax}\|_2^2}$$

Contribution of the  $i$ th coordinate

$\ell_2$  norm of  $\mathbf{Ax}$

“Importance score”: largest fraction of  $\ell_2$  norm occupied by the  $i$ th coordinate

- Many generalizations:
  - $\ell_p$  sensitivity scores (Langberg—Schulman 2010)
  - $\ell_p$  Lewis weights (Lewis 1978)
  - Online leverage scores (Cohen—Musco—Pachockski 2016)



# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Theorem (Lewis weight sampling, Cohen—Peng 2015).** For any  $\mathbf{A} \in \mathbb{R}^{n \times d}$ , there are probabilities  $q_1, q_2, \dots, q_n$  that sample

$$r = \begin{cases} \tilde{O}(\varepsilon^{-2}d) & p < 2 \\ \tilde{O}(\varepsilon^{-5}d^{p/2}) & p > 2 \end{cases}$$

rows of  $\mathbf{A}$  that forms an  $\ell_p$  subspace embedding with distortion  $\kappa = (1 + \varepsilon)$ , with probability 99%.



# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Theorem (Lewis weight sampling, Cohen—Peng 2015).** For any  $\mathbf{A} \in \mathbb{R}^{n \times d}$ , there are probabilities  $q_1, q_2, \dots, q_n$  that sample

$$r = \begin{cases} \tilde{O}(\varepsilon^{-2}d) & p < 2 \\ \tilde{O}(\varepsilon^{-5}d^{p/2}) & p > 2 \end{cases}$$

rows of  $\mathbf{A}$  that forms an  $\ell_p$  subspace embedding with distortion


$\kappa = (1 + \varepsilon)$  with probability 99%.



# Matrix Approximation

## Non-oblivious Subspace Embeddings

$$r = \begin{cases} \tilde{O}(\varepsilon^{-2}d) & p < 2 \\ \tilde{O}(\varepsilon^{-5}d^{p/2}) & p > 2 \end{cases}$$




# Matrix Approximation

## Non-oblivious Subspace Embeddings

Two questions:

$$r = \begin{cases} \tilde{O}(\varepsilon^{-2}d) & p < 2 \\ \tilde{O}(\varepsilon^{-5}d^{p/2}) & p > 2 \end{cases}$$



# Matrix Approximation

## Non-oblivious Subspace Embeddings

Two questions:

**Question 1.** Can these bounds be improved?

$$r = \begin{cases} \tilde{O}(\varepsilon^{-2}d) & p < 2 \\ \tilde{O}(\varepsilon^{-5}d^{p/2}) & p > 2 \end{cases}$$



# Matrix Approximation

## Non-oblivious Subspace Embeddings

Two questions:

**Question 1.** Can these bounds be improved?

$$r = \begin{cases} \tilde{O}(\varepsilon^{-2}d) & p < 2 \\ \tilde{O}(\varepsilon^{-5}d^{p/2}) & p > 2 \end{cases}$$

**Question 2.** What if the rows of  $\mathbf{A}$  arrive one by one in a stream?



# Matrix Approximation

## Non-oblivious Subspace Embeddings

Two questions:

**Question 1.** Can these bounds be improved?

$$r = \begin{cases} \tilde{O}(\varepsilon^{-2}d) & p < 2 \\ \tilde{O}(\varepsilon^{-5}d^{p/2}) & p > 2 \end{cases}$$

Nearly optimal (Li—Wang—Woodruff 2020)

**Question 2.** What if the rows of  $A$  arrive one by one in a stream?





# Matrix Approximation

## Non-oblivious Subspace Embeddings

Two questions:

**Question 1.** Can these bounds be improved?

$$r = \begin{cases} \tilde{O}(\varepsilon^{-2}d) & p < 2 \\ \tilde{O}(\varepsilon^{-5}d^{p/2}) & p > 2 \end{cases}$$

Nearly optimal (Li—Wang—Woodruff 2020)

$d^{p/2}$  is nearly optimal for  $\varepsilon = O(1)$   
(Li—Wang—Woodruff 2020)

**Question 2.** What if the rows of  $\mathbf{A}$  arrive one by one in a stream?



# Matrix Approximation

## Non-oblivious Subspace Embeddings

Two questions:

**Question 1.** Can these bounds be improved?

$$r = \begin{cases} \tilde{O}(\varepsilon^{-2}d) & p < 2 \\ \tilde{O}(\varepsilon^{-5}d^{p/2}) & p > 2 \end{cases}$$

Nearly optimal (Li—Wang—Woodruff 2020)

$\varepsilon^{-2}$  should be possible here!

$d^{p/2}$  is nearly optimal for  $\varepsilon = O(1)$   
(Li—Wang—Woodruff 2020)

**Question 2.** What if the rows of  $\mathbf{A}$  arrive one by one in a stream?



# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Question 1.** Can these bounds be improved?

$$r = \begin{cases} \tilde{O}(\varepsilon^{-2}d) & p < 2 \\ \tilde{O}(\varepsilon^{-5}d^{p/2}) & p > 2 \end{cases}$$

**Question 2.** What if the rows of  $\mathbf{A}$  arrive one by one in a stream?



# Matrix Approximation

## Non-oblivious Subspace Embeddings

Woodruff—Y 2023

**Question 1.** Can these bounds be improved?

$$r = \begin{cases} \tilde{O}(\varepsilon^{-2}d) & p < 2 \\ \tilde{O}(\varepsilon^{-5}d^{p/2}) & p > 2 \end{cases}$$

**Question 2.** What if the rows of  $\mathbf{A}$  arrive one by one in a stream?



# Matrix Approximation

## Non-oblivious Subspace Embeddings

Woodruff—Y 2023

**Question 1.** Can these bounds be improved?

Questions 1 and 2 can be addressed using related techniques!

$$r = \begin{cases} \tilde{O}(\varepsilon^{-2}d) & p < 2 \\ \tilde{O}(\varepsilon^{-5}d^{p/2}) & p > 2 \end{cases}$$

**Question 2.** What if the rows of  $\mathbf{A}$  arrive one by one in a stream?



# Matrix Approximation

## Non-oblivious Subspace Embeddings

Woodruff—Y 2023

**Question 1.** Can these bounds be improved?

Questions 1 and 2 can be addressed using related techniques!

$$r = \begin{cases} \tilde{O}(\varepsilon^{-2}d) & p < 2 \\ \tilde{O}(\varepsilon^{-5}d^{p/2}) & p > 2 \end{cases}$$

**Question 2.** What if the rows of  $\mathbf{A}$  arrive one by one in a stream?

Handling addition of rows in a stream  $\rightarrow$  better sampling bounds



# Matrix Approximation

## Non-oblivious Subspace Embeddings

Woodruff—Y 2023

**Question 1.** Can these bounds be improved?

Questions 1 and 2 can be addressed using related techniques!

$$r = \begin{cases} \tilde{O}(\varepsilon^{-2}d) & p < 2 \\ \tilde{O}(\varepsilon^{-2}d^{p/2}) & p > 2 \end{cases}$$

**Question 2.** What if the rows of  $\mathbf{A}$  arrive one by one in a stream?

Handling addition of rows in a stream  $\rightarrow$  better sampling bounds





# Matrix Approximation

## Non-oblivious Subspace Embeddings

Woodruff—Y 2023

**Question 1.** Can these bounds be improved?

Questions 1 and 2 can be addressed using related techniques!

$$r = \begin{cases} \tilde{O}(\varepsilon^{-2}d) & p < 2 \\ \tilde{O}(\varepsilon^{-2}d^{p/2}) & p > 2 \end{cases}$$

$\tilde{O}(\varepsilon^{-2}d^{p/2})$  (Woodruff—Y 2023)

**Question 2.** What if the rows of  $\mathbf{A}$  arrive one by one in a stream?

Handling addition of rows in a stream  $\rightarrow$  better sampling bounds





# Matrix Approximation

## Non-oblivious Subspace Embeddings

Woodruff—Y 2023

**Question 1.** Can these bounds be improved?

Questions 1 and 2 can be addressed using related techniques!

$$r = \begin{cases} \tilde{O}(\varepsilon^{-2}d) & p < 2 \\ \tilde{O}(\varepsilon^{-2}d^{p/2}) & p > 2 \end{cases}$$

$\tilde{O}(\varepsilon^{-2}d^{p/2})$  (Woodruff—Y 2023)

These bounds also hold in the streaming setting (Woodruff—Y 2023)

**Question 2.** What if the rows of  $\mathbf{A}$  arrive one by one in a stream?


Handling addition of rows in a stream → better sampling bounds



# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Question 3.** Are there similar results for other loss functions?



# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Question 3.** Are there similar results for other loss functions?

$$\|\mathbf{Ax}\|_g := \sum_{i=1}^n g([\mathbf{Ax}](i))$$




# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Question 3.** Are there similar results for other loss functions?

$$\|\mathbf{Ax}\|_g := \sum_{i=1}^n g([\mathbf{Ax}](i))$$

- There are many other loss functions used for linear regression...






# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Question 3.** Are there similar results for other loss functions?

$$\|\mathbf{Ax}\|_g := \sum_{i=1}^n g([\mathbf{Ax}](i))$$

- There are many other loss functions used for linear regression...
  - Huber/Tukey loss for robust statistics






# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Question 3.** Are there similar results for other loss functions?

$$\|\mathbf{Ax}\|_g := \sum_{i=1}^n g([\mathbf{Ax}](i))$$

- There are many other loss functions used for linear regression...
  - Huber/Tukey loss for robust statistics
  - Logistic regression for classification



# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Question 3.** Are there similar results for other loss functions?

$$\|\mathbf{Ax}\|_g := \sum_{i=1}^n g([\mathbf{Ax}](i))$$



# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Question 3.** Are there similar results for other loss functions?

$$\|\mathbf{Ax}\|_g := \sum_{i=1}^n g([\mathbf{Ax}](i))$$

Musco—Musco—Woodruff—Y 2022





# Matrix Approximation


## Non-oblivious Subspace Embeddings

**Question 3.** Are there similar results for other loss functions?

$$\|\mathbf{Ax}\|_g := \sum_{i=1}^n g([\mathbf{Ax}](i))$$

Musco—Musco—Woodruff—Y 2022

General losses with quadratic growth



# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Question 3.** Are there similar results for other loss functions?

$$\|\mathbf{Ax}\|_g := \sum_{i=1}^n g([\mathbf{Ax}](i))$$

Musco—Musco—Woodruff—Y 2022

General losses with quadratic growth

$$r = \tilde{O}(\varepsilon^{-2}d^2)$$



# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Question 3.** Are there similar results for other loss functions?



$$\|\mathbf{Ax}\|_g := \sum_{i=1}^n g([\mathbf{Ax}](i))$$

Musco—Musco—Woodruff—Y 2022

General losses with quadratic growth

$$r = \tilde{O}(\varepsilon^{-2}d^2)$$

Huber loss



# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Question 3.** Are there similar results for other loss functions?


$$\|\mathbf{Ax}\|_g := \sum_{i=1}^n g([\mathbf{Ax}](i))$$

Musco—Musco—Woodruff—Y 2022

General losses with quadratic growth

$$r = \tilde{O}(\varepsilon^{-2}d^2)$$

Huber loss



$$r = \text{poly}(\varepsilon^{-1})d^{4-2\sqrt{2}} \approx d^{1.172}$$



# Matrix Approximation

## Non-oblivious Subspace Embeddings

**Question 3.** Are there similar results for other loss functions?


$$\|\mathbf{Ax}\|_g := \sum_{i=1}^n g([\mathbf{Ax}](i))$$

Musco—Musco—Woodruff—Y 2022

General losses with quadratic growth


$$r = \tilde{O}(\varepsilon^{-2}d^2)$$

Huber loss



$$r = \text{poly}(\varepsilon^{-1})d^{4-2\sqrt{2}} \approx d^{1.172}$$

I am very interested in improving this to  $d$



# Subspace Embeddings and Linear Regression

- Oblivious  $\ell_p$  subspace embeddings: high distortion and low distortion
- Non-oblivious subspace embeddings:  $\ell_p$  Lewis weight sampling, general losses

Applications: active learning, streaming computational geometry, low rank approximation

# Matrix Approximation

Active  $\ell_p$  Linear Regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \left\| \mathbf{Ax} - \mathbf{b} \right\|_p^p$$

# Matrix Approximation

## Active $\ell_p$ Linear Regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_p^p$$

- Active learning: machine learning, where **label acquisition** is the most expensive resource



# Matrix Approximation

## Active $\ell_p$ Linear Regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \left\| \mathbf{Ax} - \mathbf{b} \right\|_p^p$$

- Active learning: machine learning, where **label acquisition** is the most expensive resource
  - Labeling could require manual labor

# Matrix Approximation

## Active $\ell_p$ Linear Regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \left\| \mathbf{Ax} - \mathbf{b} \right\|_p^p$$

- Active learning: machine learning, where **label acquisition** is the most expensive resource
  - Labeling could require manual labor
  - Labeling could require purchasing information

# Matrix Approximation

## Active $\ell_p$ Linear Regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \left\| \mathbf{Ax} - \mathbf{b} \right\|_p^p$$

- Active learning: machine learning, where **label acquisition** is the most expensive resource
  - Labeling could require manual labor
  - Labeling could require purchasing information
  - Labeling could require involve an invasive medical procedure

# Matrix Approximation

## Active $\ell_p$ Linear Regression


$$\min_{\mathbf{x} \in \mathbb{R}^d} \left\| \mathbf{Ax} - \mathbf{b} \right\|_p^p$$

- Active learning: machine learning, where **label acquisition** is the most expensive resource
  - Labeling could require manual labor
  - Labeling could require purchasing information
  - Labeling could require involve an invasive medical procedure
- Goal: **minimize the number of label entries** that are read

# Matrix Approximation

Active  $\ell_p$  Linear Regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_p^p$$


$$\mathbf{A} \mathbf{x} - \mathbf{b}$$

# Matrix Approximation

## Active $\ell_p$ Linear Regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_p^p$$

The algorithm has full access to  $\mathbf{A}$

$$\mathbf{A} \quad \mathbf{x} \quad - \quad \mathbf{b}$$

# Matrix Approximation

## Active $\ell_p$ Linear Regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_p^p$$

The algorithm has full access to  $\mathbf{A}$

$\mathbf{A}$

$\mathbf{x}$

$-$

$\mathbf{b}$

$\mathbf{b}$  vector is hidden, and the algorithm has query access to it

# Matrix Approximation

## Active $\ell_p$ Linear Regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_p^p$$

The algorithm has full access to  $\mathbf{A}$

$\mathbf{A}$

$\mathbf{x}$

$-$

$\mathbf{b}$

$\mathbf{b}$  vector is hidden, and the algorithm has query access to it


**Question.** How many entries of  $\mathbf{b}$  need to be read?



# Matrix Approximation

Active  $\ell_p$  Linear Regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_p^p$$


$$\mathbf{A} \mathbf{x} - \mathbf{b}$$


**Question.** How many entries of  $\mathbf{b}$  need to be read?

# Matrix Approximation

## Active $\ell_p$ Linear Regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_p^p$$

Sampling-based  $\ell_p$   
subspace embedding for  $\mathbf{A}$




$$\mathbf{A} \quad \mathbf{x} \quad - \quad \mathbf{b}$$

**Question.** How many entries of  $\mathbf{b}$  need to be read?

# Matrix Approximation

Active  $\ell_p$  Linear Regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \left\| \mathbf{A}\mathbf{x} - \mathbf{b} \right\|_p^p$$




**Question.** How many entries of  $\mathbf{b}$  need to be read?

# Matrix Approximation

Active  $\ell_p$  Linear Regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_p^p$$




**Question.** How many entries of  $\mathbf{b}$  need to be read?

# Matrix Approximation

Active  $\ell_p$  Linear Regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_p^p$$




**Question.** How many entries of  $\mathbf{b}$  need to be read?

# Matrix Approximation

## Active $\ell_p$ Linear Regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_p^p$$

- $p = 2$ :  $\Theta(\varepsilon^{-1}d)$  (Chen—Price 2019)




**Question.** How many entries of  $\mathbf{b}$  need to be read?

# Matrix Approximation

## Active $\ell_p$ Linear Regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_p^p$$




- $p = 2$ :  $\Theta(\varepsilon^{-1}d)$  (Chen—Price 2019)
- $p = 1$ :  $\tilde{\Theta}(\varepsilon^{-2}d)$  (Chen—Derezinski, Parulekar—Parulekar—Price 2021)

**Question.** How many entries of  $\mathbf{b}$  need to be read?

# Matrix Approximation

## Active $\ell_p$ Linear Regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_p^p$$



- $p = 2$ :  $\Theta(\varepsilon^{-1}d)$  (Chen—Price 2019)
- $p = 1$ :  $\tilde{\Theta}(\varepsilon^{-2}d)$  (Chen—Derezinski, Parulekar—Parulekar—Price 2021)
- $1 < p < 2$ :  $\tilde{O}(\varepsilon^{-2}d^2)$  (Chen—Derezinski 2021)


**Question.** How many entries of  $\mathbf{b}$  need to be read?



# Matrix Approximation

## Active $\ell_p$ Linear Regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_p^p$$




- $p = 2$ :  $\Theta(\varepsilon^{-1}d)$  (Chen—Price 2019)
- $p = 1$ :  $\tilde{\Theta}(\varepsilon^{-2}d)$  (Chen—Derezinski, Parulekar—Parulekar—Price 2021)
- $1 < p < 2$ :  $\tilde{O}(\varepsilon^{-2}d^2)$  (Chen—Derezinski 2021)

**Question.** How many entries of  $\mathbf{b}$  need to be read?

# Matrix Approximation

## Active $\ell_p$ Linear Regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_p^p$$




- $p = 2$ :  $\Theta(\varepsilon^{-1}d)$  (Chen—Price 2019)
- $p = 1$ :  $\tilde{\Theta}(\varepsilon^{-2}d)$  (Chen—Derezinski, Parulekar—Parulekar—Price 2021)
- $1 < p < 2$ :  ~~$\tilde{O}(\varepsilon^{-2}d^2)$~~  (Chen—Derezinski 2021)  
 $\tilde{\Theta}(\varepsilon^{-1}d)$  (Musco—Musco—Woodruff—Y 2022)

**Question.** How many entries of  $\mathbf{b}$  need to be read?

# Matrix Approximation

## Active $\ell_p$ Linear Regression

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_p^p$$



- $p = 2$ :  $\Theta(\varepsilon^{-1}d)$  (Chen—Price 2019)
- $p = 1$ :  $\tilde{\Theta}(\varepsilon^{-2}d)$  (Chen—Derezinski, Parulekar—Parulekar—Price 2021)
- $1 < p < 2$ :  ~~$\tilde{O}(\varepsilon^{-2}d^2)$~~  (Chen—Derezinski 2021)  
 $\tilde{\Theta}(\varepsilon^{-1}d)$  (Musco—Musco—Woodruff—Y 2022)
- $2 < p < \infty$ :  $\tilde{\Theta}(\varepsilon^{1-p}d^{p/2})$  (Woodruff—Y 2023)

**Question.** How many entries of  $\mathbf{b}$  need to be read?

# Matrix Approximation

**Streaming Löwner—John Ellipsoids**

# Matrix Approximation


## **Streaming Löwner—John Ellipsoids**

Input: symmetric polytope with  
 $2n$  faces in  $d$  dims

# Matrix Approximation

## Streaming Löwner—John Ellipsoids

Input: symmetric polytope with  
 $2n$  faces in  $d$  dims


$$\{\mathbf{x} : \langle \mathbf{a}_i, \mathbf{x} \rangle \leq 1\}$$


# Matrix Approximation

## Streaming Löwner—John Ellipsoids

Input: symmetric polytope with  
 $2n$  faces in  $d$  dims

$$\{\mathbf{x} : \langle \mathbf{a}_i, \mathbf{x} \rangle \leq 1\}$$




# Matrix Approximation

## Streaming Löwner—John Ellipsoids

Input: symmetric polytope with  
 $2n$  faces in  $d$  dims

$$\{\mathbf{x} : \langle \mathbf{a}_i, \mathbf{x} \rangle \leq 1\}$$





# Matrix Approximation

## Streaming Löwner—John Ellipsoids

Input: symmetric polytope with  
 $2n$  faces in  $d$  dims

Want the “largest” ellipsoid enclosed in polytope

$$\{\mathbf{x} : \langle \mathbf{a}_i, \mathbf{x} \rangle \leq 1\}$$


$$\{\mathbf{x} : \|\mathbf{E}\mathbf{x}\|_2^2 \leq 1\}$$
A diagram illustrating the concept of a Löwner-John ellipsoid. It shows a symmetric polytope defined by several black lines representing its faces. Inside the polytope, a blue circle represents the largest ellipsoid that can be inscribed. A yellow callout bubble points to the ellipsoid with the text 'Want the “largest” ellipsoid enclosed in polytope'. The polytope is defined by the set of points x such that the inner product of x with each of the 2n vectors a\_i is less than or equal to 1. The ellipsoid is defined by the set of points x such that the squared norm of E times x is less than or equal to 1.

# Matrix Approximation

## Streaming Löwner—John Ellipsoids

Input: symmetric polytope with  
 $2n$  faces in  $d$  dims

Want the “largest” ellipsoid enclosed in polytope



**Question.** Can Löwner—John ellipsoids be maintained in  $\text{poly}(d, \log n)$  bits of space?


# Matrix Approximation

## Streaming Löwner—John Ellipsoids

Input: symmetric polytope with  
 $2n$  faces in  $d$  dims

Want the “largest” ellipsoid enclosed in polytope

- Agarwal—Har-Peled—Varadarajan 2004:  
 $\exp(\Theta(d))$  bits




**Question.** Can Löwner—John ellipsoids be maintained in  $\text{poly}(d, \log n)$  bits of space?

# Matrix Approximation

## Streaming Löwner—John Ellipsoids

Input: symmetric polytope with  
 $2n$  faces in  $d$  dims

Want the “largest” ellipsoid enclosed in polytope



- Agarwal—Har-Peled—Varadarajan 2004:  
 $\exp(\Theta(d))$  bits
- Woodruff—Y 2022:  $O(d^2 \log^2 n)$  bits

**Question.** Can Löwner—John ellipsoids be maintained in  $\text{poly}(d, \log n)$  bits of space?

# Matrix Approximation

## Streaming Löwner—John Ellipsoids

Input: symmetric polytope with  
 $2n$  faces in  $d$  dims

Want the “largest” ellipsoid enclosed in polytope

$$\{\mathbf{x} : \langle \mathbf{a}_i, \mathbf{x} \rangle \leq 1\}$$

$$\{\mathbf{x} : \|\mathbf{E}\mathbf{x}\|_2^2 \leq 1\}$$

- Agarwal—Har-Peled—Varadarajan 2004:  
 $\exp(\Theta(d))$  bits
- Woodruff—Y 2022:  $O(d^2 \log^2 n)$  bits
- Approach: select a subset of constraints

**Question.** Can Löwner—John ellipsoids be maintained in  $\text{poly}(d, \log n)$  bits of space?

# Matrix Approximation

## Streaming Löwner—John Ellipsoids

Input: symmetric polytope with  
 $2n$  faces in  $d$  dims

Want the “largest” ellipsoid enclosed in polytope

$$\{\mathbf{x} : \langle \mathbf{a}_i, \mathbf{x} \rangle \leq 1\}$$

$$\{\mathbf{x} : \|\mathbf{E}\mathbf{x}\|_2^2 \leq 1\}$$

- Agarwal—Har-Peled—Varadarajan 2004:  
 $\exp(\Theta(d))$  bits
- Woodruff—Y 2022:  $O(d^2 \log^2 n)$  bits
- Approach: select a subset of constraints
- **Techniques:** use leverage scores to...

**Question.** Can Löwner—John ellipsoids be maintained in  $\text{poly}(d, \log n)$  bits of space?




# Matrix Approximation

## Streaming Löwner—John Ellipsoids

Input: symmetric polytope with  $2n$  faces in  $d$  dims

Want the “largest” ellipsoid enclosed in polytope



- Agarwal—Har-Peled—Varadarajan 2004:  $\exp(\Theta(d))$  bits

- Woodruff—Y 2022:  $O(d^2 \log^2 n)$  bits

- Approach: select a subset of constraints
- **Techniques:** use leverage scores to...
  - test if ellipsoid respects a new constraint  $\mathbf{a}_i$


**Question.** Can Löwner—John ellipsoids be maintained in  $\text{poly}(d, \log n)$  bits of space?

# Matrix Approximation

## Streaming Löwner—John Ellipsoids

Input: symmetric polytope with  $2n$  faces in  $d$  dims

Want the “largest” ellipsoid enclosed in polytope



- Agarwal—Har-Peled—Varadarajan 2004:  $\exp(\Theta(d))$  bits

- Woodruff—Y 2022:  $O(d^2 \log^2 n)$  bits

- Approach: select a subset of constraints
- **Techniques:** use leverage scores to...
  - test if ellipsoid respects a new constraint  $\mathbf{a}_i$
  - bound # of times we keep a new constraint  $\mathbf{a}_i$

**Question.** Can Löwner—John ellipsoids be maintained in  $\text{poly}(d, \log n)$  bits of space?




# Matrix Approximation

## Streaming Löwner—John Ellipsoids

Input: symmetric polytope with  
 $2n$  faces in  $d$  dims

Want the “largest” ellipsoid enclosed in polytope



- Agarwal—Har-Peled—Varadarajan 2004:  
 $\exp(\Theta(d))$  bits

- Woodruff—Y 2022:  $O(d^2 \log^2 n)$  bits

- Approach: select a subset of constraints
- **Techniques:** use leverage scores to...
  - test if ellipsoid respects a new constraint  $\mathbf{a}_i$
  - bound # of times we keep a new constraint  $\mathbf{a}_i$

$$\sup_{\mathbf{E}\mathbf{x} \neq 0} \frac{\langle \mathbf{a}_i, \mathbf{x} \rangle^2}{\|\mathbf{E}\mathbf{x}\|_2^2}$$


**Question.** Can Löwner—John ellipsoids be maintained in  $\text{poly}(d, \log n)$  bits of space?

# Matrix Approximation

## Streaming Löwner—John Ellipsoids

Input: symmetric polytope with  $2n$  faces in  $d$  dims

Want the “largest” ellipsoid enclosed in polytope



- Agarwal—Har-Peled—Varadarajan 2004:  $\exp(\Theta(d))$  bits

- Woodruff—Y 2022:  $O(d^2 \log^2 n)$  bits

- Approach: select a subset of constraints
- **Techniques:** use leverage scores to...
  - test if ellipsoid respects a new constraint  $\mathbf{a}_i$
  - bound # of times we keep a new constraint  $\mathbf{a}_i$

$\leq 1$  iff ellipsoid  $\mathbf{E}$   
respects constraint  $\mathbf{a}_i$

$$\sup_{\mathbf{E}\mathbf{x} \neq 0} \frac{\langle \mathbf{a}_i, \mathbf{x} \rangle^2}{\|\mathbf{E}\mathbf{x}\|_2^2}$$


**Question.** Can Löwner—John ellipsoids be maintained in  $\text{poly}(d, \log n)$  bits of space?

# Matrix Approximation

## Streaming Löwner—John Ellipsoids

Input: symmetric polytope with  
 $2n$  faces in  $d$  dims

Want the “largest” ellipsoid enclosed in polytope



**Question.** Can Löwner—John ellipsoids be maintained in  $\text{poly}(d, \log n)$  bits of space?

- Agarwal—Har-Peled—Varadarajan 2004:  
 $\exp(\Theta(d))$  bits

- Woodruff—Y 2022:  $O(d^2 \log^2 n)$  bits

- Approach: select a subset of constraints
- **Techniques:** use leverage scores to...
  - test if ellipsoid respects a new constraint  $\mathbf{a}_i$
  - bound # of times we keep a new constraint  $\mathbf{a}_i$

$\leq 1$  iff ellipsoid  $\mathbf{E}$   
respects constraint  $\mathbf{a}_i$

$$\sup_{\mathbf{E}\mathbf{x} \neq 0} \frac{\langle \mathbf{a}_i, \mathbf{x} \rangle^2}{\|\mathbf{E}\mathbf{x}\|_2^2}$$

Leverage scores:  $\tau_i(\mathbf{A}) = \sup_{\mathbf{A}\mathbf{x} \neq 0} \frac{\langle \mathbf{a}_i, \mathbf{x} \rangle^2}{\|\mathbf{A}\mathbf{x}\|_2^2}$

# Matrix Approximation

## **Low Rank Approximation with General Losses**

# Matrix Approximation


## Low Rank Approximation with General Losses

$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_F^2$$

# Matrix Approximation

## Low Rank Approximation with General Losses

$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_F^2$$



$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_{p,p}^p = \sum_{i,j} (\mathbf{A} - \mathbf{UV})_{i,j}^p$$

# Matrix Approximation

## Low Rank Approximation with General Losses

$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_F^2$$

$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_{p,p}^p = \sum_{i,j} (\mathbf{A} - \mathbf{UV})_{i,j}^p$$

$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_g = \sum_{i,j} g((\mathbf{A} - \mathbf{UV})_{i,j})$$



# Matrix Approximation

## Low Rank Approximation with General Losses

$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_F^2$$

$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_{p,p}^p = \sum_{i,j} (\mathbf{A} - \mathbf{UV})_{i,j}^p$$

$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_g = \sum_{i,j} g((\mathbf{A} - \mathbf{UV})_{i,j})$$

- NP-hard... → need approximation/bicriteria algorithms



# Matrix Approximation

## Low Rank Approximation with General Losses

$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_F^2$$

$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_{p,p}^p = \sum_{i,j} (\mathbf{A} - \mathbf{UV})_{i,j}^p$$

$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_g = \sum_{i,j} g((\mathbf{A} - \mathbf{UV})_{i,j})$$

- NP-hard... → need approximation/bicriteria algorithms
- Focus on **column subset selection** algorithms:  $\mathbf{U}$  is a set of  $\approx k$  columns of  $\mathbf{A}$

# Matrix Approximation

## Low Rank Approximation with General Losses

$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_F^2$$

$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_{p,p}^p = \sum_{i,j} (\mathbf{A} - \mathbf{UV})_{i,j}^p$$

$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_g = \sum_{i,j} g((\mathbf{A} - \mathbf{UV})_{i,j})$$

- NP-hard... → need approximation/bicriteria algorithms
- Focus on **column subset selection** algorithms:  $\mathbf{U}$  is a set of  $\approx k$  columns of  $\mathbf{A}$ 
  - Useful for unsupervised feature selection

# Matrix Approximation

## Low Rank Approximation with General Losses

$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_F^2$$

$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_{p,p}^p = \sum_{i,j} (\mathbf{A} - \mathbf{UV})_{i,j}^p$$

$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_g = \sum_{i,j} g((\mathbf{A} - \mathbf{UV})_{i,j})$$

- NP-hard... → need approximation/bicriteria algorithms
- Focus on **column subset selection** algorithms:  $\mathbf{U}$  is a set of  $\approx k$  columns of  $\mathbf{A}$ 
  - Useful for unsupervised feature selection
  - This framework gives the best known algorithms for this problem!

# Matrix Approximation

## Low Rank Approximation with General Losses

$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_g = \sum_{i,j} g((\mathbf{A} - \mathbf{UV})_{i,j})$$

# Matrix Approximation

## Low Rank Approximation with General Losses

$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \left\| \mathbf{A} - \mathbf{UV} \right\|_g = \sum_{i,j} g((\mathbf{A} - \mathbf{UV})_{i,j})$$

$$\left\| \mathbf{A} - \hat{\mathbf{U}}\hat{\mathbf{V}} \right\|_g \leq \kappa \min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \left\| \mathbf{A} - \mathbf{UV} \right\|_g$$

# Matrix Approximation

## Low Rank Approximation with General Losses

$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_g = \sum_{i,j} g((\mathbf{A} - \mathbf{UV})_{i,j})$$

$$\|\mathbf{A} - \hat{\mathbf{U}}\hat{\mathbf{V}}\|_g \leq \kappa \min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_g$$

Subset of  $\approx k$  columns of  $\mathbf{A}$

# Matrix Approximation

## Low Rank Approximation with General Losses

$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_g = \sum_{i,j} g((\mathbf{A} - \mathbf{UV})_{i,j})$$

$$\|\mathbf{A} - \hat{\mathbf{U}}\hat{\mathbf{V}}\|_g \leq \kappa \min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_g$$

Subset of  $\approx k$  columns of  $\mathbf{A}$

Approximation factor  $\kappa$

# Matrix Approximation

## Low Rank Approximation with General Losses

$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_g = \sum_{i,j} g((\mathbf{A} - \mathbf{UV})_{i,j})$$

$$\|\mathbf{A} - \hat{\mathbf{U}}\hat{\mathbf{V}}\|_g \leq \kappa \min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_g$$

Subset of  $\approx k$  columns of  $\mathbf{A}$

Approximation factor  $\kappa$

- $\ell_p, p < 2$ :  $\kappa \approx k^{1/p-1/2}$  (Mahankali—Woodruff 2021)
- $\ell_p, p > 2$ :  $\kappa \approx k^{1-1/p}$  (Dan—Wang—Zhang—Zhou—Ravikumar 2019)



# Matrix Approximation

## Low Rank Approximation with General Losses


$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_g = \sum_{i,j} g((\mathbf{A} - \mathbf{UV})_{i,j})$$

$$\|\mathbf{A} - \hat{\mathbf{U}}\hat{\mathbf{V}}\|_g \leq \kappa \min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_g$$

Subset of  $\approx k$  columns of  $\mathbf{A}$

Approximation factor  $\kappa$

- $\ell_p, p < 2$ :  $\kappa \approx k^{1/p-1/2}$  (Mahankali—Woodruff 2021)
- $\ell_p, p > 2$ :  $\kappa \approx k^{1-1/p}$  (Dan—Wang—Zhang—Zhou—Ravikumar 2019)
- Huber:  $\kappa \approx k^2$  (Song—Woodruff—Zhong 2019)



# Matrix Approximation

## Low Rank Approximation with General Losses

$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_g = \sum_{i,j} g((\mathbf{A} - \mathbf{UV})_{i,j})$$

$$\|\mathbf{A} - \hat{\mathbf{U}}\hat{\mathbf{V}}\|_g \leq \kappa \min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_g$$


Subset of  $\approx k$  columns of  $\mathbf{A}$

Approximation factor  $\kappa$

- $\ell_p, p < 2$ :  $\kappa \approx k^{1/p-1/2}$  (Mahankali—Woodruff 2021)
- $\ell_p, p > 2$ :  $\kappa \approx k^{1-1/p}$  (Dan—Wang—Zhang—Zhou—Ravikumar 2019)

$\kappa \approx k^{1/2-1/p}$  (Woodruff—Y 2023)

- Huber:  $\kappa \approx k^2$  (Song—Woodruff—Zhong 2019)



# Matrix Approximation

## Low Rank Approximation with General Losses

$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_g = \sum_{i,j} g((\mathbf{A} - \mathbf{UV})_{i,j})$$

$$\|\mathbf{A} - \hat{\mathbf{U}}\hat{\mathbf{V}}\|_g \leq \kappa \min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_g$$

Subset of  $\approx k$  columns of  $\mathbf{A}$


Approximation factor  $\kappa$

- $\ell_p, p < 2$ :  $\kappa \approx k^{1/p-1/2}$  (Mahankali—Woodruff 2021)
- $\ell_p, p > 2$ :  $\kappa \approx k^{1-1/p}$  (Dan—Wang—Zhang—Zhou—Ravikumar 2019)

$$\kappa \approx k^{1/2-1/p} \text{ (Woodruff—Y 2023)}$$

- Huber:  $\kappa \approx k^2$  (Song—Woodruff—Zhong 2019)

$$\kappa \approx k \text{ (Woodruff—Y 2023)}$$



# Matrix Approximation

## Low Rank Approximation with General Losses

$$\min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_g = \sum_{i,j} g((\mathbf{A} - \mathbf{UV})_{i,j})$$

$$\|\mathbf{A} - \hat{\mathbf{U}}\hat{\mathbf{V}}\|_g \leq \kappa \min_{\mathbf{U} \in \mathbb{R}^{n \times k}, \mathbf{V} \in \mathbb{R}^{k \times d}} \|\mathbf{A} - \mathbf{UV}\|_g$$

Subset of  $\approx k$  columns of  $\mathbf{A}$


Approximation factor  $\kappa$

- $\ell_p, p < 2$ :  $\kappa \approx k^{1/p-1/2}$  (Mahankali—Woodruff 2021)
- $\ell_p, p > 2$ :  $\kappa \approx k^{1-1/p}$  (Dan—Wang—Zhang—Zhou—Ravikumar 2019)

$$\kappa \approx k^{1/2-1/p} \text{ (Woodruff—Y 2023)}$$

- Huber:  $\kappa \approx k^2$  (Song—Woodruff—Zhong 2019)

$$\kappa \approx k \text{ (Woodruff—Y 2023)}$$



**Techniques:** well-conditioned spanning sets,  $\ell_p$  Lewis weights, ...

# Matrix Approximation

## Summary

# Matrix Approximation

## **Summary**

- We develop a wide variety of techniques for matrix approximation based on sampling and sketching techniques

# Matrix Approximation

## Summary

- We develop a wide variety of techniques for matrix approximation based on sampling and sketching techniques
  - **Tight bounds:** we obtain nearly optimal bounds for fundamental problems for matrix approximation

# Matrix Approximation

## Summary

- We develop a wide variety of techniques for matrix approximation based on sampling and sketching techniques
  - **Tight bounds:** we obtain nearly optimal bounds for fundamental problems for matrix approximation
  - **Uncertain/resource-limited settings:** we develop the first algorithms for matrix approximation in online, streaming, and other models of computation



# Matrix Approximation

## Summary

- We develop a wide variety of techniques for matrix approximation based on sampling and sketching techniques
  - **Tight bounds:** we obtain nearly optimal bounds for fundamental problems for matrix approximation
  - **Uncertain/resource-limited settings:** we develop the first algorithms for matrix approximation in online, streaming, and other models of computation
  - **Generalized loss functions:** we obtain new and improved algorithms for matrix approximation under  $\ell_p$ , Huber, and even further general loss functions

# Matrix Approximation

## Summary

- We develop a wide variety of techniques for matrix approximation based on sampling and sketching techniques
  - **Tight bounds:** we obtain nearly optimal bounds for fundamental problems for matrix approximation
  - **Uncertain/resource-limited settings:** we develop the first algorithms for matrix approximation in online, streaming, and other models of computation
  - **Generalized loss functions:** we obtain new and improved algorithms for matrix approximation under  $\ell_p$ , Huber, and even further general loss functions
  - **Applications:** we apply algorithmic techniques from matrix approximation to solve fundamental problems in machine learning and computational geometry

# Matrix Approximation

## Open Problems

# Matrix Approximation

## Open Problems

**Open Problem 1.** Is there a  $\kappa = O(1)$  distortion subspace embedding for the Huber loss with  $r \approx d$  rows?

# Matrix Approximation

## Open Problems

**Open Problem 1.** Is there a  $\kappa = O(1)$  distortion subspace embedding for the Huber loss with  $r \approx d$  rows?

Current best:  $r \approx d^{4-2\sqrt{2}}$  (Musco—Musco—Woodruff—Y 2022)

# Matrix Approximation

## Open Problems

**Open Problem 1.** Is there a  $\kappa = O(1)$  distortion subspace embedding for the Huber loss with  $r \approx d$  rows?

Current best:  $r \approx d^{4-2\sqrt{2}}$  (Musco—Musco—Woodruff—Y 2022)

**Open Problem 2.** Is  $r \approx \varepsilon^{-2} d^{p/2}$  optimal for  $\kappa = (1 + \varepsilon)$  distortion  $\ell_p$  subspace embeddings for  $p > 2$ ?

# Matrix Approximation

## Open Problems

**Open Problem 1.** Is there a  $\kappa = O(1)$  distortion subspace embedding for the Huber loss with  $r \approx d$  rows?

Current best:  $r \approx d^{4-2\sqrt{2}}$  (Musco—Musco—Woodruff—Y 2022)

**Open Problem 2.** Is  $r \approx \varepsilon^{-2} d^{p/2}$  optimal for  $\kappa = (1 + \varepsilon)$  distortion  $\ell_p$  subspace embeddings for  $p > 2$ ?

Current best:  $r \gtrsim \varepsilon^{-2} + d^{p/2}$  (Li—Wang—Woodruff 2020)

# Matrix Approximation

## Other works

- Subspace Embeddings + Applications:
  - **Sharper Bounds for  $\ell_p$  Sensitivity Sampling [preprint]**
  - New Subset Selection Algorithms for Low Rank Approximation: Offline and Online [STOC'23]
  - Online Lewis Weight Sampling [SODA'23]
  - High-Dimensional Geometric Streaming in Polynomial Space [FOCS'22]
  - Active Linear Regression for  $\ell_p$  Norms and Beyond [FOCS'22]
  - Exponentially Improved Dimensionality Reduction for  $\ell_1$ : Subspace Embeddings and Independence Testing [COLT'21]
- Low Rank Approximation
  - New Subset Selection Algorithms for Low Rank Approximation: Offline and Online [STOC'23]
  - **Improved Algorithms for Low Rank Approximation from Sparsity [SODA'22]**
- **Sequential Attention for Feature Selection [ICLR'23]**