






2-1

問1 (a), (b) は) A, B, C はアルケン C_nH_{2n}







$$1 \text{ mol} \times \frac{3n}{2} = \frac{168 \text{ L}}{22.4 \text{ L/mol}} \therefore n = 5 \quad \underline{C_5H_{10}}$$

問5 C_5H_{10} $U = 1$ $\therefore C=C \times 1$





問1 C_5H_{10}



沸点 $\rightarrow$ 分子間力 $\left\{ \begin{array}{l} \text{水素結合} \\ \text{ファンデルワールス力} \end{array} \right.$
 $\Rightarrow$ ス型の方が極性 $\odot \rightarrow$ 分子間力 $\odot$

2-2





問1 C_6H_{12}



問3 $A \neq C$



$$\frac{2.1 \text{ g}}{14n \text{ g/mol}} = \frac{4.0 \text{ g}}{160 \text{ g/mol}} \quad \therefore n=6 \quad \underline{C_6H_{12}}$$



2-[3]



問1 A $\text{CH}_2=\text{CH}-\text{CH}_2-\text{CH}_3$ 1-ブテン

B $\text{CH}_3-\text{CH}=\text{CH}-\text{CH}_3$ 2-ブテン






C $\text{CH}_3-\text{CH}_2-\text{CH}_2-\text{CH}_3$ ブタン

問2 D: 2 E: 3 (メソ体あり)

問3






2 - 4



2 - 5










(1) 5 (2) 1



2 - 6

161 102

162 C_4H_8





164 F

2 - [7]



問1 略

問2 C C_4H_8O D C_2H_4O


問3



問4.




問5




2 - [8]

問1 アセチレン, γ -ブチレン


問2 (1) A $CH_3-C \equiv C-CH_3$



(2) E $CH_3-\overset{\overset{O}{\parallel}}{C}-CH_2-CH_3$ (イ)



問3



問4.

