

Homework Assignment 1: Slab-Geometry Boltzmann Equation

Tal Ramot

September 5, 2026

Question 1 - Useful identity for the solution of the time-dependent slab-geometry Boltzmann equation

In this question we are asked to prove the following identity:

$$\psi(x, \mu, t; c) = ce^{-(1-c)v\Sigma_t t} \psi(cx, \mu, ct; 1) \quad (1)$$

We will do it in two parts. First, we will write down the equation that $\tilde{\psi} = \psi(cx, \mu, ct; 1)$ satisfies, and then we will use it in order to prove that the product $ce^{-(1-c)v\Sigma_t t} \tilde{\psi}$ satisfies the same equation as $\psi(x, \mu, t; c)$.

The equation for $\tilde{\psi}$

Notation: We will denote by $\Psi(X, \mu, T) \equiv \psi(X, \mu, T; 1)$ the $c = 1$ solution written in its own variables, and by Ψ_T and Ψ_X its partial derivatives with respect to them, so that $\tilde{\psi}(x, \mu, t) = \Psi(cx, \mu, ct)$ with $X = cx$ and $T = ct$. **This notation is used to avoid confusion when using the chain rule, as will be done immediately.** First, Ψ satisfies the $c = 1$ equation in the variables X and T :

$$\frac{1}{v} \Psi_T + \mu \Psi_X + \Sigma_t \Psi = \frac{\Sigma_t}{2} \int_{-1}^1 \Psi d\mu' \quad (2)$$

which after applying the chain rule, $\Psi_T = \frac{1}{c} \frac{\partial \tilde{\psi}}{\partial t}$ and $\Psi_X = \frac{1}{c} \frac{\partial \tilde{\psi}}{\partial x}$, and multiplying by c , becomes:

$$\frac{1}{v} \frac{\partial \tilde{\psi}}{\partial t} + \mu \frac{\partial \tilde{\psi}}{\partial x} + c \Sigma_t \tilde{\psi} = \frac{c \Sigma_t}{2} \int_{-1}^1 \tilde{\psi} d\mu' \quad (3)$$

So $\tilde{\psi}$ obeys the same equation as $\psi(x, \mu, t; c)$ except that it is removed at the rate $c \Sigma_t$ instead of Σ_t .

Proving the identity

In this part we are going to simply substitute the RHS of 1 into the LHS of the equation for $\psi(x, \mu, t; c)$ and show that it satisfies the same equation. After substitution and application of the chain rule we

get:

$$\text{LHS} = \frac{1}{v} \left[c(c-1)v\Sigma_t e^{-(1-c)v\Sigma_t t} \tilde{\psi} + c e^{-(1-c)v\Sigma_t t} \frac{\partial \tilde{\psi}}{\partial t} \right] + \mu c e^{-(1-c)v\Sigma_t t} \frac{\partial \tilde{\psi}}{\partial x} + \Sigma_t c e^{-(1-c)v\Sigma_t t} \tilde{\psi} \quad (4)$$

$$= c e^{-(1-c)v\Sigma_t t} \left[\frac{1}{v} \frac{\partial \tilde{\psi}}{\partial t} + \mu \frac{\partial \tilde{\psi}}{\partial x} + \Sigma_t \tilde{\psi} + (c-1)\Sigma_t \tilde{\psi} \right] \quad (5)$$

$$= c e^{-(1-c)v\Sigma_t t} \left[\frac{1}{v} \frac{\partial \tilde{\psi}}{\partial t} + \mu \frac{\partial \tilde{\psi}}{\partial x} + c\Sigma_t \tilde{\psi} \right] \quad (6)$$

$$= c e^{-(1-c)v\Sigma_t t} \frac{c\Sigma_t}{2} \int_{-1}^1 \tilde{\psi} d\mu' = \frac{c\Sigma_t}{2} \int_{-1}^1 c e^{-(1-c)v\Sigma_t t} \tilde{\psi} d\mu' \quad (7)$$

where 3 was used in the last line, and this is exactly the RHS of the equation for $\psi(x, \mu, t; c)$.

Question 2 - Derivation of the time-dependent planar scalar flux for a plane source

The relation between a plane source and a point source as studied in class is given by:

$$\phi_{\text{pl}}(x) = 2\pi \int_{|x|}^{\infty} \phi_{\text{pt}}(r) r dr \quad (8)$$

The solution taken from J. C. J. Paasschens is the point-source scalar flux $\phi_{\text{pt}}(r, t)$ for $c = 1$, and therefore the planar-source scalar flux is given by the integration given above.

$$\phi_{\text{pl}}(x, t) = 2\pi \int_{|x|}^{\infty} \frac{e^{-v\Sigma_t t}}{4\pi r^2} \delta(r - vt) r dr + 2\pi \int_{|x|}^{\infty} \frac{(1 - r^2/v^2 t^2)^{1/8}}{(4\pi vt/(3\Sigma_t))^{3/2}} e^{-v\Sigma_t t} G\left(v\Sigma_t t \left[1 - \frac{r^2}{v^2 t^2}\right]^{3/4}\right) \Theta(vt - r) r dr \quad (9)$$

$$= \frac{e^{-v\Sigma_t t}}{2vt} \Theta(vt - |x|) + \Theta(vt - |x|) \frac{2\pi e^{-v\Sigma_t t}}{(4\pi vt/(3\Sigma_t))^{3/2}} \int_{|x|}^{vt} \left(1 - \frac{r^2}{v^2 t^2}\right)^{1/8} G\left(v\Sigma_t t \left[1 - \frac{r^2}{v^2 t^2}\right]^{3/4}\right) r dr \quad (10)$$

$$\stackrel{\xi=r/vt}{=} \frac{e^{-v\Sigma_t t}}{2vt} \Theta(vt - |x|) + \Theta(vt - |x|) \frac{2\pi e^{-v\Sigma_t t} (vt)^{1/2}}{(4\pi/(3\Sigma_t))^{3/2}} \int_{\frac{|x|}{vt}}^1 (1 - \xi^2)^{1/8} G\left(a [1 - \xi^2]^{3/4}\right) \xi d\xi \quad (11)$$

$$\stackrel{u=1-\xi^2}{\stackrel{w=au^{3/4}}{=}} \frac{e^{-v\Sigma_t t}}{2vt} \left[1 + \sqrt{\frac{3}{\pi}} \int_0^{w_0} \sqrt{w} G(w) dw \right] \Theta(vt - |x|) \quad (12)$$

with $w_0 = a \left(1 - \frac{x^2}{(vt)^2}\right)^{3/4}$, the powers combining as $\frac{1}{6} + \frac{1}{3} = \frac{1}{2}$ and the Σ_t dependence cancelling in the last step (I used the assistance of AI in this algebraic derivation). Finally, with the interpolation formula of Paasschens, $G(w) = e^w \sqrt{1 + b/w}$ with $b = 2.026$, the integral has a closed form:

$$\int_0^{w_0} e^w \sqrt{w + b} dw = e^{w_0} \sqrt{w_0 + b} - \sqrt{b} - \frac{\sqrt{\pi}}{2} e^{-b} \left[\text{erfi}\left(\sqrt{w_0 + b}\right) - \text{erfi}\left(\sqrt{b}\right) \right] \quad (13)$$

which **closes the derivation of the planar-source scalar flux for $c = 1$** . Of course the relation proved in Q1 can be used to get the solution for $c \neq 1$ using a simple scaling and multiplication by an exponential factor of the solution for $c = 1$.

Question 3 - Classical and asymptotic diffusion against the exact solution

Derivation of the time-dependent diffusion Green's function

The time-dependent diffusion equation is:

$$\frac{1}{v} \frac{\partial \phi}{\partial t} - D \frac{\partial^2 \phi}{\partial x^2} + \Sigma_a \phi = S(x, t) \quad (14)$$

Take $S = \delta(x)\delta(t)$ and multiply (14) by v :

$$\frac{\partial \phi}{\partial t} - Dv \frac{\partial^2 \phi}{\partial x^2} + v \Sigma_a \phi = v \delta(x)\delta(t) \quad (15)$$

Step 1: the pulse source is an initial condition. Integrate (15) over $t \in (0^-, 0^+)$. The diffusion and absorption terms are bounded and contribute nothing in the limit, so only the time derivative survives:

$$\phi(x, 0^+) - \phi(x, 0^-) = v \delta(x) \quad \implies \quad \phi(x, 0^+) = v \delta(x) \quad (16)$$

since nothing precedes the pulse. For $t > 0$ the equation is source-free, so the problem is now a pure initial-value problem (this is the same trick as in Assignment 1, Question 2)

Step 2: absorption factors out exactly. As performed in the course, substituting $\phi(x, t) = e^{-v\Sigma_a t} u(x, t)$ into (15) for $t > 0$, the $-v\Sigma_a u$ produced by the time derivative cancels the absorption term identically and leaves the pure heat equation

$$\frac{\partial u}{\partial t} = Dv \frac{\partial^2 u}{\partial x^2}, \quad u(x, 0^+) = v \delta(x) \quad (17)$$

Note this step is untouched by the sign of $1 - c$: for $c > 1$ the exponential grows instead of decays, and nothing else changes.

Step 3: Fourier transform. With $\hat{u}(k, t) = \int_{-\infty}^{\infty} u(x, t) e^{-ikx} dx$, the spatial derivative becomes multiplication by $-k^2$ and the PDE becomes an ordinary differential equation in t for each k separately:

$$\frac{d\hat{u}}{dt} = -Dvk^2 \hat{u}, \quad \hat{u}(k, 0^+) = v \quad \implies \quad \hat{u}(k, t) = v e^{-Dvk^2 t} \quad (18)$$

the initial value being v because the transform of $v\delta(x)$ is the constant v — a delta excites every wavenumber equally.

Step 4: invert. Completing the square in the exponent, $-Dvk^2 t + ikx = -Dvt \left(k - \frac{ix}{2Dvt}\right)^2 - \frac{x^2}{4Dvt}$, and using $\int_{-\infty}^{\infty} e^{-\alpha k^2} dk = \sqrt{\pi/\alpha}$:

$$u(x, t) = \frac{v}{2\pi} \int_{-\infty}^{\infty} e^{-Dvk^2 t} e^{ikx} dk = \frac{v}{\sqrt{4\pi Dvt}} e^{-x^2/(4Dvt)} \quad (19)$$

Restoring the factor from Step 2 gives the Green's function:

$$\boxed{\phi_{\text{diff}}(x, t; c) = \frac{v e^{-(1-c)\Sigma_a t}}{\sqrt{4\pi Dvt}} \exp\left(-\frac{x^2}{4Dvt}\right)} \quad (20)$$

which in the units of this assignment, $\Sigma_t = v = 1$ and $D = D_0(c)$, is the form implemented in `src/homework2/diffusion.py`:

$$\phi_{\text{diff}}(x, t; c) = \frac{e^{-(1-c)t}}{\sqrt{4\pi D_0(c)t}} e^{-x^2/(4D_0(c)t)} \quad (21)$$

A second route: Fourier in space, then Laplace in time.

Define

$$\hat{\phi}(k, t) = \int_{-\infty}^{\infty} \phi(x, t) e^{-ikx} dx, \quad \tilde{\phi}(k, s) = \int_{0^-}^{\infty} \hat{\phi}(k, t) e^{-st} dt \quad (22)$$

and apply both to (15) term by term, with $\phi(x, 0^-) = 0$:

$$\frac{\partial \phi}{\partial t} \longrightarrow s \tilde{\phi} - \hat{\phi}(k, 0^-) = s \tilde{\phi} \quad (23)$$

$$-Dv \frac{\partial^2 \phi}{\partial x^2} \longrightarrow Dvk^2 \tilde{\phi} \quad (24)$$

$$v \Sigma_a \phi \longrightarrow v \Sigma_a \tilde{\phi} \quad (25)$$

$$v \delta(x) \delta(t) \longrightarrow v \quad (26)$$

so that

$$(s + Dvk^2 + v \Sigma_a) \tilde{\phi}(k, s) = v \quad (27)$$

$$\tilde{\phi}(k, s) = \frac{v}{s + Dvk^2 + v \Sigma_a} \quad (28)$$

a single simple pole at $s = -(Dvk^2 + v \Sigma_a)$, so with $\mathcal{L}^{-1}\{(s + \alpha)^{-1}\} = e^{-\alpha t}$,

$$\hat{\phi}(k, t) = v e^{-(Dvk^2 + v \Sigma_a)t} \quad (29)$$

and the Fourier inversion completes the square exactly as in Step 4:

$$\phi(x, t) = \frac{v}{2\pi} \int_{-\infty}^{\infty} e^{-(Dvk^2 + v \Sigma_a)t} e^{ikx} dk \quad (30)$$

$$= \frac{v e^{-v \Sigma_a t}}{2\pi} e^{-x^2/(4Dvt)} \int_{-\infty}^{\infty} e^{-Dvt(k - \frac{ix}{2Dvt})^2} dk \quad (31)$$

$$= \frac{v e^{-v \Sigma_a t}}{2\pi} e^{-x^2/(4Dvt)} \sqrt{\frac{\pi}{Dvt}} = \frac{v e^{-v \Sigma_a t}}{\sqrt{4\pi Dvt}} e^{-x^2/(4Dvt)} \quad (32)$$

which is (20), with $\Sigma_a = (1 - c)\Sigma_t$.

Parts 3(a) and 3(b): the comparison

Each figure fixes c and shows $t = 1, 2, 3, 4, 7, 15$. The vertical line marks the causal front $|x| = vt$, beyond which the exact solution is identically zero. Only $x \geq 0$ is plotted; all three solutions are even in x .




Figure 1: $c = 0.6$.




Figure 2: $c = 0.8$.




Figure 3: $c = 1$. The two diffusion coefficients coincide at $D_0 = 1/3$, so the classical and asymptotic curves overlie each other.




Figure 4: $c = 1.2$.




Figure 5: $c = 1.5$.