

# Rayleigh Quotients

February 19, 2025

- Quadratic forms

- Quadratic forms
- Positive (semi) definite matrices

- Quadratic forms
- Positive (semi) definite matrices
- Rayleigh quotients

- Quadratic forms
- Positive (semi) definite matrices
- Rayleigh quotients
- Generalized Rayleigh quotients

- Quadratic forms
- Positive (semi) definite matrices
- Rayleigh quotients
- Generalized Rayleigh quotients

# Symmetric Matrices and their Properties

- A symmetric matrix is a square matrices  $A$  satisfying

$$A^T = A$$

.

# Symmetric Matrices and their Properties

- A symmetric matrix is a square matrices  $A$  satisfying

$$A^T = A$$

- Real symmetric matrices  $A \in \mathbb{R}_{n \times n}$  have many nice properties:

.



# Symmetric Matrices and their Properties

- A symmetric matrix is a square matrices  $A$  satisfying

$$A^T = A$$

- Real symmetric matrices  $A \in \mathbb{R}_{n \times n}$  have many nice properties:
  - All their eigenvalues are real numbers (no complex eigenvalues)

# Symmetric Matrices and their Properties

- A symmetric matrix is a square matrices  $A$  satisfying

$$A^T = A$$

- Real symmetric matrices  $A \in \mathbb{R}_{n \times n}$  have many nice properties:

- All their eigenvalues are real numbers (no complex eigenvalues)
- They are orthogonally diagonalizable, i.e., there exist an orthogonal matrix  $Q$  and a diagonal matrix  $\Lambda$ , both of the same size as  $A$ , such that

$$A = Q\Lambda Q^T$$

(This is known as the spectral decomposition of  $A$ .)

# Symmetric Matrices and their Properties

- A symmetric matrix is a square matrices  $A$  satisfying

$$A^T = A$$

- Real symmetric matrices  $A \in \mathbb{R}_{n \times n}$  have many nice properties:

- All their eigenvalues are real numbers (no complex eigenvalues)
- They are orthogonally diagonalizable, i.e., there exist an orthogonal matrix  $Q$  and a diagonal matrix  $\Lambda$ , both of the same size as  $A$ , such that

$$A = Q\Lambda Q^T$$

(This is known as the spectral decomposition of  $A$ .)

The matrix  $\Lambda$  consists of the eigenvalues of  $A$  along the diagonal, and  $Q$  has the corresponding orthonormal eigenvectors in its columns.

# Quadratic Form

**Definition** Let  $A \in \mathbb{R}_{n \times n}$  be a symmetric matrix. A quadratic form corresponding to  $A$  is a function  $Q : \mathbb{R}^n \rightarrow \mathbb{R}$  with

# Quadratic Form

**Definition** Let  $A \in \mathbb{R}_{n \times n}$  be a symmetric matrix. A quadratic form corresponding to  $A$  is a function  $Q : \mathbb{R}^n \rightarrow \mathbb{R}$  with

$$Q(x) = x^T A x, \text{ for all } x \in \mathbb{R}^n.$$

# Quadratic Form

**Definition** Let  $A \in \mathbb{R}_{n \times n}$  be a symmetric matrix. A quadratic form corresponding to  $A$  is a function  $Q : \mathbb{R}^n \rightarrow \mathbb{R}$  with

$$Q(x) = x^T A x, \text{ for all } x \in \mathbb{R}^n.$$

**Remark.** A quadratic form is a polynomial with terms all of second order:

$$x^T A x = \sum_{i=1}^n \sum_{j=1}^n a_{ij} x_i x_j.$$

# Quadratic Form

**Definition** Let  $A \in \mathbb{R}_{n \times n}$  be a symmetric matrix. A quadratic form corresponding to  $A$  is a function  $Q : \mathbb{R}^n \rightarrow \mathbb{R}$  with

$$Q(x) = x^T A x, \text{ for all } x \in \mathbb{R}^n.$$

**Remark.** A quadratic form is a polynomial with terms all of second order:

$$x^T A x = \sum_{i=1}^n \sum_{j=1}^n a_{ij} x_i x_j.$$

**Example.** Let

$$A = \begin{pmatrix} 1 & 3 \\ 3 & 2 \end{pmatrix}, \quad x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix},$$



**Example.** Let

$$A = \begin{pmatrix} 1 & 3 \\ 3 & 2 \end{pmatrix}, \quad x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix},$$

then

$$Q(x) = x^T A x = x_1^2 + 2x_2^2 + 6x_1x_2.$$

**Definition** A symmetric matrix  $A \in \mathbb{R}_{n \times n}$  is said to be positive semidefinite if the corresponding quadratic form

$$Q(x) = x^T A x \geq 0$$

for all  $x \in \mathbb{R}^n$ .

**Definition** A symmetric matrix  $A \in \mathbb{R}_{n \times n}$  is said to be positive semidefinite if the corresponding quadratic form

$$Q(x) = x^T A x \geq 0$$

for all  $x \in \mathbb{R}^n$ .

If the equality holds true only for  $x = 0$  (i.e.,  $x^T A x > 0$  for all  $x \neq 0$ ), then  $A$  is said to be positive definite.

**Theorem** A symmetric matrix is positive definite (semidefinite) if and only if all of its eigenvalues are positive (nonnegative).

**Example** Let

$$A = \begin{pmatrix} 1 & 2 \\ 2 & 5 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix}, \quad C = \begin{pmatrix} 1 & 3 \\ 3 & 2 \end{pmatrix}.$$

According to the theorem,  $A$  is positive definite,  $B$  is positive semidefinite, and  $C$  is neither.

**Problem:** Let  $A \in \mathbb{R}_{n \times n}$  be a positive semi definite matrix. Find another matrix  $B$  of the same size such that  $A = B^2$ . We call  $B$  the **square root** of  $A$  and denote it by  $B = A^{1/2}$ .

**Solution:** Since  $A$  is symmetric and positive semi definite matrix, there exists an orthogonal matrix  $Q \in \mathbb{R}_{n \times n}$  and a diagonal matrix  $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$  with all  $\lambda_i \geq 0$  such that  $A = Q\Lambda Q^T$ .

**Solution:** Since  $A$  is symmetric and positive semi definite matrix, there exists an orthogonal matrix  $Q \in \mathbb{R}_{n \times n}$  and a diagonal matrix  $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$  with all  $\lambda_i \geq 0$  such that  $A = Q\Lambda Q^T$ . Define  $\Lambda^{1/2} = \text{diag}(\lambda_1^{1/2}, \dots, \lambda_n^{1/2})$ . Clearly,  $\Lambda^{1/2}\Lambda^{1/2} = \Lambda$ .

**Solution:** Since  $A$  is symmetric and positive semi definite matrix, there exists an orthogonal matrix  $Q \in \mathbb{R}_{n \times n}$  and a diagonal matrix  $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$  with all  $\lambda_i \geq 0$  such that  $A = Q\Lambda Q^T$ . Define  $\Lambda^{1/2} = \text{diag}(\lambda_1^{1/2}, \dots, \lambda_n^{1/2})$ . Clearly,  $\Lambda^{1/2}\Lambda^{1/2} = \Lambda$ . Let  $B = Q\Lambda^{1/2}Q^T$ . Then

$$B^2 = (Q\Lambda^{1/2}Q^T)(Q\Lambda^{1/2}Q^T) = Q\Lambda^{1/2}\Lambda^{1/2}Q^T = Q\Lambda Q^T = A.$$



**Solution:** Since  $A$  is symmetric and positive semi definite matrix, there exists an orthogonal matrix  $Q \in \mathbb{R}_{n \times n}$  and a diagonal matrix  $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$  with all  $\lambda_i \geq 0$  such that  $A = Q\Lambda Q^T$ . Define  $\Lambda^{1/2} = \text{diag}(\lambda_1^{1/2}, \dots, \lambda_n^{1/2})$ . Clearly,  $\Lambda^{1/2}\Lambda^{1/2} = \Lambda$ . Let  $B = Q\Lambda^{1/2}Q^T$ . Then

$$B^2 = (Q\Lambda^{1/2}Q^T)(Q\Lambda^{1/2}Q^T) = Q\Lambda^{1/2}\Lambda^{1/2}Q^T = Q\Lambda Q^T = A.$$

Thus,  $B = A^{1/2} = Q\Lambda^{1/2}Q^T$  is still a positive semi definite matrix.

**Example** Let

$$A = \begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix},$$

which is positive semi definite matrix because it has two nonnegative eigenvalues  $\lambda_1 = 5, \lambda_2 = 0$ . To find the matrix square root of  $A$ , we need to find its orthogonal diagonalization.

**Example** Let

$$A = \begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix},$$

which is positive semi definite matrix because it has two nonnegative eigenvalues  $\lambda_1 = 5, \lambda_2 = 0$ . To find the matrix square root of  $A$ , we need to find its orthogonal diagonalization.

$$\begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix} = \begin{pmatrix} \frac{1}{\sqrt{5}} & -\frac{2}{\sqrt{5}} \\ \frac{2}{\sqrt{5}} & \frac{1}{\sqrt{5}} \end{pmatrix} \begin{pmatrix} 5 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \frac{1}{\sqrt{5}} & -\frac{2}{\sqrt{5}} \\ \frac{2}{\sqrt{5}} & \frac{1}{\sqrt{5}} \end{pmatrix}^T.$$

**Example** Let

$$A = \begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix},$$

which is positive semi definite matrix because it has two nonnegative eigenvalues  $\lambda_1 = 5$ ,  $\lambda_2 = 0$ . To find the matrix square root of  $A$ , we need to find its orthogonal diagonalization.

$$\begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix} = \begin{pmatrix} \frac{1}{\sqrt{5}} & -\frac{2}{\sqrt{5}} \\ \frac{2}{\sqrt{5}} & \frac{1}{\sqrt{5}} \end{pmatrix} \begin{pmatrix} 5 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \frac{1}{\sqrt{5}} & -\frac{2}{\sqrt{5}} \\ \frac{2}{\sqrt{5}} & \frac{1}{\sqrt{5}} \end{pmatrix}^T.$$

It follows that

$$A^{1/2} = \begin{pmatrix} \frac{1}{\sqrt{5}} & -\frac{2}{\sqrt{5}} \\ \frac{2}{\sqrt{5}} & \frac{1}{\sqrt{5}} \end{pmatrix} \begin{pmatrix} \sqrt{5} & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \frac{1}{\sqrt{5}} & -\frac{2}{\sqrt{5}} \\ \frac{2}{\sqrt{5}} & \frac{1}{\sqrt{5}} \end{pmatrix}^T.$$

**Rayleigh Quotient** Rayleigh quotients are encountered in many statistical and machine learning problems. It is thus necessary to study them systematically.

**Rayleigh Quotient** Rayleigh quotients are encountered in many statistical and machine learning problems. It is thus necessary to study them systematically.

Rayleigh quotients have many applications:

- PCA:  $\max_{v \neq 0} \frac{v^T \Sigma v}{v^T v}$  (where  $\Sigma$  is a covariance matrix)
- LDA:  $\max_{v \neq 0} \frac{v^T S_b v}{v^T S_w v}$  (where  $S_b$  is the between-class scatter matrix and  $S_w$  is the within-class scatter matrix)
- Spectral clustering:  $\max_{v \neq 0} \frac{v^T L v}{v^T D v}$  (where  $L$  is the graph Laplacian and  $D$  is the degree matrix)

**Definition** The Rayleigh quotient for a given symmetric matrix  $A$  is a multivariate function  $f : \mathbb{R}^n - \{0\} \rightarrow \mathbb{R}$  defined by

**Definition** The Rayleigh quotient for a given symmetric matrix  $A$  is a multivariate function  $f : \mathbb{R}^n - \{0\} \rightarrow \mathbb{R}$  defined by

$$f(x) = \frac{x^T A x}{x^T x}, \quad x \neq 0.$$



**Remark.** A Rayleigh quotient is always scaling invariant. That is, for any nonzero vector  $x \in \mathbb{R}^n$ ,

$$f(kx) = \frac{(kx)^T A(kx)}{(kx)^T (kx)} = \frac{x^T A x}{x^T x} = f(x), \quad \text{for all } k \neq 0.$$

**Remark.** A Rayleigh quotient is always scaling invariant. That is, for any nonzero vector  $x \in \mathbb{R}^n$ ,

$$f(kx) = \frac{(kx)^T A(kx)}{(kx)^T (kx)} = \frac{x^T A x}{x^T x} = f(x), \quad \text{for all } k \neq 0.$$

Another way to see this is by rewriting the Rayleigh quotient as follows:

$$f(x) = \frac{x^T A x}{x^T x} = \frac{x^T A x}{\|x\|^2} = \left( \frac{x}{\|x\|} \right)^T A \left( \frac{x}{\|x\|} \right), \quad x \neq 0.$$

**Remark.** A Rayleigh quotient is always scaling invariant. That is, for any nonzero vector  $x \in \mathbb{R}^n$ ,

$$f(kx) = \frac{(kx)^T A(kx)}{(kx)^T (kx)} = \frac{x^T A x}{x^T x} = f(x), \quad \text{for all } k \neq 0.$$

Another way to see this is by rewriting the Rayleigh quotient as follows:

$$f(x) = \frac{x^T A x}{x^T x} = \frac{x^T A x}{\|x\|^2} = \left( \frac{x}{\|x\|} \right)^T A \left( \frac{x}{\|x\|} \right), \quad x \neq 0.$$

**Interpretation:** The Rayleigh quotient is essentially a quadratic form over the unit sphere.

**Example** The Rayleigh quotient for

$$A = \begin{pmatrix} 1 & 3 \\ 3 & 2 \end{pmatrix}$$

**Example** The Rayleigh quotient for

$$A = \begin{pmatrix} 1 & 3 \\ 3 & 2 \end{pmatrix}$$

is

$$f(x) = \frac{x^T A x}{x^T x} = \frac{x_1^2 + 2x_2^2 + 6x_1x_2}{x_1^2 + x_2^2}, \quad x \neq 0.$$

**Example** The Rayleigh quotient for

$$A = \begin{pmatrix} 1 & 3 \\ 3 & 2 \end{pmatrix}$$

is

$$f(x) = \frac{x^T A x}{x^T x} = \frac{x_1^2 + 2x_2^2 + 6x_1x_2}{x_1^2 + x_2^2}, \quad x \neq 0.$$

It is a function defined over  $\mathbb{R}^2$  with the origin excluded.

**Problem:** Given symmetric matrix  $A$ , find the maximum (or minimum) of the associated Rayleigh quotient

$$\max_{x \neq 0 \in \mathbb{R}^n} \frac{x^T A x}{x^T x}.$$

**Problem:** Given symmetric matrix  $A$ , find the maximum (or minimum) of the associated Rayleigh quotient

$$\max_{x \neq 0 \in \mathbb{R}^n} \frac{x^T A x}{x^T x}.$$

Equivalent formulations:

$$\max_{x \in \mathbb{R}^n, \|x\|=1} x^T A x,$$

$$\max_{x \in \mathbb{R}^n} x^T A x \quad \text{subject to} \quad \|x\|^2 = 1.$$



**Theorem** For any given symmetric matrix  $A \in \mathbb{R}_{n \times n}$ ,

$$\max_{x \in \mathbb{R}^n, x \neq 0} \frac{x^T A x}{x^T x} = \lambda_{\max},$$

**Theorem** For any given symmetric matrix  $A \in \mathbb{R}_{n \times n}$ ,

$$\max_{x \in \mathbb{R}^n, x \neq 0} \frac{x^T A x}{x^T x} = \lambda_{\max},$$

$$\min_{x \in \mathbb{R}^n, x \neq 0} \frac{x^T A x}{x^T x} = \lambda_{\min}.$$

$x$  is the eigenvector associated with the largest and smallest eigenvalue of  $A$  respectively.

**Proof:** Let  $A = V\Lambda V^T$  be the spectral decomposition, where  $V = [v_1, \dots, v_n]$  is orthogonal and  $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$  is diagonal with sorted diagonals from large to small. Then for any unit vector  $x$ ,

**Proof:** Let  $A = V\Lambda V^T$  be the spectral decomposition, where  $V = [v_1, \dots, v_n]$  is orthogonal and  $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$  is diagonal with sorted diagonals from large to small. Then for any unit vector  $x$ ,

$$x^T A x = x^T (V\Lambda V^T) x = (x^T V) \Lambda (V^T x) = y^T \Lambda y,$$

where  $y = V^T x$  is also a unit vector

**Proof:** Let  $A = V\Lambda V^T$  be the spectral decomposition, where  $V = [v_1, \dots, v_n]$  is orthogonal and  $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$  is diagonal with sorted diagonals from large to small. Then for any unit vector  $x$ ,

$$x^T A x = x^T (V\Lambda V^T) x = (x^T V) \Lambda (V^T x) = y^T \Lambda y,$$

where  $y = V^T x$  is also a unit vector

$$\|y\|^2 = y^T y = (V^T x)^T (V^T x) = x^T V V^T x = x^T x = 1.$$

**Proof:** Let  $A = V\Lambda V^T$  be the spectral decomposition, where  $V = [v_1, \dots, v_n]$  is orthogonal and  $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$  is diagonal with sorted diagonals from large to small. Then for any unit vector  $x$ ,

$$x^T A x = x^T (V\Lambda V^T) x = (x^T V) \Lambda (V^T x) = y^T \Lambda y,$$

where  $y = V^T x$  is also a unit vector

$$\|y\|^2 = y^T y = (V^T x)^T (V^T x) = x^T V V^T x = x^T x = 1.$$

So the original optimization problem becomes:

$$\max_{y \in \mathbb{R}^n, \|y\|=1} y^T \Lambda y.$$

$$y^T \Lambda y = \sum_{i=1}^n \lambda_i y_i^2 \quad \text{subject to} \quad y_1^2 + y_2^2 + \cdots + y_n^2 = 1.$$

$$y^T \Lambda y = \sum_{i=1}^n \lambda_i y_i^2 \quad \text{subject to} \quad y_1^2 + y_2^2 + \cdots + y_n^2 = 1.$$

Then clearly

$$\sum_{i=1}^n \lambda_i |y_i|^2 \leq \lambda_n \sum_{i=1}^n |y_i|^2 = \lambda_n \|y\|^2$$



$$y^T \Lambda y = \sum_{i=1}^n \lambda_i y_i^2 \quad \text{subject to} \quad y_1^2 + y_2^2 + \cdots + y_n^2 = 1.$$

Then clearly

$$\sum_{i=1}^n \lambda_i |y_i|^2 \leq \lambda_n \sum_{i=1}^n |y_i|^2 = \lambda_n \|y\|^2$$

and

$$\sum_{i=1}^n \lambda_i |y_i|^2 \geq \lambda_1 \sum_{i=1}^n |y_i|^2 = \lambda_1 \|y\|^2$$

$$y^T \Lambda y = \sum_{i=1}^n \lambda_i y_i^2 \quad \text{subject to} \quad y_1^2 + y_2^2 + \cdots + y_n^2 = 1.$$

Then clearly

$$\sum_{i=1}^n \lambda_i |y_i|^2 \leq \lambda_n \sum_{i=1}^n |y_i|^2 = \lambda_n \|y\|^2$$

and

$$\sum_{i=1}^n \lambda_i |y_i|^2 \geq \lambda_1 \sum_{i=1}^n |y_i|^2 = \lambda_1 \|y\|^2$$

therefore

$$\lambda_1 \leq R(x) \leq \lambda_n \quad \text{for all } x \neq 0.$$

**Example** For the semi positive definite matrix

$$A = \begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix},$$

**Example** For the semi positive definite matrix

$$A = \begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix},$$

we have previously obtained its eigenvalues and eigenvectors:

$$\lambda_1 = 5, \lambda_2 = 0; v_1 = \frac{1}{\sqrt{5}} \begin{pmatrix} 1 \\ 2 \end{pmatrix}, v_2 = \frac{1}{\sqrt{5}} \begin{pmatrix} -2 \\ 1 \end{pmatrix}.$$

The associated Rayleigh quotient

$$Q(x) = \frac{x_1^2 + 4x_2^2 + 4x_1x_2}{x_1^2 + x_2^2}$$

The associated Rayleigh quotient

$$Q(x) = \frac{x_1^2 + 4x_2^2 + 4x_1x_2}{x_1^2 + x_2^2}$$

has the following extreme values:

- The maximum value of  $Q(x)$  is  $\lambda_1 = 5$ , achieved at  $x = \pm v_1$ ;
- The minimum is  $\lambda_2 = 0$ , achieved at  $x = \pm v_2$ .

The overall range of the Rayleigh quotient is thus  $[0, 5]$ .