

# Ovarian Cancer Histotypes: Report of Statistical Findings

Derek Chiu

August 10, 2026

# Table of contents

|                                                        |           |
|--------------------------------------------------------|-----------|
| <b>Preface</b>                                         | <b>7</b>  |
| <b>1 Introduction</b>                                  | <b>8</b>  |
| <b>2 Methods</b>                                       | <b>9</b>  |
| 2.1 Pre-Processing . . . . .                           | 9         |
| 2.1.1 Case Selection . . . . .                         | 9         |
| 2.1.2 Quality Control . . . . .                        | 9         |
| 2.1.3 Housekeeping Genes Normalization . . . . .       | 10        |
| 2.1.4 Between CodeSet and Site Normalization . . . . . | 10        |
| 2.1.5 Final Processing . . . . .                       | 11        |
| 2.2 Classifiers . . . . .                              | 12        |
| 2.2.1 Resampling of Training Set . . . . .             | 13        |
| 2.2.2 Hyperparameter Tuning . . . . .                  | 13        |
| 2.2.3 Subsampling . . . . .                            | 14        |
| 2.2.4 Workflows . . . . .                              | 14        |
| 2.3 Two-Step Algorithm . . . . .                       | 15        |
| 2.3.1 Aggregating Predictions . . . . .                | 15        |
| 2.4 Sequential Algorithm . . . . .                     | 16        |
| 2.4.1 Aggregating Predictions . . . . .                | 17        |
| 2.5 Performance Evaluation . . . . .                   | 18        |
| 2.5.1 Class Metrics . . . . .                          | 18        |
| 2.5.2 AUC . . . . .                                    | 20        |
| 2.6 Rank Aggregation . . . . .                         | 20        |
| 2.7 Gene Optimization . . . . .                        | 20        |
| 2.7.1 Variable Importance . . . . .                    | 22        |
| <b>3 Distributions</b>                                 | <b>24</b> |
| 3.1 Histotype Distribution . . . . .                   | 24        |
| 3.2 Cohort Distribution . . . . .                      | 26        |
| 3.3 Tissue Source . . . . .                            | 26        |
| 3.4 Quality Control . . . . .                          | 27        |
| 3.4.1 Failed Samples . . . . .                         | 27        |
| 3.4.2 %GD vs. SNR . . . . .                            | 28        |
| 3.5 Pairwise Gene Expression . . . . .                 | 30        |
| <b>4 Results</b>                                       | <b>34</b> |
| 4.1 Training Set . . . . .                             | 35        |
| 4.1.1 Accuracy . . . . .                               | 35        |
| 4.1.2 Sensitivity . . . . .                            | 37        |
| 4.1.3 Specificity . . . . .                            | 39        |

|       |                                                     |    |
|-------|-----------------------------------------------------|----|
| 4.1.4 | F1-Score . . . . .                                  | 41 |
| 4.1.5 | Balanced Accuracy . . . . .                         | 43 |
| 4.1.6 | Kappa . . . . .                                     | 45 |
| 4.2   | Rank Aggregation . . . . .                          | 46 |
| 4.2.1 | Across Classes . . . . .                            | 47 |
| 4.2.2 | Across Metrics . . . . .                            | 50 |
| 4.2.3 | Top Workflows . . . . .                             | 50 |
| 4.3   | Confirmation Set . . . . .                          | 54 |
| 4.3.1 | Sequential . . . . .                                | 59 |
| 4.3.2 | 2-STEP . . . . .                                    | 60 |
| 4.3.3 | Up-RF . . . . .                                     | 61 |
| 4.3.4 | SMOTE-SVM . . . . .                                 | 62 |
| 4.3.5 | Hybrid-XGB . . . . .                                | 63 |
| 4.4   | Gene Optimization . . . . .                         | 63 |
| 4.4.1 | Hybrid-XGB . . . . .                                | 64 |
| 4.4.2 | SMOTE-SVM . . . . .                                 | 67 |
| 4.4.3 | Gene List Comparisons in Confirmation Set . . . . . | 70 |
| 4.5   | Validation Set . . . . .                            | 72 |
| 4.5.1 | Evaluation Metrics . . . . .                        | 73 |
| 4.5.2 | Confusion Matrices . . . . .                        | 74 |
| 4.5.3 | ROC Curves . . . . .                                | 75 |
| 4.5.4 | Calibration Plots . . . . .                         | 78 |
| 4.5.5 | Final Model . . . . .                               | 80 |
| 4.5.6 | Additional Explorations . . . . .                   | 88 |

|                   |           |
|-------------------|-----------|
| <b>References</b> | <b>92</b> |
|-------------------|-----------|

# List of Figures

|      |                                                                                  |    |
|------|----------------------------------------------------------------------------------|----|
| 2.1  | Venn diagram of common and unique gene targets covered by each CodeSet . . . . . | 11 |
| 2.2  | Cohorts Selection . . . . .                                                      | 12 |
| 2.3  | Visualization of Subsampling Techniques . . . . .                                | 14 |
| 2.4  | Two-Step Algorithm . . . . .                                                     | 15 |
| 2.5  | Aggregating Predictions for Two-Step Algorithm . . . . .                         | 16 |
| 2.6  | Sequential Algorithm . . . . .                                                   | 17 |
| 2.7  | Aggregating Predictions for Sequential Algorithm . . . . .                       | 18 |
| 3.1  | % Genes Detected vs. Signal to Noise Ratio . . . . .                             | 28 |
| 3.2  | % Genes Detected vs. Signal to Noise Ratio (Zoomed) . . . . .                    | 29 |
| 3.3  | Random1-Normalized CS1 vs. CS3 Gene Expression . . . . .                         | 30 |
| 3.4  | Random1-Normalized CS2 vs. CS3 Gene Expression . . . . .                         | 31 |
| 3.5  | HKgenes-Normalized CS1 vs. CS3 Gene Expression . . . . .                         | 32 |
| 3.6  | HKgenes-Normalized CS2 vs. CS3 Gene Expression . . . . .                         | 33 |
| 4.1  | Training Set Mean Accuracy . . . . .                                             | 36 |
| 4.2  | Training Set Mean Sensitivity . . . . .                                          | 38 |
| 4.3  | Training Set Mean Specificity . . . . .                                          | 40 |
| 4.4  | Training Set Mean F1-Score . . . . .                                             | 42 |
| 4.5  | Training Set Mean Balanced Accuracy . . . . .                                    | 44 |
| 4.6  | Training Set Mean Kappa . . . . .                                                | 46 |
| 4.7  | Top 5 Workflow Per-Class Evaluation Metrics by Metric . . . . .                  | 52 |
| 4.8  | Top 5 Workflow Per-Class Evaluation Metrics by Metric . . . . .                  | 53 |
| 4.9  | Evaluation Metrics on Confirmation Set Models . . . . .                          | 56 |
| 4.10 | Entropy vs. Predicted Probability in Confirmation Set . . . . .                  | 56 |
| 4.11 | Gene Optimized Workflows Per-Class Metrics in Confirmation Set . . . . .         | 57 |
| 4.12 | Confusion Matrices for Confirmation Set Models . . . . .                         | 58 |
| 4.13 | ROC Curves for Sequential Model in Confirmation Set . . . . .                    | 59 |
| 4.14 | ROC Curves for 2-STEP Model in Confirmation Set . . . . .                        | 60 |
| 4.15 | ROC Curve for Up-RF Model in Confirmation Set . . . . .                          | 61 |
| 4.16 | ROC Curve for SMOTE-SVM Model in Confirmation Set . . . . .                      | 62 |
| 4.17 | ROC Curve for Hybrid-XGB Model in Confirmation Set . . . . .                     | 63 |
| 4.18 | Gene Optimization for Hybrid-XGB Classifier using Averaged F1-Score . . . . .    | 64 |
| 4.19 | Gene Optimization for SMOTE-SVM Classifier using Averaged F1-Score . . . . .     | 67 |
| 4.20 | Gene List Comparisons of Evaluation Metrics in Confirmation Set . . . . .        | 72 |
| 4.21 | Evaluation Metrics on Validation Set Models . . . . .                            | 73 |
| 4.22 | Confusion Matrix for Training Set Models evaluated on Validation Data . . . . .  | 74 |
| 4.23 | ROC Curves for Hybrid-XGB, All Overlap Set Model in Validation Set . . . . .     | 75 |
| 4.24 | ROC Curves for Hybrid-XGB, Optimal Set Model in Validation Set . . . . .         | 76 |
| 4.25 | ROC Curves for Hybrid-XGB, Core Set Model in Validation Set . . . . .            | 77 |

|                                                                                          |    |
|------------------------------------------------------------------------------------------|----|
| 4.26 Calibration Plots for Hybrid-XGB, All Overlap Set Model in Validation Set . . . . . | 78 |
| 4.27 Calibration Plots for Hybrid-XGB, Optimal Set Model in Validation Set . . . . .     | 79 |
| 4.28 Calibration Plots for Hybrid-XGB, Core Set Model in Validation Set . . . . .        | 80 |
| 4.29 Validation Summary . . . . .                                                        | 81 |
| 4.30 Final Model Variable Importance Plot . . . . .                                      | 82 |
| 4.31 Final Model Variable Importance Plot by Histotype . . . . .                         | 83 |
| 4.32 Final Model SHAP Importance Plot . . . . .                                          | 84 |
| 4.33 SHAP Dependence Plot for HNF1B . . . . .                                            | 85 |
| 4.34 SHAP Dependence Plot for TFF3 . . . . .                                             | 86 |
| 4.35 Volcano Plots of Validation Set Predictions . . . . .                               | 89 |
| 4.36 Boxplot of Most Differentially Expressed Genes . . . . .                            | 90 |
| 4.37 Subtype Prediction Summary among Predicted HGSOc Samples . . . . .                  | 91 |

# List of Tables

|      |                                                                                     |    |
|------|-------------------------------------------------------------------------------------|----|
| 2.1  | Gene Distribution . . . . .                                                         | 21 |
| 3.1  | Histotype Distribution in Training Set by Processing Stage . . . . .                | 24 |
| 3.2  | Histotype Distribution in Training, Confirmation, and Validation Sets . . . . .     | 25 |
| 3.3  | Pre-QC Cohort Distribution by CodeSet . . . . .                                     | 26 |
| 3.4  | Tissue Source by Cohort . . . . .                                                   | 26 |
| 3.5  | Quality Control Summary . . . . .                                                   | 27 |
| 3.6  | Wilcoxon signed rank test of gene correlations between normalization methods . . .  | 33 |
| 4.1  | Training Set Mean Accuracy . . . . .                                                | 35 |
| 4.2  | Training Set Mean Sensitivity . . . . .                                             | 37 |
| 4.3  | Training Set Mean Specificity . . . . .                                             | 39 |
| 4.4  | Training Set Mean F1-Score . . . . .                                                | 41 |
| 4.5  | Training Set Mean Balanced Accuracy . . . . .                                       | 43 |
| 4.6  | Training Set Mean Kappa . . . . .                                                   | 45 |
| 4.7  | F1-Score Rank Aggregation Summary . . . . .                                         | 47 |
| 4.8  | Balanced Accuracy Rank Aggregation Summary . . . . .                                | 48 |
| 4.9  | Kappa Rank Aggregation Summary . . . . .                                            | 49 |
| 4.10 | Rank Aggregation Comparison of Metrics Used in Training Set . . . . .               | 50 |
| 4.11 | Top 5 Workflows from Final Rank Aggregation . . . . .                               | 50 |
| 4.12 | Top Workflow Per-Class Evaluation Metrics . . . . .                                 | 51 |
| 4.13 | Top Workflow Per-Class Evaluation Metrics and Ranks . . . . .                       | 53 |
| 4.14 | Evaluation Metrics on Confirmation Set Models . . . . .                             | 55 |
| 4.15 | Gene Profile of Optimal Set in Hybrid-XGB Workflow . . . . .                        | 64 |
| 4.16 | Gene Profile of Optimal Set in SMOTE-SVM Workflow . . . . .                         | 67 |
| 4.17 | Model Comparisons using Different Gene Lists in Confirmation Set . . . . .          | 71 |
| 4.18 | Evaluation Metrics on Training Set Models in Validation Set . . . . .               | 73 |
| 4.19 | Detailed Gene Membership . . . . .                                                  | 87 |
| 4.20 | Clinicopath characteristics between correct and incorrect predictions of ENOC cases | 88 |

# Preface

This report of statistical findings describes the classification of ovarian cancer histotypes using data from NanoString CodeSets.

Marina Pavanello conducted the initial exploratory data analysis, Cathy Tang implemented class imbalance techniques, Derek Chiu conducted the normalization and statistical analysis, and Lauren Tindale and Aline Talhouk are the project leads.

# 1 Introduction

Ovarian cancer has five major histotypes: high-grade serous carcinoma (HGSOC), low-grade serous carcinoma (LGSOC), endometrioid carcinoma (ENOC), mucinous carcinoma (MUOC), and clear cell carcinoma (CCOC). A common problem with classifying these histotypes is that there is a class imbalance issue. HGSOC dominates the distribution, commonly accounting for 70% of cases in many patient cohorts, while the other four histotypes are spread over the rest of the cases. Subsampling methods like up-sampling, down-sampling, and SMOTE can be used to mitigate this problem.

The supervised learning is performed under a consensus framework: we consider various classification algorithms and use evaluation metrics like accuracy, F1-score, and Kappa, to inform the decision of which methods to carry forward for prediction in confirmation and validation sets.

## 2 Methods

### 2.1 Pre-Processing

#### 2.1.1 Case Selection

Prior to pre-processing, samples were split into a training, a confirmation, and a validation set.

- Training
  - CS1: OOU, OOUE, VOA, MAYO, MTL
  - CS2: OOU, OOUE, VOA, MAYO, OVAR3, OVAR11, JAPAN, MTL, POOL-CTRL
  - CS3: OOU, OOUE, VOA, POOL-1, POOL-2, POOL-3
- Confirmation:
  - CS3: TNCO
- Validation:
  - CS3: DOVE4

#### 2.1.2 Quality Control

Before normalization, we calculated several quality control measures and excluded samples that failed to achieve sample quality in one or more of these measures.

- **Linearity of positive control genes:** If the R-squared from a linear model of positive controls and their concentrations is less than 0.95 or missing, then the sample is flagged.
- **Imaging quality:** The sample is flagged if the field of view percentage is less than 75%.
- **Positive Control flag:** We consider the two smallest positive controls at concentrations 0.5 and 1. If these two controls are less than the lower limit of detection (defined as two standard deviations below the mean of the negative control expression), or if the mean negative control expression is 0, the sample is flagged.
- **The signal-to-noise ratio or percent of genes detected:** These two measures are defined as the ratio of the average housekeeping gene expression over the upper limit of detection, defined as two standard deviations above the mean of the negative control expression (or 0 if this limit is less than 0.001), and the proportion of endogenous genes with expression greater than the upper limit of detection. These measures are flagged if they are below a pre-specified threshold, which is determined visually by considering their bivariate distribution in a scatterplot. In this case, we used 100 for the SNR threshold and 50% for the threshold for genes detected. Note: these thresholds were determined by examining the relationship in [Section 3.4.2](#).

### 2.1.3 Housekeeping Genes Normalization

The full training set (n=1257) comprised of data from three CodeSets (CS) 1, 2, and 3. Data normalization removes technical variation from high-throughput platforms to improve the validity of comparative analyses.

Each CodeSet was first normalized to housekeeping genes: *ACTB*, *RPL19*, *POLR1B*, *SDHA*, and *PGK1*. Housekeeping genes encode proteins responsible for basic cell function and have consistent expression in all cells. All expression values were log2 transformed. Normalization to housekeeping genes corrects the viable RNA from each sample. This is achieved by subtracting the average log (base 2)-transformed expression of the housekeeping genes from the log (base 2)-transformed expression of each gene:

$$\log_2(\text{endogenous gene expression}) - \text{average}(\log_2(\text{housekeeping gene expression})) = \text{relative expression} \quad (2.1)$$

### 2.1.4 Between CodeSet and Site Normalization

To normalize between CodeSets, we randomly selected five specimens, one from each histotype, among specimens repeated in all three CodeSets. This formed the reference set (Random 1). We selected only one sample from each histotype to use as few samples as possible for normalization and retain the rest for analysis.

A reference-based approach (Talhouk et al. (2016)) was used to normalize CS1 to CS3 and CS2 to CS3 across their common genes:

$$\begin{aligned} \text{X-Norm}_{\text{CS1}} &= X_{\text{CS1}} + \bar{R}_{\text{CS3}} - \bar{R}_{\text{CS1}} \\ \text{X-Norm}_{\text{CS2}} &= X_{\text{CS2}} + \bar{R}_{\text{CS3}} - \bar{R}_{\text{CS2}} \end{aligned} \quad (2.2)$$

Samples in CS3 were processed at three different locations; we also had to normalize for “site” in this CodeSet. Finally, the CS3 expression samples were included in the training set without further normalization:

$$\begin{aligned} \text{X-Norm}_{\text{CS3-USC}} &= X_{\text{CS3-USC}} + \bar{R}_{\text{CS3-VAN}} - \bar{R}_{\text{CS3-USC}} \\ \text{X-Norm}_{\text{CS3-AOC}} &= X_{\text{CS3-AOC}} + \bar{R}_{\text{CS3-VAN}} - \bar{R}_{\text{CS3-AOC}} \end{aligned} \quad (2.3)$$

The CS3 expression samples were included in the training set without further normalization. CS3 cross-site replicates were all chosen from Vancouver, thus we did not retain any samples from USC or AOC. Finally, the initial training set is assembled by combining the normalized datasets for CS1 and CS2 along with the CS3 expression not used in normalization:

$$\begin{aligned} \text{Training Set} &= \text{X-Norm}_{\text{CS1}} + \text{X-Norm}_{\text{CS2}} + X_{\text{CS3-VAN}} + \text{X-Norm}_{\text{CS3-USC}} + \text{X-Norm}_{\text{CS3-AOC}} \\ &= \text{X-Norm}_{\text{CS1}} + \text{X-Norm}_{\text{CS2}} + X_{\text{CS3}} \end{aligned} \quad (2.4)$$




Figure 2.1: Venn diagram of common and unique gene targets covered by each CodeSet

### 2.1.5 Final Processing

We map ovarian histotypes to all remaining samples and keep the major histotypes for building the predictive model: high-grade serous ovarian carcinoma (HGSOC), clear cell ovarian carcinoma (CCOC), endometrioid ovarian carcinoma (ENOC), mucinous ovarian carcinoma (MUOC), and low-grade serous ovarian carcinoma (LGSOC).

Duplicate cases (two samples with the same ottaID) were removed before generating the final training set to use for fitting the classification models. All CS3 cases were preferred over CS1

and CS2, and CS3-Vancouver cases were preferred over CS3-AOC and CS3-USC when selecting duplicates.

The final training set used only genes that were common across all three CodeSets.

Figure 2.2: Cohorts Selection

## 2.2 Classifiers

We used a supervised learning framework to train four multinomial classification algorithms in the Training Set. The full prediction model pipeline was run using SLURM batch jobs submitted to a partition on a Rocky 9 server. All resampling techniques, pre-processing, model specification, hyperparameter tuning, and evaluation metrics were implemented using the tidymodels suite of packages in R (v4.5.2). The classifiers we used are:

- Random Forest (`rf`) ([Wright and Ziegler 2017](#))
- Support Vector Machine (`svm`) ([Platt et al. 1999](#))
- XGBoost (`xgb`) ([Chen and Guestrin 2016](#))
- Regularized Multinomial Regression (`mr`) ([Hastie et al. 2015](#))

### 2.2.1 Resampling of Training Set

We used a nested cross-validation design to evaluate each classifier and perform hyperparameter tuning. An outer 5-fold CV stratified by histotype, along with an inner 5-fold CV with two repeats also stratified by histotype, ensured that the test sets of the inner dataset contained a reasonable number of cases from the smallest minority class.

The hyperparameter combination that yielded the highest average F1 score across the inner training sets was selected for assessing prediction performance in the outer training loop. We avoided using the bootstrap method for outer resampling to prevent overlap between the training and test sets caused by sampling with replacement. This overlap could lead to inflated performance estimates, as some observations would be used for both training and evaluation in the inner loop.

### 2.2.2 Hyperparameter Tuning

The following specifications for each classifier were used for tuning hyperparameters:

- **rf**: The number of trees was fixed at 500. Two other hyperparameters were tuned across 10 randomly selected points in a space-filling parameter grid design:
  - `mtry`: number of randomly selected predictors
  - `min_n`: minimal node size
- **xgb**: The number of trees was fixed at 500. Seven other hyperparameters were tuned across 10 randomly selected points in a space-filling parameter grid design:
  - `mtry`: number of randomly selected predictors
  - `min_n`: minimal node size
  - `tree_depth`: maximum depth of the tree (splits)
  - `learn_rate`: rate at which the boosting algorithm adapts from iteration-to-iteration (shrinkage parameter)
  - `loss_reduction`: reduction in the loss function required to split further
  - `sample_size`: proportion of observations exposed to fitting routine
  - `stop_iter`: number of iterations without improvement before stopping
- **svm**: The cost and sigma hyperparameters were tuned across 10 randomly selected points in a space-filling parameter grid design. We tuned the cost parameter in the range [1, 8]. The range for tuning the sigma parameter was obtained from the 10% and 90% quantiles of the estimates from the `kernlab::sigest()` function.
- **mr**: The penalty (lambda) and mixture (alpha) hyperparameters were tuned across 100 randomly selected points in a space-filling parameter grid design.

The hyperparameter combination that resulted in the highest average F1-score across the inner training sets was selected for each classifier to use as the model for assessing prediction performance in the outer training loop.

### 2.2.3 Subsampling

One of the main concerns in the analysis is that the distribution of the five main histotypes is highly imbalanced, which has implications for the accuracy of the classifier. To mitigate this, we investigated several strategies based on subsampling approaches:

- **None:** No subsampling is performed
- **Down-sampling:** All levels except the minority class are sampled down to the same frequency as the minority class ([Hvitfeldt 2025](#))
- **Up-sampling:** All levels except the majority class are sampled up to the same frequency as the majority class ([Hvitfeldt 2025](#))
- **SMOTE:** All levels except the majority class have synthetic data generated until they have the same frequency as the majority class ([Chawla et al. 2002](#))
- **Hybrid:** All levels except the majority class have synthetic data generated up to 50% of the frequency of the majority class (i.e. SMOTE with 0.5 ratio of minority to majority), then the majority class is down-sampled to the same frequency as the rest.

Figure 2.3 helps visualize how the distribution of classes changes when we apply subsampling techniques to handle class imbalance:




Figure 2.3: Visualization of Subsampling Techniques

All subsampling methods to mitigate class imbalance were applied within the inner CV loop.

### 2.2.4 Workflows

The four **algorithms** and five **subsampling** methods created 20 unique classification **workflows**. For example, the `hybrid_xgb` workflow pre-processes the training set by applying a hybrid sub-

sampling method and subsequently uses the XGBoost algorithm to classify ovarian histotypes. We additionally tested two ensemble algorithms, the two-step, and the sequential algorithms described in the following sections.

## 2.3 Two-Step Algorithm

The HGSOC histotype comprises most cases among ovarian carcinoma patients, while the remaining patients are uniformly split across ENOC, CCOC, MUOC, and LGSOC histotypes. We can implement a two-step algorithm as such:

- Step 1: Select the best classifier to predict HGSOC vs. non-HGSOC, using the workflow with the best per-class F1-score for HGSOC when fit to a 5-class prediction model
- Step 2: remove HGSOCs from the dataset and re-train a 4-class multinomial classifier to predict the four remaining non-HGSOC histotypes. Select the workflow with the best macro-averaged F1-score when fit to a 4-class prediction model.

Let

$$\begin{aligned}
 X_k &= \text{Training data with } k \text{ classes} \\
 C_k &= \text{Class with highest } F_1 \text{ score from training } X_k \\
 W_k &= \text{Workflow associated with } C_k
 \end{aligned} \tag{2.5}$$

Figure 2.4 shows how the two-step algorithm works:




Figure 2.4: Two-Step Algorithm

### 2.3.1 Aggregating Predictions

The aggregation for two-step predictions is quite straightforward:

1. Predict HGSOC vs. non-HGSOC

2. Among all non-HGSOC cases, predict CCOC vs. LGSOC vs. MUOC vs. ENOC




Figure 2.5: Aggregating Predictions for Two-Step Algorithm

## 2.4 Sequential Algorithm

Instead of training on  $k$  classes simultaneously using multinomial classifiers, we can use a sequential algorithm that iteratively performs one-vs-all binary classifications  $k-1$  times to obtain a final prediction of all cases. At each step in the sequence, we classify one class vs. all other classes, where the classes that make up the “other” class are those not equal to the current “one” class and exclude all “one” classes from previous steps. For example, if the “one” class in step 1 was HGSOC, the “other” classes would include CCOC, ENOC, LGSOC, and MUOC. If the “one” class in step 2 was CCOC, the “other” classes would include ENOC, LGSOC, and MUOC.

The order of classes and workflows to use at each step in the sequential algorithm is determined after removing the data associated with an already classified class and retraining using the remaining data multinomial classifiers, as described before. The class and workflow to use for the next step in the sequence are selected based on the best per-class evaluation metric value (e.g., F1-score).

Figure 2.6 illustrates how the sequential algorithm works for  $K=5$ , using ovarian histotypes as an example for the classes.




Figure 2.6: Sequential Algorithm

### 2.4.1 Aggregating Predictions

We have to aggregate the one-vs-all predictions from each of the sequential algorithm workflows in order to obtain a final class prediction on a holdout test set. Each sequential workflow has to be assessed on every sample to ensure that cases classified into the “other” class from a previous step of the sequence are eventually assigned to a predicted class. For example, if based on certain class-specific metrics, we determined that the order of classes in the sequential algorithm was to predict HGSC vs. non-HGSC, CCOC vs. non-CCOC, LGSOC vs. non-LGSOC, and then MUOC vs. ENOC. Figure 2.7 illustrates how the final predictions are assigned:




Figure 2.7: Aggregating Predictions for Sequential Algorithm

## 2.5 Performance Evaluation

### 2.5.1 Class Metrics

We use accuracy, sensitivity, specificity, F1-score, kappa, and balanced accuracy, as class metrics to measure both training and test performance between different workflows. Multiclass extensions of these metrics can be calculated except for F1-score, where we use macro-averaging to obtain an overall metric. Class-specific metrics are calculated by recoding classes into one-vs-all categories for each class.

#### 2.5.1.1 Accuracy

The accuracy is defined as the proportion of correct predictions out of all cases:

$$\text{accuracy} = \frac{TP}{TP + FP + FN + TN} \quad (2.6)$$

#### 2.5.1.2 Sensitivity

Sensitivity is the proportional of correctly predicted positive cases, out of all cases that were truly positive

$$\text{sensitivity} = \frac{TP}{TP + FN} \quad (2.7)$$

### 2.5.1.3 Specificity

Specificity is the proportional of correctly predicted negative cases, out of all cases that were truly negative.

$$\text{specificity} = \frac{TN}{TN + FP} \quad (2.8)$$

### 2.5.1.4 F1-Score

The F-measure can be thought of as a harmonic mean between precision and recall:

$$F_{meas} = \frac{(1 + \beta^2) \times \text{precision} \times \text{recall}}{(\beta^2 \times \text{precision}) + \text{recall}} \quad (2.9)$$

The  $\beta$  value can be adjusted to place more weight upon precision or recall. The most common value is  $\beta$  is 1, which is also commonly known as the F1-score. A multiclass extension doesn't exist for the F1-score, so we use macro-averaging to calculate this metric when there are more than two classes. For example, with  $k$  classes, the macro-averaged F1-score is equal to:

$$F_{1_{macro}} = \frac{1}{k} \sum_{i=1}^k F_{1_i} \quad (2.10)$$

where each  $F_{1_i}$  is the F1-score computed from recoding classes into  $k = i$  vs.  $k \neq i$ .

In situations where there is not at least one predicted case for each of the classes (e.g. for a poor classifier),  $F_{1_i}$  is undefined because the per-class precision of class  $i$  is undefined. Those  $F_{1_i}$  terms are removed from the  $F_{1_{macro}}$  equation and the resulting value may be inflated. Interpreting the F1-score in such a case would be misleading.

### 2.5.1.5 Balanced Accuracy

Balanced accuracy is the arithmetic mean of sensitivity and specificity.

$$\text{Balanced Accuracy} = \frac{\text{Sensitivity} + \text{Specificity}}{2} \quad (2.11)$$

### 2.5.1.6 Kappa

Kappa is the defined as:

$$\text{kappa} = \frac{p_0 - p_e}{1 - p_e} \quad (2.12)$$

where  $p_0$  is the observed agreement among raters and  $p_e$  is the hypothetical probability of agreement due to random chance.

### 2.5.2 AUC

The area under the receiver operating curve (AUC) is calculated by adding up the area under the curve formed by plotting sensitivity vs. 1 - specificity. The Hand-till method is used as a multiclass extension for the AUC.

We did not use AUC to measure class-specific training set performance because combining predicted probabilities in a one-vs-all fashion might be potentially misleading. The sum of probabilities that add up to the “other” class is not equivalent to the predicted probability of the “other” class when using a multiclass classifier.

Instead, we only reported ROC curves and their associated AUCs for test set performance among the highest ranked algorithms.

## 2.6 Rank Aggregation

To select the best algorithm, we implemented a two-stage rank aggregation procedure using the Genetic Algorithm (Pihur et al. 2009). First, we ranked all workflows based on per-class F1-scores, balanced accuracy, and kappa to see which workflows performed well in predicting all five histotypes. Then, we took the ranks from these three performance metrics and performed a second run of rank aggregation. The top 5 workflows were determined from the final rank aggregation result.

## 2.7 Gene Optimization

We want to discover an optimal set of genes for the classifiers while including specific genes from other studies such as PrOTYPE and SPOT. A total of 72 genes are used in the classifier training set.

The classifier set includes 16 genes that overlap with the PrOTYPE classifier: COL11A1, CD74, CD2, TIMP3, LUM, CYTIP, COL3A1, THBS2, TCF7L1, HMGA2, FN1, POSTN, COL1A2, COL5A2, PDZK1IP1, FBN1.

The classifier set also includes 13 genes that overlap with the SPOT signature: HIF1A, CXCL10, DUSP4, SOX17, MITF, CDKN3, BRCA2, CEACAM5, ANXA4, SERPINE1, TCF7L1, CRABP2, DNAJC9.

We obtain 28 genes from the union of PrOTYPE and SPOT genes that we want to include in the final classifier, regardless of model performance. We then incrementally add genes one at a time from the remaining 44 candidate genes based on a variable importance rank to the set of 28 core genes and recalculate performance metrics. The optimal number of genes is determined by the number of genes needed to achieve the highest average F1-Score across classes, including the overall metric in the average. We use an F1-score that is averaged across cross-validation folds (5) and class groups (6: Overall, HGSOC, CCOC, ENOC, MUOC, LGSOC) to compare performance between different number of genes selected.

Here is the breakdown of genes used and whether they belong to the PrOTYPE and/or SPOT sets:

Table 2.1: Gene Distribution

| Genes     | PrOTYPE | SPOT |
|-----------|---------|------|
| TCF7L1    | v       | v    |
| COL11A1   | v       |      |
| CD74      | v       |      |
| CD2       | v       |      |
| TIMP3     | v       |      |
| LUM       | v       |      |
| CYTIP     | v       |      |
| COL3A1    | v       |      |
| THBS2     | v       |      |
| HMGA2     | v       |      |
| FN1       | v       |      |
| POSTN     | v       |      |
| COL1A2    | v       |      |
| COL5A2    | v       |      |
| PDZK1IP1  | v       |      |
| FBN1      | v       |      |
| HIF1A     |         | v    |
| CXCL10    |         | v    |
| DUSP4     |         | v    |
| SOX17     |         | v    |
| MITF      |         | v    |
| CDKN3     |         | v    |
| BRCA2     |         | v    |
| CEACAM5   |         | v    |
| ANXA4     |         | v    |
| SERPINE1  |         | v    |
| CRABP2    |         | v    |
| DNAJC9    |         | v    |
| C10orf116 |         |      |
| GAD1      |         |      |
| TPX2      |         |      |
| KGFLP2    |         |      |
| EGFL6     |         |      |
| KLK7      |         |      |
| PBX1      |         |      |
| LIN28B    |         |      |
| TFF3      |         |      |
| MUC5B     |         |      |
| FUT3      |         |      |
| STC1      |         |      |
| BCL2      |         |      |
| PAX8      |         |      |
| GCNT3     |         |      |
| GPR64     |         |      |

ADCYAP1R1  
IGKC  
BRCA1  
IGJ  
TFF1  
MET  
CYP2C18  
CYP4B1  
SLC3A1  
EPAS1  
HNF1B  
IL6  
ATP5G3  
DKK4  
SENP8  
CAPN2  
C1orf173  
CPNE8  
IGFBP1  
WT1  
TP53  
SEMA6A  
SERPINA5  
ZBED1  
TSPAN8  
SCGB1D2  
LGALS4  
MAP1LC3A

---

### 2.7.1 Variable Importance

Variable importance is calculated using either a model-based approach if it is available, or a permutation-based VI score otherwise (e.g., for SVM). The variable importance scores are averaged across the outer training folds, and then ranked from highest to lowest.

For the sequential and two-step classifiers, we calculate an overall VI rank by taking the cumulative union of genes at each variable importance rank across all sequences, until all genes have been included.

The variable importance measures are:

- Random Forest: impurity measure (Gini index)
- XGBoost: gain (fractional contribution of each feature to the model based on the total gain of the corresponding features's splits)
- SVM: permutation based p-values

- Multinomial regression: absolute value of estimated coefficients at cross-validated lambda value

## 3 Distributions

### 3.1 Histotype Distribution

Table 3.1: Histotype Distribution in Training Set by Processing Stage

| Variable                  | Levels  | CS3        | CS1       | CS2       | Total       |
|---------------------------|---------|------------|-----------|-----------|-------------|
| <b>Selected Cohorts</b>   |         |            |           |           |             |
| Histotype                 | HGSOC   | 1808 (78%) | 128 (47%) | 655 (78%) | 2591 (75%)  |
|                           | CCOC    | 164 (7%)   | 48 (18%)  | 62 (7%)   | 274 (8%)    |
|                           | ENOC    | 250 (11%)  | 60 (22%)  | 49 (6%)   | 359 (10%)   |
|                           | MUOC    | 68 (3%)    | 17 (6%)   | 58 (7%)   | 143 (4%)    |
|                           | LGSOC   | 36 (2%)    | 19 (7%)   | 20 (2%)   | 75 (2%)     |
|                           | Missing | 151        | 22        | 59        | 232         |
| Total                     | N (%)   | 2477 (67%) | 294 (8%)  | 903 (25%) | 3674 (100%) |
| <b>QC</b>                 |         |            |           |           |             |
| Histotype                 | HGSOC   | 1676 (78%) | 122 (46%) | 641 (78%) | 2439 (75%)  |
|                           | CCOC    | 158 (7%)   | 48 (18%)  | 62 (8%)   | 268 (8%)    |
|                           | ENOC    | 213 (10%)  | 60 (23%)  | 47 (6%)   | 320 (10%)   |
|                           | MUOC    | 65 (3%)    | 16 (6%)   | 56 (7%)   | 137 (4%)    |
|                           | LGSOC   | 36 (2%)    | 18 (7%)   | 20 (2%)   | 74 (2%)     |
|                           | Missing | 125        | 22        | 56        | 203         |
| Total                     | N (%)   | 2273 (66%) | 286 (8%)  | 882 (26%) | 3441 (100%) |
| <b>Main Histotypes</b>    |         |            |           |           |             |
| Histotype                 | HGSOC   | 1676 (78%) | 122 (46%) | 641 (78%) | 2439 (75%)  |
|                           | CCOC    | 158 (7%)   | 48 (18%)  | 62 (8%)   | 268 (8%)    |
|                           | ENOC    | 213 (10%)  | 60 (23%)  | 47 (6%)   | 320 (10%)   |
|                           | MUOC    | 65 (3%)    | 16 (6%)   | 56 (7%)   | 137 (4%)    |
|                           | LGSOC   | 36 (2%)    | 18 (7%)   | 20 (2%)   | 74 (2%)     |
| Total                     | N (%)   | 2148 (66%) | 264 (8%)  | 826 (26%) | 3238 (100%) |
| <b>Removed Duplicates</b> |         |            |           |           |             |
|                           | HGSOC   | 1578 (78%) | 118 (48%) | 623 (78%) | 2319 (76%)  |

|                                  |       |            |           |           |             |
|----------------------------------|-------|------------|-----------|-----------|-------------|
| Histotype                        | CCOC  | 146 (7%)   | 45 (18%)  | 56 (7%)   | 247 (8%)    |
|                                  | ENOC  | 200 (10%)  | 56 (23%)  | 43 (5%)   | 299 (10%)   |
|                                  | MUOC  | 55 (3%)    | 13 (5%)   | 54 (7%)   | 122 (4%)    |
|                                  | LGSOC | 32 (2%)    | 14 (6%)   | 19 (2%)   | 65 (2%)     |
| Total                            | N (%) | 2011 (66%) | 246 (8%)  | 795 (26%) | 3052 (100%) |
| <b>Normalized and Recombined</b> |       |            |           |           |             |
| Histotype                        | HGSOC | 454 (97%)  | 117 (49%) | 622 (79%) | 1193 (79%)  |
|                                  | CCOC  | 4 (1%)     | 44 (18%)  | 55 (7%)   | 103 (7%)    |
|                                  | ENOC  | 4 (1%)     | 55 (23%)  | 42 (5%)   | 101 (7%)    |
|                                  | MUOC  | 4 (1%)     | 12 (5%)   | 53 (7%)   | 69 (5%)     |
|                                  | LGSOC | 4 (1%)     | 13 (5%)   | 18 (2%)   | 35 (2%)     |
| Total                            | N (%) | 470 (31%)  | 241 (16%) | 790 (53%) | 1501 (100%) |
| <b>Removed Replicates</b>        |       |            |           |           |             |
| Histotype                        | HGSOC | 454 (97%)  | 9 (12%)   | 552 (78%) | 1015 (81%)  |
|                                  | CCOC  | 4 (1%)     | 24 (31%)  | 53 (7%)   | 81 (6%)     |
|                                  | ENOC  | 4 (1%)     | 38 (49%)  | 40 (6%)   | 82 (7%)     |
|                                  | MUOC  | 4 (1%)     | 3 (4%)    | 50 (7%)   | 57 (5%)     |
|                                  | LGSOC | 4 (1%)     | 3 (4%)    | 15 (2%)   | 22 (2%)     |
| Total                            | N (%) | 470 (37%)  | 77 (6%)   | 710 (56%) | 1257 (100%) |

Table 3.2: Histotype Distribution in Training, Confirmation, and Validation Sets

| Variable  | Levels  | Training   | Confirmation | Validation |
|-----------|---------|------------|--------------|------------|
| Histotype | CCOC    | 81 (50%)   | 72 (40%)     | 69 (44%)   |
|           | ENOC    | 82 (50%)   | 107 (60%)    | 88 (56%)   |
|           | Missing | 1094       | 463          | 737        |
| Total     | N (%)   | 1257 (45%) | 642 (23%)    | 894 (32%)  |

### 3.2 Cohort Distribution

Table 3.3: Pre-QC Cohort Distribution by CodeSet

| CodeSet   | CS1<br>N = 294 | CS2<br>N = 903 | CS3<br>N = 2,477 |
|-----------|----------------|----------------|------------------|
| Cohort    |                |                |                  |
| OOU       | 108 (37%)      | 43 (4.8%)      | 19 (0.8%)        |
| OOUE      | 32 (11%)       | 30 (3.3%)      | 11 (0.4%)        |
| VOA       | 145 (49%)      | 122 (14%)      | 538 (22%)        |
| OVAR3     | 0 (0%)         | 150 (17%)      | 0 (0%)           |
| OVAR11    | 0 (0%)         | 416 (46%)      | 0 (0%)           |
| MAYO      | 6 (2.0%)       | 63 (7.0%)      | 0 (0%)           |
| DOVE4     | 0 (0%)         | 0 (0%)         | 1,160 (47%)      |
| TNCO      | 0 (0%)         | 0 (0%)         | 691 (28%)        |
| MTL       | 3 (1.0%)       | 59 (6.5%)      | 0 (0%)           |
| JAPAN     | 0 (0%)         | 8 (0.9%)       | 0 (0%)           |
| POOL-CTRL | 0 (0%)         | 12 (1.3%)      | 0 (0%)           |
| POOL-1    | 0 (0%)         | 0 (0%)         | 31 (1.3%)        |
| POOL-2    | 0 (0%)         | 0 (0%)         | 14 (0.6%)        |
| POOL-3    | 0 (0%)         | 0 (0%)         | 13 (0.5%)        |

<sup>1</sup> n (%)

### 3.3 Tissue Source

Table 3.4: Tissue Source by Cohort

| Data         | Cohort | FFPE | Frozen |
|--------------|--------|------|--------|
| Training     | OVAR11 | 365  | 0      |
|              | JAPAN  | 0    | 8      |
|              | MAYO   | 56   | 0      |
|              | MTL    | 19   | 41     |
|              | OOU    | 98   | 0      |
|              | OOUE   | 30   | 0      |
|              | OVAR3  | 135  | 0      |
|              | VOA    | 494  | 11     |
| Confirmation | TNCO   | 642  | 0      |
| Validation   | DOVE4  | 894  | 0      |

## 3.4 Quality Control

### 3.4.1 Failed Samples

We use an aggregated **QCFlag** that considers a sample to have failed QC if any of the following QC conditions are flagged:

- Linearity
- Imaging
- Smallest Positive Control
- Normality

Table 3.5: Quality Control Summary

| Quality Control Flag      | Training<br>N = 1,753 | Confirmation<br>N = 691 | Validation<br>N = 1,160 | Total<br>N = 3,604 |
|---------------------------|-----------------------|-------------------------|-------------------------|--------------------|
| Linearity                 | 4 (0.2%)              | 0 (0%)                  | 0 (0%)                  | 4 (0.1%)           |
| Imaging                   | 3 (0.2%)              | 0 (0%)                  | 4 (0.3%)                | 7 (0.2%)           |
| Smallest Positive Control | 2 (0.1%)              | 0 (0%)                  | 0 (0%)                  | 2 (<0.1%)          |
| Normality                 | 26 (1.5%)             | 1 (0.1%)                | 197 (17%)               | 224 (6.2%)         |
| Overall QC                | 31 (1.8%)             | 1 (0.1%)                | 201 (17%)               | 233 (6.5%)         |

<sup>1</sup> n (%)

### 3.4.2 %GD vs. SNR

% Genes Detected vs. Signal-to-Noise Ratio




Figure 3.1: % Genes Detected vs. Signal to Noise Ratio

### % Genes Detected vs. Signal-to-Noise Ratio (Zoomed)




Figure 3.2: % Genes Detected vs. Signal to Noise Ratio (Zoomed)

### 3.5 Pairwise Gene Expression




Figure 3.3: Random1-Normalized CS1 vs. CS3 Gene Expression




Figure 3.4: Random1-Normalized CS2 vs. CS3 Gene Expression




Figure 3.5: HKgenes-Normalized CS1 vs. CS3 Gene Expression




Figure 3.6: HKgenes-Normalized CS2 vs. CS3 Gene Expression

Table 3.6: Wilcoxon signed rank test of gene correlations between normalization methods

| Correlation | Housekeeping Genes N = 72 <sup>1</sup> | Random1 N = 72 <sup>1</sup> | p-value <sup>2</sup> |
|-------------|----------------------------------------|-----------------------------|----------------------|
| CS1 vs. CS3 | 0.84 (0.74, 0.90)                      | 0.84 (0.67, 0.89)           | 0.079                |
| CS2 vs. CS3 | 0.80 (0.69, 0.88)                      | 0.84 (0.72, 0.88)           | 0.003                |

<sup>1</sup>Median (Q1, Q3)

<sup>2</sup>Wilcoxon signed rank test with continuity correction

## 4 Results

We summarize cross-validated training performance of class metrics in the training set. The accuracy, F1-score, and kappa, are the metrics of interest. Workflows are ordered by their mean estimates across the outer folds of the nested CV for each metric.

## 4.1 Training Set

### 4.1.1 Accuracy

Table 4.1: Training Set Mean Accuracy

| Subsampling | Algorithms | Overall | Histotypes |       |       |       |       |
|-------------|------------|---------|------------|-------|-------|-------|-------|
|             |            |         | HGSOC      | CCOC  | ENOC  | MUOC  | LGSOC |
| none        | rf         | 0.922   | 0.948      | 0.983 | 0.953 | 0.975 | 0.984 |
|             | svm        | 0.92    | 0.947      | 0.982 | 0.955 | 0.975 | 0.982 |
|             | xgb        | 0.807   | 0.809      | 0.936 | 0.933 | 0.955 | 0.982 |
|             | mr         | 0.807   | 0.807      | 0.936 | 0.935 | 0.955 | 0.982 |
| down        | rf         | 0.804   | 0.853      | 0.975 | 0.903 | 0.959 | 0.92  |
|             | svm        | 0.793   | 0.833      | 0.973 | 0.884 | 0.969 | 0.928 |
|             | xgb        | 0.783   | 0.833      | 0.968 | 0.897 | 0.961 | 0.907 |
|             | mr         | 0.829   | 0.87       | 0.977 | 0.912 | 0.961 | 0.938 |
| up          | rf         | 0.928   | 0.955      | 0.983 | 0.959 | 0.979 | 0.982 |
|             | svm        | 0.92    | 0.951      | 0.974 | 0.951 | 0.979 | 0.985 |
|             | xgb        | 0.928   | 0.959      | 0.982 | 0.96  | 0.974 | 0.982 |
|             | mr         | 0.891   | 0.924      | 0.976 | 0.949 | 0.966 | 0.967 |
| smote       | rf         | 0.921   | 0.955      | 0.98  | 0.953 | 0.97  | 0.984 |
|             | svm        | 0.92    | 0.95       | 0.979 | 0.954 | 0.978 | 0.981 |
|             | xgb        | 0.923   | 0.955      | 0.98  | 0.955 | 0.971 | 0.984 |
|             | mr         | 0.897   | 0.931      | 0.979 | 0.951 | 0.967 | 0.967 |
| hybrid      | rf         | 0.922   | 0.953      | 0.98  | 0.952 | 0.975 | 0.983 |
|             | svm        | 0.916   | 0.944      | 0.982 | 0.951 | 0.975 | 0.979 |
|             | xgb        | 0.924   | 0.954      | 0.984 | 0.958 | 0.971 | 0.982 |
|             | mr         | 0.892   | 0.925      | 0.978 | 0.947 | 0.967 | 0.966 |




Figure 4.1: Training Set Mean Accuracy

### 4.1.2 Sensitivity

Table 4.2: Training Set Mean Sensitivity

| Subsampling | Algorithms | Overall | Histotypes |       |       |       |       |
|-------------|------------|---------|------------|-------|-------|-------|-------|
|             |            |         | HGSOC      | CCOC  | ENOC  | MUOC  | LGSOC |
| none        | rf         | 0.636   | 0.994      | 0.794 | 0.498 | 0.728 | 0.167 |
|             | svm        | 0.615   | 0.989      | 0.762 | 0.576 | 0.751 | 0     |
|             | xgb        | 0.2     | 1          | 0     | 0     | 0     | 0     |
|             | mr         | 0.2     | 1          | 0     | 0     | 0     | 0     |
| down        | rf         | 0.742   | 0.83       | 0.827 | 0.565 | 0.657 | 0.833 |
|             | svm        | 0.779   | 0.805      | 0.759 | 0.611 | 0.836 | 0.883 |
|             | xgb        | 0.714   | 0.802      | 0.843 | 0.589 | 0.67  | 0.667 |
|             | mr         | 0.776   | 0.848      | 0.831 | 0.62  | 0.77  | 0.808 |
| up          | rf         | 0.673   | 0.988      | 0.793 | 0.635 | 0.766 | 0.183 |
|             | svm        | 0.714   | 0.976      | 0.735 | 0.625 | 0.71  | 0.525 |
|             | xgb        | 0.718   | 0.981      | 0.805 | 0.624 | 0.781 | 0.4   |
|             | mr         | 0.788   | 0.921      | 0.849 | 0.666 | 0.786 | 0.717 |
| smote       | rf         | 0.658   | 0.984      | 0.748 | 0.618 | 0.75  | 0.192 |
|             | svm        | 0.759   | 0.966      | 0.757 | 0.681 | 0.748 | 0.642 |
|             | xgb        | 0.777   | 0.965      | 0.831 | 0.615 | 0.804 | 0.667 |
|             | mr         | 0.788   | 0.93       | 0.835 | 0.654 | 0.802 | 0.717 |
| hybrid      | rf         | 0.72    | 0.972      | 0.805 | 0.638 | 0.804 | 0.383 |
|             | svm        | 0.764   | 0.957      | 0.793 | 0.696 | 0.733 | 0.642 |
|             | xgb        | 0.77    | 0.966      | 0.851 | 0.624 | 0.818 | 0.592 |
|             | mr         | 0.783   | 0.923      | 0.835 | 0.676 | 0.767 | 0.717 |




Figure 4.2: Training Set Mean Sensitivity

### 4.1.3 Specificity

Table 4.3: Training Set Mean Specificity

| Subsampling | Algorithms | Overall | Histotypes |       |       |       |       |
|-------------|------------|---------|------------|-------|-------|-------|-------|
|             |            |         | HGSOC      | CCOC  | ENOC  | MUOC  | LGSOC |
| none        | rf         | 0.946   | 0.764      | 0.996 | 0.985 | 0.988 | 0.999 |
|             | svm        | 0.947   | 0.775      | 0.997 | 0.98  | 0.986 | 1     |
|             | xgb        | 0.802   | 0.009      | 1     | 0.998 | 1     | 1     |
|             | mr         | 0.8     | 0          | 1     | 1     | 1     | 1     |
| down        | rf         | 0.952   | 0.955      | 0.984 | 0.928 | 0.973 | 0.921 |
|             | svm        | 0.949   | 0.952      | 0.987 | 0.902 | 0.976 | 0.928 |
|             | xgb        | 0.949   | 0.965      | 0.976 | 0.918 | 0.975 | 0.911 |
|             | mr         | 0.958   | 0.963      | 0.986 | 0.931 | 0.971 | 0.94  |
| up          | rf         | 0.957   | 0.822      | 0.996 | 0.98  | 0.989 | 0.997 |
|             | svm        | 0.958   | 0.844      | 0.99  | 0.973 | 0.992 | 0.993 |
|             | xgb        | 0.964   | 0.867      | 0.994 | 0.982 | 0.983 | 0.993 |
|             | mr         | 0.967   | 0.938      | 0.985 | 0.968 | 0.975 | 0.971 |
| smote       | rf         | 0.958   | 0.839      | 0.996 | 0.975 | 0.981 | 0.998 |
|             | svm        | 0.964   | 0.88       | 0.993 | 0.971 | 0.989 | 0.987 |
|             | xgb        | 0.969   | 0.91       | 0.99  | 0.978 | 0.98  | 0.99  |
|             | mr         | 0.968   | 0.934      | 0.988 | 0.971 | 0.976 | 0.971 |
| hybrid      | rf         | 0.964   | 0.874      | 0.992 | 0.974 | 0.984 | 0.994 |
|             | svm        | 0.965   | 0.89       | 0.994 | 0.968 | 0.987 | 0.985 |
|             | xgb        | 0.969   | 0.904      | 0.992 | 0.98  | 0.979 | 0.989 |
|             | mr         | 0.967   | 0.934      | 0.987 | 0.965 | 0.977 | 0.97  |




Figure 4.3: Training Set Mean Specificity

#### 4.1.4 F1-Score

Table 4.4: Training Set Mean F1-Score

| Subsampling | Algorithms | Overall | Histotypes |       |       |       |       |
|-------------|------------|---------|------------|-------|-------|-------|-------|
|             |            |         | HGSOC      | CCOC  | ENOC  | MUOC  | LGSOC |
| none        | rf         | 0.733   | 0.968      | 0.848 | 0.568 | 0.723 | 0.4   |
|             | svm        | 0.782   | 0.968      | 0.834 | 0.607 | 0.719 | NaN   |
|             | xgb        | 0.712   | 0.894      | NaN   | 0     | NaN   | NaN   |
|             | mr         | 0.893   | 0.893      | NaN   | NaN   | NaN   | NaN   |
| down        | rf         | 0.594   | 0.901      | 0.799 | 0.419 | 0.589 | 0.262 |
|             | svm        | 0.617   | 0.886      | 0.778 | 0.412 | 0.713 | 0.295 |
|             | xgb        | 0.576   | 0.886      | 0.771 | 0.42  | 0.608 | 0.194 |
|             | mr         | 0.628   | 0.913      | 0.817 | 0.47  | 0.638 | 0.302 |
| up          | rf         | 0.755   | 0.972      | 0.849 | 0.652 | 0.757 | 0.362 |
|             | svm        | 0.725   | 0.97       | 0.778 | 0.605 | 0.75  | 0.521 |
|             | xgb        | 0.723   | 0.974      | 0.844 | 0.651 | 0.728 | 0.416 |
|             | mr         | 0.695   | 0.951      | 0.816 | 0.619 | 0.674 | 0.413 |
| smote       | rf         | 0.68    | 0.972      | 0.82  | 0.616 | 0.688 | 0.304 |
|             | svm        | 0.739   | 0.969      | 0.811 | 0.638 | 0.752 | 0.524 |
|             | xgb        | 0.744   | 0.972      | 0.833 | 0.628 | 0.71  | 0.579 |
|             | mr         | 0.704   | 0.956      | 0.827 | 0.629 | 0.691 | 0.418 |
| hybrid      | rf         | 0.724   | 0.971      | 0.829 | 0.631 | 0.741 | 0.45  |
|             | svm        | 0.733   | 0.965      | 0.838 | 0.635 | 0.72  | 0.507 |
|             | xgb        | 0.741   | 0.971      | 0.86  | 0.643 | 0.716 | 0.515 |
|             | mr         | 0.696   | 0.952      | 0.821 | 0.613 | 0.682 | 0.412 |




Figure 4.4: Training Set Mean F1-Score

### 4.1.5 Balanced Accuracy

Table 4.5: Training Set Mean Balanced Accuracy

| Subsampling | Algorithms | Overall | Histotypes |       |       |       |       |
|-------------|------------|---------|------------|-------|-------|-------|-------|
|             |            |         | HGSOC      | CCOC  | ENOC  | MUOC  | LGSOC |
| none        | rf         | 0.791   | 0.879      | 0.895 | 0.741 | 0.858 | 0.583 |
|             | svm        | 0.781   | 0.882      | 0.879 | 0.778 | 0.868 | 0.5   |
|             | xgb        | 0.501   | 0.505      | 0.5   | 0.499 | 0.5   | 0.5   |
|             | mr         | 0.5     | 0.5        | 0.5   | 0.5   | 0.5   | 0.5   |
| down        | rf         | 0.847   | 0.892      | 0.905 | 0.746 | 0.815 | 0.877 |
|             | svm        | 0.864   | 0.879      | 0.873 | 0.757 | 0.906 | 0.906 |
|             | xgb        | 0.832   | 0.883      | 0.91  | 0.753 | 0.822 | 0.789 |
|             | mr         | 0.867   | 0.906      | 0.909 | 0.776 | 0.87  | 0.874 |
| up          | rf         | 0.815   | 0.905      | 0.894 | 0.808 | 0.878 | 0.59  |
|             | svm        | 0.836   | 0.91       | 0.862 | 0.799 | 0.851 | 0.759 |
|             | xgb        | 0.841   | 0.924      | 0.9   | 0.803 | 0.882 | 0.696 |
|             | mr         | 0.878   | 0.93       | 0.917 | 0.817 | 0.881 | 0.844 |
| smote       | rf         | 0.808   | 0.911      | 0.872 | 0.797 | 0.865 | 0.595 |
|             | svm        | 0.862   | 0.923      | 0.875 | 0.826 | 0.869 | 0.814 |
|             | xgb        | 0.873   | 0.938      | 0.911 | 0.797 | 0.892 | 0.828 |
|             | mr         | 0.878   | 0.932      | 0.912 | 0.813 | 0.889 | 0.844 |
| hybrid      | rf         | 0.842   | 0.923      | 0.898 | 0.806 | 0.894 | 0.689 |
|             | svm        | 0.864   | 0.923      | 0.893 | 0.832 | 0.86  | 0.814 |
|             | xgb        | 0.87    | 0.935      | 0.922 | 0.802 | 0.899 | 0.791 |
|             | mr         | 0.875   | 0.928      | 0.911 | 0.821 | 0.872 | 0.843 |




Figure 4.5: Training Set Mean Balanced Accuracy

### 4.1.6 Kappa

Table 4.6: Training Set Mean Kappa

| Subsampling | Algorithms | Overall | Histotypes |       |        |       |       |
|-------------|------------|---------|------------|-------|--------|-------|-------|
|             |            |         | HGSOC      | CCOC  | ENOC   | MUOC  | LGSOC |
| none        | rf         | 0.744   | 0.822      | 0.84  | 0.545  | 0.711 | 0.237 |
|             | svm        | 0.741   | 0.818      | 0.825 | 0.583  | 0.706 | 0     |
|             | xgb        | 0.007   | 0.016      | 0     | -0.003 | 0     | 0     |
|             | mr         | 0       | 0          | 0     | 0      | 0     | 0     |
| down        | rf         | 0.55    | 0.621      | 0.785 | 0.37   | 0.568 | 0.24  |
|             | svm        | 0.539   | 0.584      | 0.764 | 0.359  | 0.697 | 0.275 |
|             | xgb        | 0.519   | 0.584      | 0.754 | 0.37   | 0.588 | 0.17  |
|             | mr         | 0.595   | 0.657      | 0.805 | 0.426  | 0.618 | 0.282 |
| up          | rf         | 0.776   | 0.85       | 0.84  | 0.63   | 0.746 | 0.213 |
|             | svm        | 0.749   | 0.835      | 0.764 | 0.579  | 0.739 | 0.513 |
|             | xgb        | 0.781   | 0.865      | 0.835 | 0.63   | 0.714 | 0.408 |
|             | mr         | 0.708   | 0.778      | 0.803 | 0.592  | 0.657 | 0.399 |
| smote       | rf         | 0.756   | 0.853      | 0.81  | 0.591  | 0.673 | 0.299 |
|             | svm        | 0.761   | 0.839      | 0.8   | 0.614  | 0.741 | 0.516 |
|             | xgb        | 0.773   | 0.856      | 0.822 | 0.604  | 0.695 | 0.571 |
|             | mr         | 0.72    | 0.793      | 0.816 | 0.603  | 0.675 | 0.405 |
| hybrid      | rf         | 0.767   | 0.85       | 0.818 | 0.605  | 0.728 | 0.443 |
|             | svm        | 0.753   | 0.825      | 0.828 | 0.609  | 0.707 | 0.497 |
|             | xgb        | 0.778   | 0.855      | 0.852 | 0.62   | 0.702 | 0.506 |
|             | mr         | 0.707   | 0.779      | 0.81  | 0.585  | 0.665 | 0.399 |




Figure 4.6: Training Set Mean Kappa

## 4.2 Rank Aggregation

Multi-step methods:

- **sequential:** sequential algorithm sequence of subsampling methods and algorithms used are:

- HGSOC vs. non-HGSOC using upsampling and XGBoost
  - CCOC vs. non-CCOC using SMOTE subsampling and XGBoost
  - ENOC vs. non-ENOC using no subsampling and support vector machine
  - MUOC vs. LGSOC using SMOTE subsampling and random forest
- **two\_step**: two-step algorithm sequence of subsampling methods and algorithms used are:
    - HGSOC vs. non-HGSOC using upsampling and XGBoost
    - CCOC vs. ENOC vs. MUOC vs. LGSOC using SMOTE subsampling and support vector machine

We conduct rank aggregation using a two-stage nested approach:

1. First we rank aggregate the per-class metrics for F1-score, balanced accuracy and kappa.
2. Then we take the aggregated lists from the three metrics and perform a final rank aggregation.
3. The top workflows from the final rank aggregation are used for gene optimization in the confirmation set

## 4.2.1 Across Classes

### 4.2.1.1 F1-Score

Table 4.7: F1-Score Rank Aggregation Summary

| Workflow   | Rank | HGSOC | CCOC  | ENOC  | MUOC  | LGSOC |
|------------|------|-------|-------|-------|-------|-------|
| sequential | 1    | 0.973 | 0.877 | 0.86  | 0.966 | 0.91  |
| two_step   | 2    | 0.973 | 0.899 | 0.755 | 0.788 | 0.733 |
| up_rf      | 3    | 0.972 | 0.849 | 0.652 | 0.757 | 0.362 |
| smote_svm  | 4    | 0.969 | 0.811 | 0.638 | 0.752 | 0.524 |
| hybrid_xgb | 5    | 0.971 | 0.86  | 0.643 | 0.716 | 0.515 |
| up_xgb     | 6    | 0.974 | 0.844 | 0.651 | 0.728 | 0.416 |
| hybrid_svm | 7    | 0.965 | 0.838 | 0.635 | 0.72  | 0.507 |
| hybrid_rf  | 8    | 0.971 | 0.829 | 0.631 | 0.741 | 0.45  |
| smote_xgb  | 9    | 0.972 | 0.833 | 0.628 | 0.71  | 0.579 |
| smote_mr   | 10   | 0.956 | 0.827 | 0.629 | 0.691 | 0.418 |
| up_svm     | 11   | 0.97  | 0.778 | 0.605 | 0.75  | 0.521 |
| smote_rf   | 12   | 0.972 | 0.82  | 0.616 | 0.688 | 0.304 |
| none_rf    | 13   | 0.968 | 0.848 | 0.568 | 0.723 | 0.4   |
| hybrid_mr  | 14   | 0.952 | 0.821 | 0.613 | 0.682 | 0.412 |
| up_mr      | 15   | 0.951 | 0.816 | 0.619 | 0.674 | 0.413 |
| down_mr    | 16   | 0.913 | 0.817 | 0.47  | 0.638 | 0.302 |
| down_svm   | 17   | 0.886 | 0.778 | 0.412 | 0.713 | 0.295 |
| down_rf    | 18   | 0.901 | 0.799 | 0.419 | 0.589 | 0.262 |
| down_xgb   | 19   | 0.886 | 0.771 | 0.42  | 0.608 | 0.194 |

### 4.2.1.2 Balanced Accuracy

Table 4.8: Balanced Accuracy Rank Aggregation Summary

| Workflow   | Rank | HGSOC | CCOC  | ENOC  | MUOC  | LGSOC |
|------------|------|-------|-------|-------|-------|-------|
| sequential | 1    | 0.926 | 0.907 | 0.856 | 0.943 | 0.943 |
| hybrid_xgb | 2    | 0.935 | 0.922 | 0.802 | 0.899 | 0.791 |
| up_mr      | 3    | 0.93  | 0.917 | 0.817 | 0.881 | 0.844 |
| smote_mr   | 4    | 0.932 | 0.912 | 0.813 | 0.889 | 0.844 |
| smote_xgb  | 5    | 0.938 | 0.911 | 0.797 | 0.892 | 0.828 |
| two_step   | 6    | 0.926 | 0.917 | 0.814 | 0.875 | 0.872 |
| hybrid_mr  | 7    | 0.928 | 0.911 | 0.821 | 0.872 | 0.843 |
| up_xgb     | 8    | 0.924 | 0.9   | 0.803 | 0.882 | 0.696 |
| hybrid_svm | 9    | 0.923 | 0.893 | 0.832 | 0.86  | 0.814 |
| smote_svm  | 10   | 0.923 | 0.875 | 0.826 | 0.869 | 0.814 |
| down_mr    | 11   | 0.906 | 0.909 | 0.776 | 0.87  | 0.874 |
| hybrid_rf  | 12   | 0.923 | 0.898 | 0.806 | 0.894 | 0.689 |
| up_svm     | 13   | 0.91  | 0.862 | 0.799 | 0.851 | 0.759 |
| up_rf      | 14   | 0.905 | 0.894 | 0.808 | 0.878 | 0.59  |
| smote_rf   | 15   | 0.911 | 0.872 | 0.797 | 0.865 | 0.595 |
| down_rf    | 16   | 0.892 | 0.905 | 0.746 | 0.815 | 0.877 |
| down_xgb   | 17   | 0.883 | 0.91  | 0.753 | 0.822 | 0.789 |
| none_svm   | 18   | 0.882 | 0.879 | 0.778 | 0.868 | 0.5   |
| none_rf    | 19   | 0.879 | 0.895 | 0.741 | 0.858 | 0.583 |
| down_svm   | 20   | 0.879 | 0.873 | 0.757 | 0.906 | 0.906 |
| none_mr    | 21   | 0.5   | 0.5   | 0.5   | 0.5   | 0.5   |
| none_xgb   | 22   | 0.505 | 0.5   | 0.499 | 0.5   | 0.5   |

### 4.2.1.3 Kappa

Table 4.9: Kappa Rank Aggregation Summary

| Workflow   | Rank | HGSOC | CCOC  | ENOC   | MUOC  | LGSOC |
|------------|------|-------|-------|--------|-------|-------|
| sequential | 1    | 0.858 | 0.815 | 0.712  | 0.877 | 0.877 |
| two_step   | 2    | 0.858 | 0.85  | 0.635  | 0.716 | 0.718 |
| smote_svm  | 3    | 0.839 | 0.8   | 0.614  | 0.741 | 0.516 |
| up_rf      | 4    | 0.85  | 0.84  | 0.63   | 0.746 | 0.213 |
| up_xgb     | 5    | 0.865 | 0.835 | 0.63   | 0.714 | 0.408 |
| hybrid_xgb | 6    | 0.855 | 0.852 | 0.62   | 0.702 | 0.506 |
| hybrid_svm | 7    | 0.825 | 0.828 | 0.609  | 0.707 | 0.497 |
| hybrid_rf  | 8    | 0.85  | 0.818 | 0.605  | 0.728 | 0.443 |
| smote_xgb  | 9    | 0.856 | 0.822 | 0.604  | 0.695 | 0.571 |
| up_svm     | 10   | 0.835 | 0.764 | 0.579  | 0.739 | 0.513 |
| smote_mr   | 11   | 0.793 | 0.816 | 0.603  | 0.675 | 0.405 |
| smote_rf   | 12   | 0.853 | 0.81  | 0.591  | 0.673 | 0.299 |
| none_svm   | 13   | 0.818 | 0.825 | 0.583  | 0.706 | 0     |
| none_rf    | 14   | 0.822 | 0.84  | 0.545  | 0.711 | 0.237 |
| hybrid_mr  | 15   | 0.779 | 0.81  | 0.585  | 0.665 | 0.399 |
| up_mr      | 16   | 0.778 | 0.803 | 0.592  | 0.657 | 0.399 |
| down_mr    | 17   | 0.657 | 0.805 | 0.426  | 0.618 | 0.282 |
| down_rf    | 18   | 0.621 | 0.785 | 0.37   | 0.568 | 0.24  |
| down_xgb   | 19   | 0.584 | 0.754 | 0.37   | 0.588 | 0.17  |
| down_svm   | 20   | 0.584 | 0.764 | 0.359  | 0.697 | 0.275 |
| none_mr    | 21   | 0     | 0     | 0      | 0     | 0     |
| none_xgb   | 22   | 0.016 | 0     | -0.003 | 0     | 0     |

### 4.2.2 Across Metrics

Table 4.10: Rank Aggregation Comparison of Metrics Used in Training Set

| Rank | F1         | Balanced Accuracy | Kappa      |
|------|------------|-------------------|------------|
| 1    | sequential | sequential        | sequential |
| 2    | two_step   | hybrid_xgb        | two_step   |
| 3    | up_rf      | up_mr             | smote_svm  |
| 4    | smote_svm  | smote_mr          | up_rf      |
| 5    | hybrid_xgb | smote_xgb         | up_xgb     |
| 6    | up_xgb     | two_step          | hybrid_xgb |
| 7    | hybrid_svm | hybrid_mr         | hybrid_svm |
| 8    | hybrid_rf  | up_xgb            | hybrid_rf  |
| 9    | smote_xgb  | hybrid_svm        | smote_xgb  |
| 10   | smote_mr   | smote_svm         | up_svm     |
| 11   | up_svm     | down_mr           | smote_mr   |
| 12   | smote_rf   | hybrid_rf         | smote_rf   |
| 13   | none_rf    | up_svm            | none_svm   |
| 14   | hybrid_mr  | up_rf             | none_rf    |
| 15   | up_mr      | smote_rf          | hybrid_mr  |
| 16   | down_mr    | down_rf           | up_mr      |
| 17   | down_svm   | down_xgb          | down_mr    |
| 18   | down_rf    | none_svm          | down_rf    |
| 19   | down_xgb   | none_rf           | down_xgb   |
| 20   | NA         | down_svm          | down_svm   |
| 21   | NA         | none_mr           | none_mr    |
| 22   | NA         | none_xgb          | none_xgb   |

Table 4.11: Top 5 Workflows from Final Rank Aggregation

| Rank | Workflow   |
|------|------------|
| 1    | sequential |
| 2    | two_step   |
| 3    | up_rf      |
| 4    | smote_svm  |
| 5    | hybrid_xgb |

### 4.2.3 Top Workflows

We look at the per-class evaluation metrics of the top 5 workflows.

Table 4.12: Top Workflow Per-Class Evaluation Metrics

| Metric            | Workflow   | Histotypes        |                   |                   |                   |                   |
|-------------------|------------|-------------------|-------------------|-------------------|-------------------|-------------------|
|                   |            | HGSOC             | CCOC              | ENOC              | MUOC              | LGSOC             |
| Accuracy          | Sequential | 0.96 (0.95, 0.96) | 0.92 (0.88, 0.96) | 0.86 (0.77, 0.91) | 0.95 (0.88, 1)    | 0.95 (0.88, 1)    |
|                   | 2-STEP     | 0.96 (0.95, 0.96) | 0.93 (0.9, 0.96)  | 0.84 (0.73, 0.9)  | 0.89 (0.81, 0.96) | 0.97 (0.94, 1)    |
|                   | Up-RF      | 0.95 (0.92, 0.98) | 0.98 (0.97, 0.99) | 0.96 (0.95, 0.97) | 0.98 (0.96, 0.99) | 0.98 (0.97, 0.99) |
|                   | SMOTE-SVM  | 0.95 (0.94, 0.97) | 0.98 (0.97, 0.99) | 0.95 (0.94, 0.96) | 0.98 (0.96, 0.98) | 0.98 (0.97, 0.98) |
|                   | Hybrid-XGB | 0.95 (0.94, 0.97) | 0.98 (0.97, 0.99) | 0.96 (0.94, 0.97) | 0.97 (0.96, 0.98) | 0.98 (0.97, 0.99) |
| Sensitivity       | Sequential | 0.98 (0.97, 0.99) | 0.88 (0.81, 1)    | 0.86 (0.81, 0.94) | 0.97 (0.91, 1)    | 0.92 (0.6, 1)     |
|                   | 2-STEP     | 0.98 (0.97, 0.99) | 0.87 (0.82, 0.93) | 0.73 (0.56, 0.82) | 0.84 (0.75, 0.92) | 0.77 (0, 1)       |
|                   | Up-RF      | 0.99 (0.98, 1)    | 0.79 (0.57, 0.88) | 0.64 (0.44, 0.81) | 0.77 (0.46, 0.91) | 0.18 (0, 0.33)    |
|                   | SMOTE-SVM  | 0.97 (0.96, 0.98) | 0.76 (0.57, 0.85) | 0.68 (0.5, 0.86)  | 0.75 (0.54, 0.91) | 0.64 (0.5, 0.75)  |
|                   | Hybrid-XGB | 0.97 (0.96, 0.98) | 0.85 (0.57, 0.95) | 0.62 (0.44, 0.77) | 0.82 (0.54, 0.91) | 0.59 (0.38, 0.75) |
| Specificity       | Sequential | 0.88 (0.85, 0.9)  | 0.94 (0.85, 1)    | 0.85 (0.73, 0.88) | 0.92 (0.6, 1)     | 0.97 (0.91, 1)    |
|                   | 2-STEP     | 0.88 (0.85, 0.9)  | 0.97 (0.94, 1)    | 0.89 (0.83, 0.94) | 0.91 (0.83, 0.97) | 0.98 (0.95, 1)    |
|                   | Up-RF      | 0.82 (0.74, 0.89) | 1 (0.99, 1)       | 0.98 (0.97, 0.99) | 0.99 (0.98, 1)    | 1 (0.99, 1)       |
|                   | SMOTE-SVM  | 0.88 (0.8, 0.92)  | 0.99 (0.99, 1)    | 0.97 (0.96, 0.98) | 0.99 (0.98, 1)    | 0.99 (0.98, 1)    |
|                   | Hybrid-XGB | 0.9 (0.85, 0.95)  | 0.99 (0.99, 1)    | 0.98 (0.97, 0.99) | 0.98 (0.97, 0.99) | 0.99 (0.98, 0.99) |
| F1-Score          | Sequential | 0.97 (0.97, 0.98) | 0.88 (0.81, 0.94) | 0.86 (0.79, 0.91) | 0.97 (0.92, 1)    | 0.91 (0.75, 1)    |
|                   | 2-STEP     | 0.97 (0.97, 0.98) | 0.9 (0.85, 0.93)  | 0.75 (0.61, 0.85) | 0.79 (0.67, 0.92) | 0.73 (0, 1)       |
|                   | Up-RF      | 0.97 (0.95, 0.99) | 0.85 (0.7, 0.94)  | 0.65 (0.47, 0.79) | 0.76 (0.57, 0.87) | 0.36 (0.29, 0.4)  |
|                   | SMOTE-SVM  | 0.97 (0.96, 0.98) | 0.81 (0.67, 0.92) | 0.64 (0.5, 0.81)  | 0.75 (0.58, 0.83) | 0.52 (0.36, 0.71) |
|                   | Hybrid-XGB | 0.97 (0.96, 0.98) | 0.86 (0.67, 0.95) | 0.64 (0.5, 0.81)  | 0.72 (0.61, 0.8)  | 0.51 (0.44, 0.67) |
| Balanced Accuracy | Sequential | 0.93 (0.91, 0.94) | 0.91 (0.86, 0.95) | 0.86 (0.77, 0.91) | 0.94 (0.8, 1)     | 0.94 (0.8, 1)     |
|                   | 2-STEP     | 0.93 (0.91, 0.94) | 0.92 (0.88, 0.95) | 0.81 (0.69, 0.88) | 0.87 (0.79, 0.95) | 0.87 (0.48, 1)    |
|                   | Up-RF      | 0.9 (0.86, 0.94)  | 0.89 (0.78, 0.94) | 0.81 (0.71, 0.9)  | 0.88 (0.73, 0.95) | 0.59 (0.5, 0.66)  |
|                   | SMOTE-SVM  | 0.92 (0.89, 0.95) | 0.88 (0.78, 0.92) | 0.83 (0.74, 0.92) | 0.87 (0.76, 0.95) | 0.81 (0.75, 0.87) |
|                   | Hybrid-XGB | 0.94 (0.91, 0.96) | 0.92 (0.78, 0.97) | 0.8 (0.72, 0.88)  | 0.9 (0.76, 0.95)  | 0.79 (0.68, 0.87) |
| Kappa             | Sequential | 0.86 (0.83, 0.88) | 0.82 (0.72, 0.91) | 0.71 (0.55, 0.82) | 0.88 (0.68, 1)    | 0.88 (0.68, 1)    |
|                   | 2-STEP     | 0.86 (0.83, 0.88) | 0.85 (0.77, 0.9)  | 0.63 (0.4, 0.77)  | 0.72 (0.54, 0.9)  | 0.72 (-0.03, 1)   |
|                   | Up-RF      | 0.85 (0.77, 0.9)  | 0.84 (0.68, 0.93) | 0.63 (0.45, 0.77) | 0.75 (0.55, 0.86) | 0.21 (0, 0.4)     |
|                   | SMOTE-SVM  | 0.84 (0.78, 0.88) | 0.8 (0.65, 0.91)  | 0.61 (0.48, 0.79) | 0.74 (0.56, 0.83) | 0.52 (0.35, 0.71) |
|                   | Hybrid-XGB | 0.85 (0.83, 0.89) | 0.85 (0.65, 0.95) | 0.62 (0.48, 0.79) | 0.7 (0.59, 0.79)  | 0.51 (0.44, 0.66) |




Figure 4.7: Top 5 Workflow Per-Class Evaluation Metrics by Metric

Table 4.13: Top Workflow Per-Class Evaluation Metrics and Ranks

| Workflow                 | Rank | HGSOC | CCOC  | ENOC  | MUOC  | LGSOC |
|--------------------------|------|-------|-------|-------|-------|-------|
| <b>F1-Score</b>          |      |       |       |       |       |       |
| Sequential               | 1    | 0.973 | 0.877 | 0.860 | 0.966 | 0.910 |
| 2-STEP                   | 2    | 0.973 | 0.899 | 0.755 | 0.788 | 0.733 |
| Up-RF                    | 3    | 0.972 | 0.849 | 0.652 | 0.757 | 0.362 |
| SMOTE-SVM                | 4    | 0.969 | 0.811 | 0.638 | 0.752 | 0.524 |
| Hybrid-XGB               | 5    | 0.971 | 0.860 | 0.643 | 0.716 | 0.515 |
| <b>Balanced Accuracy</b> |      |       |       |       |       |       |
| Sequential               | 1    | 0.926 | 0.907 | 0.856 | 0.943 | 0.943 |
| Hybrid-XGB               | 2    | 0.935 | 0.922 | 0.802 | 0.899 | 0.791 |
| 2-STEP                   | 6    | 0.926 | 0.917 | 0.814 | 0.875 | 0.872 |
| SMOTE-SVM                | 10   | 0.923 | 0.875 | 0.826 | 0.869 | 0.814 |
| Up-RF                    | 14   | 0.905 | 0.894 | 0.808 | 0.878 | 0.590 |
| <b>Kappa</b>             |      |       |       |       |       |       |
| Sequential               | 1    | 0.858 | 0.815 | 0.712 | 0.877 | 0.877 |
| 2-STEP                   | 2    | 0.858 | 0.850 | 0.635 | 0.716 | 0.718 |
| SMOTE-SVM                | 3    | 0.839 | 0.800 | 0.614 | 0.741 | 0.516 |
| Up-RF                    | 4    | 0.850 | 0.840 | 0.630 | 0.746 | 0.213 |
| Hybrid-XGB               | 6    | 0.855 | 0.852 | 0.620 | 0.702 | 0.506 |




Figure 4.8: Top 5 Workflow Per-Class Evaluation Metrics by Metric

Misclassified cases from a previous step of the sequence of classifiers are not included in subsequent steps of the training set CV folds. Thus, we cannot piece together the test set predictions from the sequential and two-step algorithms to obtain overall metrics.

### **4.3 Confirmation Set**

Now we'd like to see how our best five workflows perform in the confirmation set. The class-specific F1-scores will be used. The top performing method will be selected for gene optimization.

Table 4.14: Evaluation Metrics on Confirmation Set Models

| Method     | Metric            | Overall | Histotypes |       |       |       |       |
|------------|-------------------|---------|------------|-------|-------|-------|-------|
|            |                   |         | HGSOC      | CCOC  | ENOC  | MUOC  | LGSOC |
| Sequential | Accuracy          | 0.829   | 0.866      | 0.969 | 0.883 | 0.966 | 0.974 |
|            | Sensitivity       | 0.584   | 0.953      | 0.861 | 0.467 | 0.556 | 0.083 |
|            | Specificity       | 0.924   | 0.697      | 0.982 | 0.966 | 0.984 | 0.990 |
|            | F1-Score          | 0.604   | 0.904      | 0.861 | 0.571 | 0.577 | 0.105 |
|            | Balanced Accuracy | 0.754   | 0.825      | 0.922 | 0.717 | 0.770 | 0.537 |
|            | Kappa             | 0.646   | 0.685      | 0.844 | 0.508 | 0.559 | 0.093 |
| 2-STEP     | Accuracy          | 0.838   | 0.866      | 0.970 | 0.891 | 0.977 | 0.972 |
|            | Sensitivity       | 0.613   | 0.953      | 0.875 | 0.486 | 0.667 | 0.083 |
|            | Specificity       | 0.926   | 0.697      | 0.982 | 0.972 | 0.990 | 0.989 |
|            | F1-Score          | 0.635   | 0.904      | 0.869 | 0.598 | 0.706 | 0.100 |
|            | Balanced Accuracy | 0.769   | 0.825      | 0.929 | 0.729 | 0.828 | 0.536 |
|            | Kappa             | 0.666   | 0.685      | 0.852 | 0.538 | 0.694 | 0.086 |
| Up-RF      | Accuracy          | 0.835   | 0.857      | 0.975 | 0.883 | 0.974 | 0.981 |
|            | Sensitivity       | 0.613   | 0.972      | 0.875 | 0.383 | 0.667 | 0.167 |
|            | Specificity       | 0.918   | 0.633      | 0.988 | 0.983 | 0.987 | 0.997 |
|            | F1-Score          | 0.648   | 0.900      | 0.887 | 0.522 | 0.679 | 0.250 |
|            | Balanced Accuracy | 0.765   | 0.802      | 0.931 | 0.683 | 0.827 | 0.582 |
|            | Kappa             | 0.646   | 0.654      | 0.873 | 0.466 | 0.665 | 0.243 |
| SMOTE-SVM  | Accuracy          | 0.827   | 0.866      | 0.958 | 0.888 | 0.972 | 0.970 |
|            | Sensitivity       | 0.650   | 0.939      | 0.861 | 0.477 | 0.556 | 0.417 |
|            | Specificity       | 0.927   | 0.725      | 0.970 | 0.970 | 0.990 | 0.981 |
|            | F1-Score          | 0.656   | 0.902      | 0.821 | 0.586 | 0.625 | 0.345 |
|            | Balanced Accuracy | 0.788   | 0.832      | 0.916 | 0.723 | 0.773 | 0.699 |
|            | Kappa             | 0.651   | 0.690      | 0.797 | 0.525 | 0.611 | 0.330 |
| Hybrid-XGB | Accuracy          | 0.830   | 0.869      | 0.961 | 0.893 | 0.967 | 0.970 |
|            | Sensitivity       | 0.662   | 0.943      | 0.861 | 0.458 | 0.630 | 0.417 |
|            | Specificity       | 0.928   | 0.725      | 0.974 | 0.979 | 0.982 | 0.981 |
|            | F1-Score          | 0.657   | 0.905      | 0.832 | 0.587 | 0.618 | 0.345 |
|            | Balanced Accuracy | 0.795   | 0.834      | 0.917 | 0.719 | 0.806 | 0.699 |
|            | Kappa             | 0.657   | 0.696      | 0.810 | 0.531 | 0.601 | 0.330 |

Evaluation Metrics on Confirmation Set Models




Figure 4.9: Evaluation Metrics on Confirmation Set Models




Figure 4.10: Entropy vs. Predicted Probability in Confirmation Set




Figure 4.11: Gene Optimized Workflows Per-Class Metrics in Confirmation Set

## Confusion Matrices for Confirmation Set Models




Figure 4.12: Confusion Matrices for Confirmation Set Models

### 4.3.1 Sequential

ROC Curves for Sequential Model in Confirmation Set





Figure 4.13: ROC Curves for Sequential Model in Confirmation Set

### 4.3.2 2-STEP

ROC Curves for 2-STEP Model in Confirmation Set

A

HGSOC vs. non\_HGSOC



B

CCOC vs. ENOC vs. MUOC vs. LGSOC

AUC = 0.879




Figure 4.14: ROC Curves for 2-STEP Model in Confirmation Set

### 4.3.3 Up-RF




Figure 4.15: ROC Curve for Up-RF Model in Confirmation Set

#### 4.3.4 SMOTE-SVM




Figure 4.16: ROC Curve for SMOTE-SVM Model in Confirmation Set

### 4.3.5 Hybrid-XGB




Figure 4.17: ROC Curve for Hybrid-XGB Model in Confirmation Set

## 4.4 Gene Optimization

From Figure 4.9, we see that both Hybrid-XGB and SMOTE-SVM have the highest overall F1-Score and are relatively better at predicting the rarest histotype LGSOC (sensitivity = 0.42). Thus we choose both of these two workflows for gene optimization in the confirmation set. The optimal number of genes is determined by the highest average F1-Score across classes, including the overall metric in the average. We use an F1-score that is averaged across cross-validation folds (5) and class groups (6: Overall, HGSOC, CCOC, ENOC, MUOC, LGSOC) to compare performance between different number of genes selected.

#### 4.4.1 Hybrid-XGB




Figure 4.18: Gene Optimization for Hybrid-XGB Classifier using Averaged F1-Score

In the Hybrid-XGB classifier, the optimal number of genes is highlighted in red in Figure 4.18. Hence the optimal number of total genes used will be  $n=28+19=47$ .

The gene profile of the optimal set of genes used is displayed in Table 4.15. Core genes in the PrOTYPE and SPOT sets are annotated with green circles, and the added genes are annotated with yellow circles. The added genes are: HNF1B, TFF1, TPX2, WT1, IGFBP1, LGALS4, TFF3, KLK7, CYP2C18, GPR64, CAPN2, MET, GCNT3, GAD1, SLC3A1, EGFL6, C1orf173, DKK4 and C10orf116. Unused genes are annotated with red crosses.

Table 4.15: Gene Profile of Optimal Set in Hybrid-XGB Workflow

| Set | Genes   | PrOTYPE | SPOT | Optimal Set | Candidate Rank |
|-----|---------|---------|------|-------------|----------------|
|     | COL11A1 | v       |      | (*)         |                |
|     | CD74    | v       |      | (*)         |                |
|     | CD2     | v       |      | (*)         |                |
|     | TIMP3   | v       |      | (*)         |                |
|     | LUM     | v       |      | (*)         |                |
|     | CYTIP   | v       |      | (*)         |                |
|     | COL3A1  | v       |      | (*)         |                |

|      |          |   |     |        |
|------|----------|---|-----|--------|
| Core | THBS2    | v | (*) |        |
|      | TCF7L1   | v | v   | (*)    |
|      | HMGA2    | v |     | (*)    |
|      | FN1      | v |     | (*)    |
|      | POSTN    | v |     | (*)    |
|      | COL1A2   | v |     | (*)    |
|      | COL5A2   | v |     | (*)    |
|      | PDZK1IP1 | v |     | (*)    |
|      | FBN1     | v |     | (*)    |
|      | HIF1A    |   | v   | (*)    |
|      | CXCL10   |   | v   | (*)    |
|      | DUSP4    |   | v   | (*)    |
|      | SOX17    |   | v   | (*)    |
|      | MITF     |   | v   | (*)    |
|      | CDKN3    |   | v   | (*)    |
|      | BRCA2    |   | v   | (*)    |
|      | CEACAM5  |   | v   | (*)    |
|      | ANXA4    |   | v   | (*)    |
|      | SERPINE1 |   | v   | (*)    |
|      | CRABP2   |   | v   | (*)    |
|      | DNAJC9   |   | v   | (*)    |
|      | HNF1B    |   |     | (*) 1  |
|      | TFF1     |   |     | (*) 2  |
|      | TPX2     |   |     | (*) 3  |
|      | WT1      |   |     | (*) 4  |
|      | IGFBP1   |   |     | (*) 5  |
|      | LGALS4   |   |     | (*) 6  |
|      | TFF3     |   |     | (*) 7  |
|      | KLK7     |   |     | (*) 8  |
|      | CYP2C18  |   |     | (*) 9  |
|      | GPR64    |   |     | (*) 10 |
|      | CAPN2    |   |     | (*) 11 |
|      | MET      |   |     | (*) 12 |
|      | GCNT3    |   |     | (*) 13 |

|            |           |     |    |
|------------|-----------|-----|----|
| Candidates | GAD1      | (*) | 14 |
|            | SLC3A1    | (*) | 15 |
|            | EGFL6     | (*) | 16 |
|            | C1orf173  | (*) | 17 |
|            | DKK4      | (*) | 18 |
|            | C10orf116 | (*) | 19 |
|            | FUT3      | (x) | 20 |
|            | PBX1      | (x) | 21 |
|            | MUC5B     | (x) | 22 |
|            | KGFLP2    | (x) | 23 |
|            | IGKC      | (x) | 24 |
|            | IL6       | (x) | 25 |
|            | CPNE8     | (x) | 26 |
|            | CYP4B1    | (x) | 27 |
|            | TP53      | (x) | 28 |
|            | PAX8      | (x) | 29 |
|            | SERPINA5  | (x) | 30 |
|            | SENP8     | (x) | 31 |
|            | BRCA1     | (x) | 32 |
|            | STC1      | (x) | 33 |
|            | SEMA6A    | (x) | 34 |
|            | TSPAN8    | (x) | 35 |
|            | LIN28B    | (x) | 36 |
|            | EPAS1     | (x) | 37 |
|            | ATP5G3    | (x) | 38 |
|            | IGJ       | (x) | 39 |
|            | SCGB1D2   | (x) | 40 |
|            | BCL2      | (x) | 41 |
|            | ADCYAP1R1 | (x) | 42 |
|            | MAP1LC3A  | (x) | 43 |

#### 4.4.2 SMOTE-SVM




Figure 4.19: Gene Optimization for SMOTE-SVM Classifier using Averaged F1-Score

In the SMOTE-svm classifier, the optimal number of genes is achieved at the highest averaged F1-score with 41 genes added, highlighted in red in Figure 4.19. Hence the optimal number of total genes used will be  $n=28+41=69$ .

The gene profile of the optimal set of genes used is displayed in Table 4.16. Core genes in the PrOTYPE and SPOT sets are annotated with green circles, and the added genes are annotated with yellow circles. The added genes are: EGFL6, IGJ, IGKC, TP53, DKK4, MUC5B, SLC3A1, MAP1LC3A, IGFBP1, CPNE8, SERPINA5, SCGB1D2, STC1, EPAS1, BRCA1, KGFLP2, SENP8, BCL2, PBX1, KLK7, C10orf116, LIN28B, LGALS4, ADCYAP1R1, IL6, ZBED1, WT1, TFF1, GCNT3, HNF1B, TFF3, CYP4B1, CYP2C18, TSPAN8, FUT3, MET, ATP5G3, SEMA6A, GPR64, PAX8 and C1orf173. Unused genes are annotated with red crosses.

Table 4.16: Gene Profile of Optimal Set in SMOTE-SVM Workflow

| Set | Genes   | PrOTYPE | SPOT | Optimal Set | Candidate Rank |
|-----|---------|---------|------|-------------|----------------|
|     | COL11A1 | v       |      | (*)         |                |
|     | CD74    | v       |      | (*)         |                |
|     | CD2     | v       |      | (*)         |                |
|     | TIMP3   | v       |      | (*)         |                |
|     | LUM     | v       |      | (*)         |                |

|      |          |   |     |     |
|------|----------|---|-----|-----|
| Core | CYTIP    | v | (*) |     |
|      | COL3A1   | v | (*) |     |
|      | THBS2    | v | (*) |     |
|      | TCF7L1   | v | v   | (*) |
|      | HMGA2    | v | (*) |     |
|      | FN1      | v | (*) |     |
|      | POSTN    | v | (*) |     |
|      | COL1A2   | v | (*) |     |
|      | COL5A2   | v | (*) |     |
|      | PDZK1IP1 | v | (*) |     |
|      | FBN1     | v | (*) |     |
|      | HIF1A    |   | v   | (*) |
|      | CXCL10   |   | v   | (*) |
|      | DUSP4    |   | v   | (*) |
|      | SOX17    |   | v   | (*) |
|      | MITF     |   | v   | (*) |
|      | CDKN3    |   | v   | (*) |
|      | BRCA2    |   | v   | (*) |
|      | CEACAM5  |   | v   | (*) |
|      | ANXA4    |   | v   | (*) |
|      | SERPINE1 |   | v   | (*) |
|      | CRABP2   |   | v   | (*) |
|      | DNAJC9   |   | v   | (*) |
|      | EGFL6    |   | (*) | 1   |
|      | IGJ      |   | (*) | 2   |
|      | IGKC     |   | (*) | 3   |
|      | TP53     |   | (*) | 4   |
|      | DKK4     |   | (*) | 5   |
|      | MUC5B    |   | (*) | 6   |
|      | SLC3A1   |   | (*) | 7   |
|      | MAP1LC3A |   | (*) | 8   |
|      | IGFBP1   |   | (*) | 9   |
|      | CPNE8    |   | (*) | 10  |
|      | SERPINA5 |   | (*) | 11  |

|            |           |     |    |
|------------|-----------|-----|----|
| Candidates | SCGB1D2   | (*) | 12 |
|            | STC1      | (*) | 13 |
|            | EPAS1     | (*) | 14 |
|            | BRCA1     | (*) | 15 |
|            | KGFLP2    | (*) | 16 |
|            | SENP8     | (*) | 17 |
|            | BCL2      | (*) | 18 |
|            | PBX1      | (*) | 19 |
|            | KLK7      | (*) | 20 |
|            | C10orf116 | (*) | 21 |
|            | LIN28B    | (*) | 22 |
|            | LGALS4    | (*) | 23 |
|            | ADCYAP1R1 | (*) | 24 |
|            | IL6       | (*) | 25 |
|            | ZBED1     | (*) | 26 |
|            | WT1       | (*) | 27 |
|            | TFF1      | (*) | 28 |
|            | GCNT3     | (*) | 29 |
|            | HNF1B     | (*) | 30 |
|            | TFF3      | (*) | 31 |
|            | CYP4B1    | (*) | 32 |
|            | CYP2C18   | (*) | 33 |
|            | TSPAN8    | (*) | 34 |
|            | FUT3      | (*) | 35 |
|            | MET       | (*) | 36 |
|            | ATP5G3    | (*) | 37 |
|            | SEMA6A    | (*) | 38 |
|            | GPR64     | (*) | 39 |
|            | PAX8      | (*) | 40 |
|            | C1orf173  | (*) | 41 |
|            | GAD1      | (x) | 42 |
|            | CAPN2     | (x) | 43 |

### 4.4.3 Gene List Comparisons in Confirmation Set

We train the Hybrid-XGB and SMOTE-SVM workflows using the core and optimal gene lists in the training set. The models are evaluated on the confirmation set. Overall and per-class results are shown in Table [4.17](#). The gene lists are:

1. Core (n=28): among the overlapping genes, the core set from the PrOTYPE and SPOT lists
2. Optimal (n=47, 69): among the overlapping genes, the core set plus the additional number of genes that result in the optimal value for a selected evaluation metric, as assessed in Figure [4.18](#) and Figure [4.19](#)

Table 4.17: Model Comparisons using Different Gene Lists in Confirmation Set

| Method              | Metric            | Overall | Histotypes |       |       |       |       |
|---------------------|-------------------|---------|------------|-------|-------|-------|-------|
|                     |                   |         | HGSOC      | CCOC  | ENOC  | MUOC  | LGSOC |
| Hybrid-XGB, Optimal | Accuracy          | 0.836   | 0.872      | 0.966 | 0.893 | 0.966 | 0.977 |
|                     | Sensitivity       | 0.694   | 0.946      | 0.861 | 0.458 | 0.704 | 0.500 |
|                     | Specificity       | 0.930   | 0.729      | 0.979 | 0.979 | 0.977 | 0.986 |
|                     | F1-Score          | 0.684   | 0.907      | 0.849 | 0.587 | 0.633 | 0.444 |
|                     | Balanced Accuracy | 0.812   | 0.838      | 0.920 | 0.719 | 0.840 | 0.743 |
|                     | Kappa             | 0.670   | 0.703      | 0.830 | 0.531 | 0.616 | 0.433 |
| Hybrid-XGB, Core    | Accuracy          | 0.810   | 0.858      | 0.952 | 0.872 | 0.970 | 0.967 |
|                     | Sensitivity       | 0.613   | 0.929      | 0.875 | 0.393 | 0.704 | 0.167 |
|                     | Specificity       | 0.923   | 0.720      | 0.961 | 0.968 | 0.982 | 0.983 |
|                     | F1-Score          | 0.606   | 0.896      | 0.803 | 0.506 | 0.667 | 0.160 |
|                     | Balanced Accuracy | 0.768   | 0.825      | 0.918 | 0.680 | 0.843 | 0.575 |
|                     | Kappa             | 0.620   | 0.673      | 0.775 | 0.440 | 0.651 | 0.143 |
| SMOTE-SVM, Optimal  | Accuracy          | 0.821   | 0.860      | 0.958 | 0.879 | 0.974 | 0.972 |
|                     | Sensitivity       | 0.666   | 0.932      | 0.847 | 0.458 | 0.593 | 0.500 |
|                     | Specificity       | 0.925   | 0.720      | 0.972 | 0.963 | 0.990 | 0.981 |
|                     | F1-Score          | 0.665   | 0.898      | 0.819 | 0.557 | 0.653 | 0.400 |
|                     | Balanced Accuracy | 0.796   | 0.826      | 0.910 | 0.710 | 0.791 | 0.740 |
|                     | Kappa             | 0.640   | 0.676      | 0.795 | 0.490 | 0.639 | 0.386 |
| SMOTE-SVM, Core     | Accuracy          | 0.815   | 0.860      | 0.952 | 0.874 | 0.970 | 0.974 |
|                     | Sensitivity       | 0.672   | 0.903      | 0.889 | 0.514 | 0.556 | 0.500 |
|                     | Specificity       | 0.930   | 0.775      | 0.960 | 0.946 | 0.989 | 0.983 |
|                     | F1-Score          | 0.660   | 0.895      | 0.805 | 0.576 | 0.612 | 0.414 |
|                     | Balanced Accuracy | 0.801   | 0.839      | 0.924 | 0.730 | 0.772 | 0.741 |
|                     | Kappa             | 0.641   | 0.685      | 0.778 | 0.503 | 0.597 | 0.401 |




Figure 4.20: Gene List Comparisons of Evaluation Metrics in Confirmation Set

## 4.5 Validation Set

From the results in Table 4.17 and Figure 4.20, we see that the Hybrid-XGB, optimal workflow as the highest overall F1-Score and balanced accuracy. Thus, we choose the Hybrid-XGB model trained on the training set for the all overlap, optimal, and core gene lists, and evaluate performance in the validation set.

### 4.5.1 Evaluation Metrics

Table 4.18: Evaluation Metrics on Training Set Models in Validation Set

| Method                  | Metric            | Overall | Histotypes |       |       |       |       |
|-------------------------|-------------------|---------|------------|-------|-------|-------|-------|
|                         |                   |         | HGSOC      | CCOC  | ENOC  | MUOC  | LGSOC |
| Hybrid-XGB, All Overlap | Accuracy          | 0.889   | 0.905      | 0.969 | 0.955 | 0.978 | 0.972 |
|                         | Sensitivity       | 0.799   | 0.913      | 1.000 | 0.682 | 0.870 | 0.533 |
|                         | Specificity       | 0.958   | 0.877      | 0.966 | 0.985 | 0.980 | 0.980 |
|                         | F1-Score          | 0.715   | 0.938      | 0.831 | 0.750 | 0.667 | 0.390 |
|                         | Balanced Accuracy | 0.879   | 0.895      | 0.983 | 0.833 | 0.925 | 0.756 |
|                         | Kappa             | 0.725   | 0.739      | 0.815 | 0.726 | 0.656 | 0.377 |
| Hybrid-XGB, Optimal     | Accuracy          | 0.876   | 0.894      | 0.966 | 0.949 | 0.974 | 0.969 |
|                         | Sensitivity       | 0.789   | 0.904      | 0.971 | 0.625 | 0.913 | 0.533 |
|                         | Specificity       | 0.952   | 0.856      | 0.966 | 0.984 | 0.976 | 0.976 |
|                         | F1-Score          | 0.692   | 0.930      | 0.817 | 0.705 | 0.646 | 0.364 |
|                         | Balanced Accuracy | 0.870   | 0.880      | 0.969 | 0.804 | 0.944 | 0.755 |
|                         | Kappa             | 0.694   | 0.709      | 0.799 | 0.677 | 0.634 | 0.349 |
| Hybrid-XGB, Core        | Accuracy          | 0.843   | 0.877      | 0.954 | 0.923 | 0.966 | 0.966 |
|                         | Sensitivity       | 0.721   | 0.897      | 0.942 | 0.386 | 0.913 | 0.467 |
|                         | Specificity       | 0.937   | 0.805      | 0.955 | 0.981 | 0.968 | 0.975 |
|                         | F1-Score          | 0.615   | 0.919      | 0.760 | 0.496 | 0.583 | 0.318 |
|                         | Balanced Accuracy | 0.829   | 0.851      | 0.949 | 0.684 | 0.940 | 0.721 |
|                         | Kappa             | 0.611   | 0.661      | 0.736 | 0.458 | 0.568 | 0.303 |




Figure 4.21: Evaluation Metrics on Validation Set Models

## 4.5.2 Confusion Matrices

Confusion Matrix for Training Set Models evaluated on Validation Data

| A          |       | Hybrid-XGB, All Overlap |      |      |      |       |       |
|------------|-------|-------------------------|------|------|------|-------|-------|
| Prediction | HGSOC | 638                     | 0    | 21   | 0    | 3     |       |
|            | CCOC  | 20                      | 69   | 5    | 2    | 1     |       |
|            | ENOC  | 9                       | 0    | 60   | 1    | 2     |       |
|            | MUOC  | 14                      | 0    | 2    | 20   | 1     |       |
|            | LGSOC | 18                      | 0    | 0    | 0    | 8     |       |
|            |       | HGSOC                   | CCOC | ENOC | MUOC | LGSOC | Truth |
| B          |       | Hybrid-XGB, Optimal     |      |      |      |       |       |
| Prediction | HGSOC | 632                     | 1    | 24   | 0    | 3     |       |
|            | CCOC  | 20                      | 67   | 6    | 1    | 1     |       |
|            | ENOC  | 10                      | 0    | 55   | 1    | 2     |       |
|            | MUOC  | 16                      | 1    | 3    | 21   | 1     |       |
|            | LGSOC | 21                      | 0    | 0    | 0    | 8     |       |
|            |       | HGSOC                   | CCOC | ENOC | MUOC | LGSOC | Truth |
| C          |       | Hybrid-XGB, Core        |      |      |      |       |       |
| Prediction | HGSOC | 627                     | 0    | 35   | 0    | 3     |       |
|            | CCOC  | 26                      | 65   | 10   | 1    | 0     |       |
|            | ENOC  | 9                       | 1    | 34   | 1    | 4     |       |
|            | MUOC  | 18                      | 3    | 6    | 21   | 1     |       |
|            | LGSOC | 19                      | 0    | 3    | 0    | 7     |       |
|            |       | HGSOC                   | CCOC | ENOC | MUOC | LGSOC | Truth |

Figure 4.22: Confusion Matrix for Training Set Models evaluated on Validation Data

### 4.5.3 ROC Curves

#### 4.5.3.1 All Overlap




Figure 4.23: ROC Curves for Hybrid-XGB, All Overlap Set Model in Validation Set

#### 4.5.3.2 Optimal Set




Figure 4.24: ROC Curves for Hybrid-XGB, Optimal Set Model in Validation Set

#### 4.5.3.3 Core Set




Figure 4.25: ROC Curves for Hybrid-XGB, Core Set Model in Validation Set

## 4.5.4 Calibration Plots

### 4.5.4.1 All Overlap




Figure 4.26: Calibration Plots for Hybrid-XGB, All Overlap Set Model in Validation Set

#### 4.5.4.2 Optimal Set




Figure 4.27: Calibration Plots for Hybrid-XGB, Optimal Set Model in Validation Set

#### 4.5.4.3 Core Set




Figure 4.28: Calibration Plots for Hybrid-XGB, Core Set Model in Validation Set

#### 4.5.5 Final Model

A summary of the Hybrid-XGB, Optimal model results are shown in Figure [4.29](#).

# Hybrid-XGB, Optimal Set Summary in Validation Set




Figure 4.29: Validation Summary

## 4.5.5.1 Feature Importance

For a tree model, the “gain” is the most frequently used variable importance measure. It represents the fractional contribution of each feature to the model based on the total gain of this feature’s splits. A higher percentage means higher importance. Figure Figure 4.30 shows the gain for the top 10 features of the final model (Hybrid-XGB).




Figure 4.30: Final Model Variable Importance Plot

XGBoost builds a separate set of trees sequentially during each boosting round. We are able to calculate variable importance for each of the histotypes by including only the trees that are associated with their class index.

For example, the class index labels from our model are ordered CCOC, ENOC, HGSOC, LGSOC, MUOC. Assuming a total of 214 boosting rounds, every 5th tree is selected using these indices:

- CCOC: 1, 6, 11, ..., 1066
- ENOC: 2, 7, 12, ..., 1067
- HGSOC: 3, 8, 13, ..., 1068
- LGSOC: 4, 9, 14, ..., 1069
- MUOC: 5, 10, 15, ..., 1070

### Feature Importance by Histotype




Figure 4.31: Final Model Variable Importance Plot by Histotype

A SHAP importance plot incorporates directionality and distribution of the importance patterns (Lundberg and Lee 2017). Figure 4.32 illustrates a beeswarm plot for each histotype. We only show the top 10 features, with the most important at the top to the least important at the bottom. Positive SHAP values increase the prediction and negative SHAP values decrease the prediction. The colours represent feature values, with low values coloured purple and high values coloured yellow. Finally, the density of the points indicate whether there is a high concentration of observations distributed at a particular SHAP value.

For example, we see that for CCOC, higher values of the gene HNF1B increase the prediction and lower values decrease the prediction. Conversely, lower values of CYP2C18 decrease the prediction of HGSOC and vice versa.

SHAP Summary Plot by Histotype




Figure 4.32: Final Model SHAP Importance Plot

We can use a scatterplot to visualize how the actual values of individual features affect SHAP values, and subsequently the model predictions. The relationship may be linear, curved, or some other complex pattern. Fig Figure 4.33 and Figure 4.34 show the SHAP dependence plots for HNF1B and TFF3 respectively.

We know from the beeswarm plot that HNF1B was the most important feature for the CCOC class. SHAP values are relatively flat until HNF1B increases past -4, at which point the SHAP values start to increase sharply and prediction is strongly driven toward CCOC. At the same value of -4, the inverse pattern is observed for HGSOC.

SHAP Dependence Plot for HNF1B




Figure 4.33: SHAP Dependence Plot for HNF1B

TFF3 was the most important feature for ENOC prediction. SHAP values start to flip to positive after TFF3 increases past -5. On the other hand for CCOC, SHAP values start to flip to negative as TFF3 increases past -2, driving prediction away from the CCOC class.

SHAP Dependence Plot for TFF3




Figure 4.34: SHAP Dependence Plot for TFF3

## 4.5.5.2 Gene Membership

Table 4.19: Detailed Gene Membership

| Copy                             | PDF                              | Show                             | 50                               | entries                          | Search: <input type="text"/>     |                                  |                                  |                                  |                                  |                                  |  |
|----------------------------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|--|
| Genes                            | CS1                              | CS2                              | CS3                              | ProTYPE                          | SPOT                             | Core                             | All Overlap                      | Complete                         | Optimal-HXGB                     | Optimal-SSVM                     |  |
| <input type="text" value="All"/> | <input type="text" value="All"/> | <input type="text" value="All"/> | <input type="text" value="All"/> | <input type="text" value="All"/> | <input type="text" value="All"/> | <input type="text" value="All"/> | <input type="text" value="All"/> | <input type="text" value="All"/> | <input type="text" value="All"/> | <input type="text" value="All"/> |  |
| COL11A1                          | v                                | v                                | v                                | v                                |                                  | v                                | v                                | v                                | v                                | v                                |  |
| MARCH2                           | v                                | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| C10orf116                        | v                                | v                                | v                                |                                  |                                  |                                  | v                                |                                  | v                                | v                                |  |
| AADAT                            | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| TMTC1                            | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| HIF1A                            | v                                | v                                | v                                |                                  | v                                | v                                | v                                | v                                | v                                | v                                |  |
| NOTUM                            | v                                | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| GAD1                             | v                                | v                                | v                                |                                  |                                  |                                  | v                                |                                  | v                                |                                  |  |
| CCRL1                            | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| SPRY2                            | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| TPX2                             | v                                | v                                | v                                |                                  |                                  |                                  | v                                |                                  | v                                |                                  |  |
| COL6A1                           | v                                | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| ADD3                             | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| C4orf27                          | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| RGS7BP                           | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| SEPT1                            | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| KGFLP2                           | v                                | v                                | v                                |                                  |                                  |                                  | v                                |                                  |                                  | v                                |  |
| EGFL6                            | v                                | v                                | v                                |                                  |                                  |                                  | v                                |                                  | v                                | v                                |  |
| ARNTL2                           | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| HIBCH                            | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| LDHC_V3                          | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| BBOX1                            | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| CELSR2                           | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| IL15                             | v                                | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| KLK7                             | v                                | v                                | v                                |                                  |                                  |                                  | v                                |                                  | v                                | v                                |  |
| PBX1                             | v                                | v                                | v                                |                                  |                                  |                                  | v                                |                                  |                                  | v                                |  |
| TMEFF1                           | v                                | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| FGF20                            | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| LIN28B                           | v                                | v                                | v                                |                                  |                                  |                                  | v                                |                                  |                                  | v                                |  |
| XIAP                             | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| AZGP1                            | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| CD74                             | v                                | v                                | v                                | v                                |                                  | v                                | v                                | v                                | v                                | v                                |  |
| BGN                              | v                                | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| RASGRF1                          | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| ARG2                             | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| TFF3                             | v                                | v                                | v                                |                                  |                                  |                                  | v                                |                                  | v                                | v                                |  |
| MUC5B                            | v                                | v                                | v                                |                                  |                                  |                                  | v                                |                                  |                                  | v                                |  |
| CD2                              | v                                | v                                | v                                | v                                |                                  | v                                | v                                | v                                | v                                | v                                |  |
| MARCH1                           | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| CXCL10                           | v                                | v                                | v                                |                                  | v                                | v                                | v                                | v                                | v                                | v                                |  |
| HNRNPA0                          | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| FUT3                             | v                                | v                                | v                                |                                  |                                  |                                  | v                                |                                  |                                  | v                                |  |
| GAS1                             | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| IL11RA                           | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| DEFB1                            | v                                | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| IRF1                             | v                                | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| C12orf60                         | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| MST4                             | v                                |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |                                  |  |
| TIMP3                            | v                                | v                                | v                                | v                                |                                  | v                                | v                                | v                                | v                                | v                                |  |
| STC1                             | v                                | v                                | v                                |                                  |                                  |                                  | v                                |                                  |                                  | v                                |  |

Showing 1 to 50 of 869 entries

Previous

1

2

3

4

5

...

18

Next

Showing 1 to 50 of 869 entries

Previous [1](#) [2](#) [3](#) [4](#) [5](#) ... [18](#) Next

#### 4.5.6 Additional Explorations

Table 4.20: Clinicopath characteristics between correct and incorrect predictions of ENOC cases

| Characteristic            | Predicted ENOC Correctly N = 55 <sup>1</sup> | Missed ENOC N = 33 <sup>1</sup> |
|---------------------------|----------------------------------------------|---------------------------------|
| <b>Age at diagnosis</b>   | 53 (46, 63)                                  | 56 (51, 62)                     |
| <b>Tumour grade</b>       |                                              |                                 |
| low grade                 | 42 (93%)                                     | 17 (63%)                        |
| high grade                | 3 (6.7%)                                     | 10 (37%)                        |
| Unknown                   | 10                                           | 6                               |
| <b>FIGO tumour stage</b>  |                                              |                                 |
| I                         | 42 (78%)                                     | 20 (61%)                        |
| II-IV                     | 12 (22%)                                     | 13 (39%)                        |
| Unknown                   | 1                                            | 0                               |
| <b>Race</b>               |                                              |                                 |
| white                     | 50 (93%)                                     | 26 (90%)                        |
| non-white                 | 4 (7.4%)                                     | 3 (10%)                         |
| Unknown                   | 1                                            | 4                               |
| <b>ARID1A</b>             |                                              |                                 |
| absent/subclonal          | 10 (18%)                                     | 6 (18%)                         |
| present                   | 45 (82%)                                     | 27 (82%)                        |
| <b>WT1</b>                |                                              |                                 |
| diffuse (>50%)            | 2 (3.6%)                                     | 3 (9.1%)                        |
| focal (1-50%)             | 3 (5.5%)                                     | 0 (0%)                          |
| negative                  | 50 (91%)                                     | 30 (91%)                        |
| <b>TP53</b>               |                                              |                                 |
| mutated                   | 1 (1.8%)                                     | 4 (13%)                         |
| wild type                 | 54 (98%)                                     | 28 (88%)                        |
| Unknown                   | 0                                            | 1                               |
| <b>PR</b>                 |                                              |                                 |
| diffuse (>50%)            | 35 (64%)                                     | 11 (33%)                        |
| focal (1-50%)             | 9 (16%)                                      | 6 (18%)                         |
| negative                  | 11 (20%)                                     | 16 (48%)                        |
| <b>P16</b>                |                                              |                                 |
| abnormal block            | 2 (3.6%)                                     | 4 (12%)                         |
| abnormal complete absence | 13 (24%)                                     | 12 (36%)                        |
| normal                    | 40 (73%)                                     | 17 (52%)                        |
| <b>NAPSIN A</b>           |                                              |                                 |
| negative                  | 53 (96%)                                     | 32 (100%)                       |
| positive                  | 2 (3.6%)                                     | 0 (0%)                          |
| Unknown                   | 0                                            | 1                               |

<sup>1</sup>Median (Q1, Q3); n (%)

<sup>2</sup>Wilcoxon rank sum test; Fisher's exact test; Pearson's Chi-squared test




Figure 4.35: Volcano Plots of Validation Set Predictions

Boxplot of Most Differentially Expressed Genes




Figure 4.36: Boxplot of Most Differentially Expressed Genes

### Subtype Prediction Summary among Predicted HGSOc Samples




Figure 4.37: Subtype Prediction Summary among Predicted HGSOc Samples

# References

- Chawla, Nitesh V, Kevin W Bowyer, Lawrence O Hall, and W Philip Kegelmeyer. 2002. “SMOTE: Synthetic Minority over-Sampling Technique.” *Journal of Artificial Intelligence Research* 16: 321–57.
- Chen, Tianqi, and Carlos Guestrin. 2016. “XGBoost: A Scalable Tree Boosting System.” *arXiv*, ahead of print. <https://doi.org/10.48550/ARXIV.1603.02754>.
- Hastie, Trevor, Robert Tibshirani, and Martin Wainwright. 2015. “Statistical Learning with Sparsity.” *Monographs on Statistics and Applied Probability* 143 (143): 8.
- Hvitfeldt, Emil. 2025. *Themis: Extra Recipes Steps for Dealing with Unbalanced Data*. <https://doi.org/10.32614/CRAN.package.themis>.
- Lundberg, Scott M, and Su-In Lee. 2017. “A Unified Approach to Interpreting Model Predictions.” *Advances in Neural Information Processing Systems* 30.
- Pihur, Vasyli, Susmita Datta, and Somnath Datta. 2009. “RankAggreg, an r Package for Weighted Rank Aggregation.” *BMC Bioinformatics* 10 (1): 62.
- Platt, John et al. 1999. “Probabilistic Outputs for Support Vector Machines and Comparisons to Regularized Likelihood Methods.” *Advances in Large Margin Classifiers* 10 (3): 61–74.
- Talhouk, Aline, Stefan Kommos, Robertson Mackenzie, et al. 2016. “Single-Patient Molecular Testing with NanoString nCounter Data Using a Reference-Based Strategy for Batch Effect Correction.” *PLOS ONE* 11 (4): e0153844. <https://doi.org/10.1371/journal.pone.0153844>.
- Wright, Marvin N., and Andreas Ziegler. 2017. “**Ranger**: A Fast Implementation of Random Forests for High Dimensional Data in *C++* and *R*.” *Journal of Statistical Software* 77 (1). <https://doi.org/10.18637/jss.v077.i01>.