

## **STATISTICS WORKSHEET-5**

**Q1 to Q10 are MCQs with only one correct answer. Choose the correct option.**

1. Using a goodness of fit, we can assess whether a set of obtained frequencies differ from a set of frequencies.
  - a) Mean
  - b) Actual
  - c) Predicted
  - d) Expected
2. Chi-square is used to analyse
  - a) Score
  - b) Rank
  - c) Frequencies
  - d) All of these
3. What is the mean of a Chi Square distribution with 6 degrees of freedom?
  - a) 4
  - b) 12
  - c) 6
  - d) 8
4. Which of these distributions is used for a goodness of fit testing?
  - a) Normal distribution
  - b) Chi-squared distribution
  - c) Gamma distribution
  - d) Poisson distribution
5. Which of the following distributions is Continuous
  - a) Binomial Distribution
  - b) Hypergeometric Distribution
  - c) F Distribution
  - d) Poisson Distribution
6. A statement made about a population for testing purpose is called?
  - a) Statistic
  - b) Hypothesis
  - c) Level of Significance
  - d) Test Statistic
7. If the assumed hypothesis is tested for rejection considering it to be true is called?
  - a) Null Hypothesis
  - b) Statistical Hypothesis
  - c) Simple Hypothesis
  - d) Composite Hypothesis
8. If the Critical region is evenly distributed then the test is referred as?
  - a) Two tailed
  - b) One tailed
  - c) Three tailed
  - d) Zero tailed
9. Alternative Hypothesis is also called as?
  - a) Composite hypothesis

- b) Research Hypothesis
  - c) Simple Hypothesis
  - d) Null Hypothesis
10. In a Binomial Distribution, if 'n' is the number of trials and 'p' is the probability of success, then the mean value is given by \_
- a) np
  - b) n

### Answers

- 1. Expected (d)
- 2. Frequencies (C)
- 3. 6 (C )
- 4. Chisqared distribution (B)
- 5. F Distribution ( C)
- 6. Hypothesis( B )
- 7. Null Hypothesis (A )
- 8. One tailed (B)
- 9. Research Hypothesis (B )
- 10. np (A)