

## Sex Differences in Stroke Epidemiology A Systematic Review

Peter Appelros, MD, PhD; Birgitta Stegmayr, PhD; Andreas Terént, MD, PhD

**Background and Purpose**—Epidemiological studies, mainly based on Western European surveys, have shown that stroke is more common in men than in women. In recent years, sex-specific data on stroke incidence, prevalence, subtypes, severity and case-fatality have become available from other parts of the world. The purpose of this article is to give a worldwide review on sex differences in stroke epidemiology.

**Methods**—We searched PubMed, tables-of-contents, review articles, and reference lists for community-based studies including information on sex differences. In some areas, such as secular trends, ischemic subtypes and stroke severity, noncommunity-based studies were also reviewed. Male/female ratios were calculated.

**Results**—We found 98 articles that contained relevant sex-specific information, including 59 incidence studies from 19 countries and 5 continents. The mean age at first-ever stroke was 68.6 years among men, and 72.9 years among women. Male stroke incidence rate was 33% higher and stroke prevalence was 41% higher than the female, with large variations between age bands and between populations. The incidence rates of brain infarction and intracerebral hemorrhage were higher among men, whereas the rate of subarachnoidal hemorrhage was higher among women, although this difference was not statistically significant. Stroke tended to be more severe in women, with a 1-month case fatality of 24.7% compared with 19.7% for men.

**Conclusions**—Worldwide, stroke is more common among men, but women are more severely ill. The mismatch between the sexes is larger than previously described. (*Stroke*. 2009;40:1082-1090.)

**Key Words:** stroke ■ epidemiology ■ sex ■ gender

A previous review article, mainly based on studies from Western Europe, showed that stroke incidence was about 30% higher in men than in women. For cerebral infarction the excess was 45%, whereas there was little difference between the sexes regarding intracerebral hemorrhage. For subarachnoidal hemorrhage, the relationship between the sexes was opposite, with a male deficit of about 50%.<sup>1</sup> Male patients are on average younger than female when they got their first stroke.<sup>2</sup> During recent years, a considerable number of articles have been published which add knowledge on epidemiological differences between the sexes in different parts of the world. The purpose of this review is to give an update on the current knowledge within this field. We have extended this review to also include data on stroke prevalence, severity, trends and case fatality, which hitherto have been given little attention in review articles.

### Methods

We searched PubMed using the algorithms (incidence[ti] OR epidemiology[ti] OR prevalence[ti]) AND (cerebrovascular[ti] OR stroke[ti]); subtype[ti] AND (cerebrovascular[ti] OR stroke[ti]). We also used existing review articles on community-based incidence and prevalence studies,<sup>3-11</sup> as well as reference lists in articles. Furthermore, tables-of-contents in relevant journals have been scrutinized. The period of manual search reached from the year 2000 and lasted

until the end of 2007, while the above mentioned review articles cover earlier periods. The names of the journals that were manually searched are listed in the supplemental Appendix (available online at <http://stroke.ahajournals.org>). The ambition was to include comparable community-based studies only.<sup>3,12</sup> The main criteria of such studies are: complete community-based case ascertainment, multiple overlapping sources, standard definition of stroke (WHO), first-ever strokes only, prospective design, all ages included, standard presentation (mid-decade age bands). All incidence studies (except one, see below) fulfilled these criteria. For prevalence studies, we required a community-based procedure using telephone interviews or questionnaires. Regarding stroke trends and ischemic subtypes, other studies than community-based have been accepted as well. When noncommunity-based studies have been referred to, this has been emphasized. Only articles in English have been included in this review.

Most results in this review are given as male/female (m/f) ratios (eg, incidence rate ratios and prevalence ratios). When calculating the overall age-standardized m/f ratios, it is not possible to divide the overall (all ages) male rate with the overall female rate because males and females have different age distributions. To obtain the total incidence rate ratio for each study, the male and female crude incidence rates have to be standardized to the age structure of the total (male+female) population. In our case, standardization was performed to the WHO world standard population<sup>13</sup> using the direct method.<sup>14</sup> The number of strokes as well as the number of people at risk (person-years) has to be known for each age band. If this information was not

Received October 24, 2008; final revision received November 25, 2008; accepted December 15, 2008.


From the Department of Neurology (P.A.), Örebro University Hospital, Örebro, Sweden; The National Board of Health and Welfare (B.S.), Stockholm, Sweden; and the Department of Medical Sciences (A.T.), Internal Medicine, Uppsala University Hospital, Uppsala, Sweden.

Correspondence to Dr Peter Appelros, Department of Neurology, Örebro University Hospital, SE-701 85 Örebro, Sweden. E-mail [peter.appelros@orebroll.se](mailto:peter.appelros@orebroll.se)

© 2009 American Heart Association, Inc.

Stroke is available at <http://stroke.ahajournals.org>

DOI: 10.1161/STROKEAHA.108.540781



**Figure 1.** Male/female stroke incidence rate ratios with 95% CIs for different populations. Ages  $\geq 45$  years are included. (Adjusted to the WHO world population.) Test for heterogeneity:  $\chi^2=144.18$ ,  $df=43$ ,  $P<0.00001$ ,  $I^2=70\%$ ; test for overall effect:  $Z=6.48$  ( $P<0.00001$ ).

obtainable, the study could not be included in the overall age-standardized results.

The STATA software<sup>15</sup> was used to calculate the standardized m/f rate ratios. For the stroke incidence studies, only studies which have given number of strokes separately for men and women, along with person-years at risk for the age bands—45 to 54, 55 to 64, 65 to 74, 75 to 84, and  $\geq 85$ —were included in the results. Because of the lack of information in many studies, the age bands below 45 years could not be included in this calculation. Theoretically, this means that the total rate ratio is biased against younger individuals, but in practice, these age bands have little influence on the total rate ratio, as only 2.9% of all strokes occur in ages below 45 in these studies. (For example: for Örebro, Sweden, the world standardized m/f rate ratio is 1.33, whether ages  $<45$  years are included or not.) For stroke subtypes, however, we chose to also include ages below 45 years, because in these populations as many as 36% of all subarachnoid hemorrhages (SAH) occurred in ages below 45 years, and therefore this age group may heavily influence the total m/f SAH rate ratio.

To achieve the total age-specific ratios, data were pooled from individual studies (number of stroke cases for each study, as well as denominators were summarized). Confidence intervals (CIs) for different age bands were calculated according to the method described by Morris and Gardner.<sup>16</sup>

For the prevalence studies, as well as for main stroke types, the STATA software<sup>15</sup> was used to calculate ratios with CIs. For tests of heterogeneity, Review Manager (RevMan)<sup>17</sup> was used. Non-age-standardized data were used for the tests of heterogeneity.


## Results

### Incidence

In total, we found 59 different incidence studies from 19 countries and 5 continents which fulfilled our inclusion criteria.<sup>18–59</sup> With the exception of the Rochester study from the United States, which was included because of its long duration and careful design, these studies essentially comply with the requirements of comparable community-based studies described above.<sup>3,12</sup> There were 14 149 male and 16 255 female strokes, which means that females outnumbered men by the factor 1.15.

Figure 1 shows the world population age-adjusted m/f incidence rate ratios for 44 different populations, with 95% CIs, as well as the pooled result. The pooled age-adjusted rate ratio was 1.33 (95% CI, 1.30 to 1.37), meaning that stroke is 33% more incident in males than in females. For Europe (24 studies, 11 687 cases), the pooled age adjusted rate ratio was 1.24 (95% CI, 1.20 to 1.29), and for the Australasia and the Americas (20 studies, 10 304 cases) the rate ratio was 1.45 (95% CI, 1.39 to 1.51).

Figure 2 shows rate ratios for different age bands. There was a significant drop in the m/f incidence rate ratio between the 65 to 74 and the 75 to 84 age bands, and a tendency to further drop in the  $\geq 85$  age band. Further details of individual



**Figure 2.** Male/female stroke incidence rate ratios for different age bands with 95% CIs. Summary estimates. The same populations as in Figure 1 form the basis for this figure. Tests for heterogeneity: Ages 25 to 34:  $\chi^2=13.70$ ,  $df=15$ ,  $P=0.55$ ,  $I^2=0\%$ ; test for overall effect:  $Z=1.49$  ( $P=0.14$ ). Ages 35 to 44:  $\chi^2=19.92$ ,  $df=30$ ,  $P=0.92$ ,  $I^2=0\%$ ; test for overall effect:  $Z=3.92$  ( $P<0.0001$ ). Ages 45 to 54:  $\chi^2=46.52$ ,  $df=43$ ,  $P=0.33$ ,  $I^2=8\%$ ; test for overall effect:  $Z=8.00$  ( $P<0.00001$ ). Ages 55 to 64:  $\chi^2=80.65$ ,  $df=43$ ,  $P=0.0004$ ,  $I^2=47\%$ ; test for overall effect:  $Z=12.97$  ( $P<0.00001$ ). Ages 65 to 74:  $\chi^2=78.00$ ,  $df=43$ ,  $P=0.0009$ ,  $I^2=45\%$ ; test for overall effect:  $Z=15.23$  ( $P<0.00001$ ). Ages 75 to 84:  $\chi^2=65.05$ ,  $df=43$ ,  $P=0.02$ ,  $I^2=34\%$ ; test for overall effect:  $Z=8.18$  ( $P<0.00001$ ). Ages 85+ :  $\chi^2=64.73$ ,  $df=43$ ,  $P=0.02$ ,  $I^2=34\%$ ; test for overall effect:  $Z=1.81$  ( $P=0.07$ ).

studies are given in supplemental Table I (available online at <http://stroke.ahajournals.org>).

### Mean Age at First-Ever Stroke

Overall, the weighted mean age (according to study size) for men was 68.6 years, and 72.9 years for women, which means that women get their first strokes on an average 4.3 years later than men. The mean age for male patients varies between 60.8 years (Uzhgorod, Ukraine) and 75.3 years (Innherred, Norway), and for females between 65.3 years (Matão, Brazil) and 80.4 years (Lund, Sweden).

The studies from Eastern Europe (Novosibirsk, Tartu, Uzhgorod, Tblisi) average at 63.6/69.1 years (m/f), the studies from Western Europe at 71.8/76.0 years, and the studies from North America and Australasia average at 68.6/73.3.

### Secular Trends

In many populations, during the last 3 decades, there has been a decrease in stroke incidence in both sexes.<sup>20,32,60–63</sup> Some studies have found the declining stroke incidence to be greater in men.<sup>64–66</sup> A few studies did not find any significant changes over time.<sup>67,68</sup>

Although one study found that mean age at first stroke increased among women, and was unchanged among men,<sup>20</sup> many studies have shown a trend toward higher age in both sexes with time.<sup>24,39,67</sup> Some studies have shown an unchanged mean age,<sup>30,32</sup> and one study even a declining age at first-ever stroke.<sup>66</sup>

### Prevalence

We found 13 different prevalence studies which had given prevalence figure for men and women separately.<sup>5,9,69–78</sup> Figure 3 shows 9 of these studies presenting sex-specific data for the age bands 55 to 64, 65 to 74, 75 to 84 and  $\geq 85$  years. Age-adjusted<sup>13</sup> m/f prevalence ratios are shown for individual studies, as well as the pooled estimate (with 95% CIs). The pooled ratio was 1.41 (95% CI, 1.12 to 1.59), meaning that

Rate ratios (95% confidence intervals)	No. of studies	No. of patients m/f
1.42 (0.86-2.37)	19	41/29
1.49 (1.21-1.85)	31	221/148
1.42 (1.29-1.57)	44	945/711
1.48 (1.38-1.57)	44	2099/1665
1.43 (1.37-1.50)	44	3423/3327
1.21 (1.15-1.27)	44	2845/4150
1.09 (1.00-1.18)	44	850/1976

stroke was 41% more prevalent among men than women in these studies.

Supplemental Table II gives age-specific data for all 13 studies with 95% CIs, as well as pooled ratios. The m/f ratio peaks in the age band 65 to 74, while it tends to drop in the 75 to 84 age band, and significantly so in the  $\geq 85$  year age band.

### Types of Stroke

We found 17 studies which have given sex- and age-specific data on the main types of stroke, comprising 7783 male strokes and 8371 female strokes.<sup>41–43,45,48,52,55,58,59,79–84</sup> Figure 4 shows the age-adjusted m/f incidence rate ratios for different main types of stroke, with 95% CIs. For ischemic stroke, the ratio was 1.55 (95% CI, 1.48 to 1.61), for intracerebral hemorrhage 1.60 (95% CI, 1.47 to 1.74), for SAH 0.84 (95% CI, 0.69 to 1.04), and for stroke of undetermined cause 1.08 (0.98 to 1.20). Supplemental Table III gives age-specific values for individual populations, as well as the pooled results with CIs.


Studies on ischemic subtypes often give proportions or rates that are difficult to compare. Very few such studies have included age bands for subtypes. In supplemental Table IV we have compiled studies that have included information on numbers of strokes per sex for each subtype.<sup>25,52,82,85–94</sup> The table is based on 4808 male and 3869 female strokes. The most consistent finding is that cardioembolic stroke is proportionally more common in women, while large vessel and small vessel strokes are proportionally more common in men.

### Stroke Severity

Several studies have shown that women have more severe strokes than men,<sup>90,95,96</sup> although there were only trends in some studies.<sup>97,98</sup> The differences are modest—in our local study, the average score on the National Institutes of Health Stroke Scale (range 0 to 42)<sup>99</sup> was 10.0 for women and 8.2 for men ( $P=0.06$ ,  $n=377$ ).<sup>100</sup>

### Case Fatality

Figure 5 shows community-based studies that have accounted for 28-day (in some cases 30-day or 1-month) case fatality



**Figure 3.** Male/female stroke prevalence ratios for different populations with 95% CIs. Ages  $\geq 55$  years are included. (Adjusted to the WHO world population.) Test for heterogeneity:  $\chi^2=35.53$ ,  $df=8$ ,  $P<0.0001$ ,  $I^2=77\%$ ; test for overall effect:  $Z=9.55$  ( $P<0.00001$ ).

separately for men and women<sup>20,24,30,32,34,39,42,43,52-54,56,58,101-103</sup> Case fatality was higher among women than among men in 26 of 31 studies, and higher in male patients in only 3 studies. Supplemental Table V shows the case fatality in more detail for individual populations.


The pooled case fatality for men was 19.7%, and for women 24.7%. Thus, case fatality was 1.25 times higher among women (95% CI, 1.17 to 1.34). Twenty-one of the populations are included in this calculation (7561 male and 8639 female strokes).

## Discussion

Male sex may be a stronger risk factor for stroke than previously stated. Textbooks have often mentioned the incidence rates to be about 25% to 30% higher among men,<sup>104</sup> but the present review accounts for an overall risk increase for men by 33%. An explanation may be that community-based studies from new parts of the world have been published, and


that these studies have changed the picture. There was a significantly larger male predominance in studies from Australasia and the Americas compared to studies from Europe.

The  $\geq 85$ -year age band is open-ended, which has certain implications. In Sweden, the expected remaining length of life at 85 years is 5.24 years among men, and 6.50 years among women.<sup>105</sup> Thus, at 85 years, women have a 24% longer remaining lifetime than men that they are exposed to the risk of getting a stroke. The implication is that the male and female rates are not directly comparable in this age band. For example, in Örebro, Sweden, for ages  $\geq 85$  years, the age-specific incidence rate is 2878 per 100 000 per year for men, and 3285 for women.<sup>52</sup> In order to compensate for a shorter exposition time, the incidence rate for men has to be multiplied with 1.24. The adjusted incidence rate for men would then be 3569, which also would affect the total age adjusted rate ratios (m/f), in this case from 1.33 to 1.39. Consideration has not been taken to the  $\geq 85$  year age band



**Figure 4.** Male/female incidence rate ratios for main types of stroke with 95% CIs. (Adjusted to the WHO world population.) Summary estimates. Tests for heterogeneity: cerebral infarction:  $\chi^2=33.26$ ,  $df=16$ ,  $P=0.007$ ,  $I^2=52\%$ ; test for overall effect:  $Z=3.15$  ( $P=0.002$ ); intracerebral hemorrhage:  $\chi^2=22.14$ ,  $df=16$ ,  $P=0.14$ ,  $I^2=28\%$ ; test for overall effect:  $Z=4.69$  ( $P<0.00001$ ); subarachnoidal hemorrhage:  $\chi^2=16.11$ ,  $df=16$ ,  $P=0.45$ ,  $I^2=1\%$ ; test for overall effect:  $Z=2.47$  ( $P=0.01$ ); undetermined stroke:  $\chi^2=16.95$ ,  $df=16$ ,  $P=0.26$ ,  $I^2=17\%$ ; test for overall effect:  $Z=10.60$  ( $P<0.00001$ ).





\*Studies with an asterisk are included in the pooled results.

**Figure 5.** Male and female case fatality percentages at 1 month for different stroke populations. The order is based on average m/f case fatality. \*Studies with an asterisk are included in the pooled results.

problem in our review, however, because the population structure in the age band  $\geq 85$  years is not available for individual studies. Because of the different life expectancy in the sexes in ages  $\geq 85$  years, the overall m/f rate ratio is likely to be a few percentage units higher than the value accounted for.

The question is: Why do women have lower stroke incidence than men? One possible answer is genetic factors. We have not found support for this assumption in the literature. On the contrary, a recent systematic review found that women with stroke are more likely than men to have a parental history of stroke.<sup>106</sup> Another answer might be the positive effects of estrogen on the cerebral circulation.<sup>107</sup> A lifetime exposure to ovarian estrogens may protect against ischemic stroke, at least of the noncardioembolic type,<sup>108</sup> an effect that seems to cease with menopause.<sup>109</sup> These observations have been one reason for the use of postmenopausal hormone replacement therapy. Randomized controlled trials have, however, failed to show advantages for treated groups,<sup>110</sup> and the current recommendation is that postmenopausal hormone therapy should not be used in the primary prevention of stroke.<sup>111</sup> A third factor may be blood pressure. Studies have shown that blood pressure values are higher in men than women of similar ages.<sup>112,113</sup> Moreover, ischemic heart disease,<sup>86,95,114,115</sup> peripheral artery disease,<sup>86,90</sup> and cigarette smoking<sup>86,90,95,116,117</sup> are more prevalent among male stroke patients. These conditions are associated with large-vessel disease.

Textbook knowledge states that SAH is more common among women than men,<sup>104</sup> although this may not be the case in all populations. In a study from the WHO MONICA project, the populations from Eastern Europe and Finland had higher incidence rates of SAH in men than in women.<sup>118</sup> The MONICA study covers ages 25 through 64 years in most populations. A recent systematic review has shown that the female predominance begins in the upper part of this age

band, at the age of 55 through 64, and is most pronounced in the age band 75 through 85 years.<sup>11</sup> The same review found that the incidence of SAH was 1.24 times higher in women than in men, which in its inverted form,  $1/1.24=0.81$ , is not far from our figure of 0.84. It is not known why SAH is more common among women, but hormonal factors have been proposed.<sup>119</sup>

Atrial fibrillation is more prevalent among elderly people, and female stroke patients are older when they get their first stroke.<sup>86,90,114,120</sup> Several studies have shown that women are generally at higher risk than men for atrial fibrillation-related, cardioembolic stroke,<sup>115,121-124</sup> but a few studies also have failed to show any significant sex difference.<sup>125,126</sup> A biological reason for an increased risk for stroke in women with atrial fibrillation is still lacking,<sup>127</sup> but a study from the Swedish Stroke Register has shown that female patients with atrial fibrillation receive oral anticoagulant therapy less often than men.<sup>117</sup> The higher prevalence of embolic strokes among women may to a large part explain their higher stroke severity.

The compilation of ischemic subtypes (supplemental Table IV) suffers from low numbers and varying definitions, and that some studies have included an undetermined group as well (not accounted for in the table). Nonetheless, most studies show that embolic stroke is proportionally more common among women. Whether the incidence of cardioembolic stroke among higher in women is still an open question. Unfortunately, few studies have given age specific data on men and women separately for ischemic subtypes. We have only been able to find one such study,<sup>82</sup> and data from that study, as well as unpublished data from our own incidence study,<sup>52</sup> do not support that cardioembolic stroke is more common among women.

In almost all studies reviewed here, women have higher 1-month case fatality than men. It must be observed that confounders have not been adjusted for. Women are older

when they get their first stroke, and SAH and cardioembolic strokes are proportionally more common among women. Also, the results are not adjusted for differences in other baseline characteristics (eg, risk factors), which may play a role. Therefore, this result is not in conflict with prognostic studies, where multivariate analysis emphasizes other factors, and in which sex rarely becomes an independent factor.<sup>128</sup>

One obvious limitation of this review is that the distribution of contributing studies is geographically uneven. Africa, Asia and South America are underrepresented. This is of importance, because our results show that the m/f ratios may differ between different parts of the world. Also, because of limitations in our search strategy, we may have missed some studies. Despite this, we think that the main trends in this review are supported by most studies. Some of these epidemiological factors, as for example age distribution and distribution of stroke types, may however vary considerably between places. The case fatality ratios are not age-standardized, and age differences may therefore bias the results. Differences in stroke care may also strongly affect case fatality.

Our review has confirmed that stroke is more common among men than women. The difference tends to decrease with age. Women get their first stroke about four and a half years later than men. The incidence of SAH tends to be higher in women, even if the results of our review were not statistically significant. Both brain infarction and intracerebral hemorrhage are more common in men, but cardioembolic stroke, which is more severe, accounts for a larger proportion of strokes among women. Strokes are more severe in women, and case fatality at one month is higher among women.

### Acknowledgments

We thank Anders Magnuson, BSc, for statistical advice.

### Sources of Funding

This study was supported by grants from the Research Funds of the Neurology Department, Örebro University Hospital, Örebro, Sweden.

### Disclosures

None.

### References

- Haberman S, Capildeo R, Rose FC. Sex differences in the incidence of cerebrovascular disease. *J Epidemiol Community Health*. 1981;35:45–50.
- Wyller TB. Stroke and gender. *J Gend Specif Med*. 1999;2:41–45.
- Malmgren R, Warlow C, Bamford J, Sandercock P. Geographical and secular trends in stroke incidence. *Lancet*. 1987;2:1196–1200.
- Sudlow CL, Warlow CP. Comparable studies of the incidence of stroke and its pathological types: results from an international collaboration. International Stroke Incidence Collaboration. *Stroke*. 1997;28:491–499.
- Di Carlo A, Launer LJ, Breteler MM, Fratiglioni L, Lobo A, Martinez-Lage J, Schmidt R, Hofman A. Frequency of stroke in Europe: a collaborative study of population-based cohorts. ILSA Working Group and the Neurologic Diseases in the Elderly Research Group. Italian Longitudinal Study on Aging. *Neurology*. 2000;54:S28–S33.
- Feigin VL, Lawes CM, Bennett DA, Anderson CA. Stroke epidemiology: a review of population-based studies of incidence, prevalence, and case-fatality in the late 20th century. *Lancet Neurol*. 2003;2:43–53.
- Saposnik G, Del Brutto OH. Stroke in South America: a systematic review of incidence, prevalence, and stroke subtypes. *Stroke*. 2003;34:2103–2107.
- Truelsen T, Piechowski-Józwiak B, Bonita R, Mathers C, Bogouslavsky J, Boysen G. Stroke incidence and prevalence in Europe: a review of available data. *Eur J Neurol*. 2006;13:581–598.
- Boix R, del Barrio JL, Saz P, Reñe R, Manubens JM, Lobo A, Gascón J, de Arce A, Díaz-Guzmán J, Bergareche A, Bermejo-Pareja F, de Pedro-Cuesta J. Stroke prevalence among the Spanish elderly: an analysis based on screening surveys. *BMC Neurol*. 2006;6:36.
- Lavados PM, Hennis AJ, Fernandes JG, Medina MT, Legetic B, Hoppe A, Sacks C, Jadue L, Salinas R. Stroke epidemiology, prevention, and management strategies at a regional level: Latin America and the Caribbean. *Lancet Neurol*. 2007;6:362–372.
- de Rooij NK, Linn FH, van der Plas JA, Algra A, Rinkel GJ. Incidence of subarachnoid haemorrhage: a systematic review with emphasis on region, age, gender and time trends. *J Neurol Neurosurg Psychiatry*. 2007;78:1365–1372.
- Sudlow CL, Warlow CP. Comparing stroke incidence worldwide: What makes studies comparable? *Stroke*. 1996;27:550–558.
- Ahmad OB, Boschi-Pinto C, Lopez AD, Murray CJL, Lozano L, Inoue M. Age standardization of rates. A new WHO standard. World Health Organization. Available from: [http://www.who.int/infobase/publicfiles/who\\_standardpopulation\\_discussion31.pdf](http://www.who.int/infobase/publicfiles/who_standardpopulation_discussion31.pdf). Accessed August 22, 2008.
- Rothman KJ, Greenland S. Measures of disease frequency. In: Rothman KJ, Greenland S, eds. *Modern Epidemiology II ed*. Philadelphia: Lippincott Williams & Wilkins;1998:29–46.
- STATA. *Statistical Package v. 7.0*. College Station, Texas: Stata Corporation; 2001.
- Morris JA, Gardner MJ. Epidemiological studies. In: Altman DG, Machin D, Bryant TN, Gardner MJ, eds. *Statistics with Confidence*. II ed. London: BMJ Books;2000:57–72.
- The Nordic Cochrane Centre. *Review Manager (RevMan) version 5.0.15*. Copenhagen: The Cochrane Collaboration; 2008. Available from: <http://www.cc-ims.net/RevMan>. Accessed 2008–09-01.
- Homer D, Whisnant JP, Schoenberg BS. Trends in the incidence rates of stroke in Rochester, Minnesota, since 1935. *Ann Neurol*. 1987;22:245–251.
- Whisnant JP, Fitzgibbons JP, Kurland LT, Sayre GP. Natural history of stroke in Rochester, Minnesota, 1945 through 1954. *Stroke*. 1971;2:11–22.
- Brown RD, Whisnant JP, Sicks JD, O'Fallon WM, Wiebers DO. Stroke incidence, prevalence, and survival: secular trends in Rochester, Minnesota, through 1989. *Stroke*. 1996;27:373–380.
- Jørgensen HS, Plesner AM, Hübbe P, Larsen K. Marked increase of stroke incidence in men between 1972 and 1990 in Frederiksberg, Denmark. *Stroke*. 1992;23:1701–1704.
- Terént A. Increasing incidence of stroke among Swedish women. *Stroke*. 1988;19:598–603.
- Tanaka H, Ueda Y, Date C, Baba T, Yamashita H, Hayashi M, Shoji H, Owada K, Baba KI, Shibuya M, Kon T, Detels R. Incidence of stroke in Shibata, Japan: 1976–1978. *Stroke*. 1981;12:460–466.
- Morikawa Y, Nakagawa H, Naruse Y, Nishijo M, Miura K, Tabata M, Hirokawa W, Kagamimori S, Honda M, Yoshita K, Hayashi K. Trends in stroke incidence and acute case fatality in a Japanese rural area: the Oyabe study. *Stroke*. 2000;31:1583–1587.
- Herman B, Leyten AC, van Luijk JH, Frenken CW, Op de Coul AA, Schulte BP. Epidemiology of stroke in Tilburg, The Netherlands. The population-based stroke incidence register: 2. Incidence, initial clinical picture and medical care, and three-week case fatality. *Stroke*. 1982;13:629–634.
- Christie D. Stroke in Melbourne, Australia: an epidemiological study. *Stroke*. 1981;12:467–469.
- Thrift AG, Dewey HM, Macdonell RAL, McNeil JJ, Donnan GA. Stroke incidence on the east coast of Australia. The North East Melbourne stroke incidence study. *Stroke*. 2000;31:2087–2092.
- Kotila M. Declining incidence and mortality of stroke? *Stroke*. 1984;15:255–259.
- Bonita R, Beaglehole R, North JD. Event, incidence and case fatality rates of cerebrovascular disease in Auckland, New Zealand. *Am J Epidemiol*. 1984;120:236–243.
- Bonita R, Broad JB, Beaglehole R. Changes in stroke incidence and case-fatality in Auckland, New Zealand, 1981–91. *Lancet*. 1993;342:1470–1473.
- Rothwell PM, Coull AJ, Giles MF, Howard SC, Silver LE, Bull LM, Gutnikov SA, Edwards P, Mant D, Sackley CM, Farmer A, Sandercock PA, Dennis MS, Warlow CP, Bamford JM, Anslow P. Change in stroke incidence, mortality, case-fatality, severity, and risk factors in

- Oxfordshire, UK from 1981 to 2004 (Oxford Vascular Study). *Lancet*. 2004;363:1925–1933.
32. Feigin VL, Wiebers DO, Whisnant JP, O'Fallon WM. Stroke incidence and 30-day case-fatality rates in Novosibirsk, Russia, 1982 through 1992. *Stroke*. 1995;26:924–929.
  33. Giroud M, Beuriat P, Vion P, D'Ath PH, Dusserre L, Dumas R. Stroke in a French prospective population study. *Neuroepidemiology*. 1989;8:97–104.
  34. Hulter Åsberg K, Parrow A. Event, incidence, and fatality rates of cerebrovascular diseases in Enköping-Häbo, Sweden, 1986–1988. *Scand J Soc Med*. 1991;19:134–139.
  35. Ricci S, Celani MG, La Rosa F, Vitali R, Duca E, Ferraguzzi R, Paolotti M, Seppoloni D, Caputo N, Chiurulla C, Scaroni R, Signorini E. SEPIVAC: a community-based study of stroke incidence in Umbria, Italy. *J Neurol Neurosurg Psychiatry*. 1991;54:695–698.
  36. D'Alessandro G, Bottacchi E, Di Giovanni M, Martinazzo C, Sironi L, Lia C, Carenini L, Corso G, Gerbaz V, Polillo C, Compagnoni MP. Temporal trends of stroke in Valle d'Aosta, Italy. Incidence and 30-day fatality rates. *Neurol Sci*. 2000;21:13–18.
  37. Anderson CS, Jamrozik KD, Burvill PW, Chakera TM, Johnson GA, Stewart-Wynne EG. Ascertaining the true incidence of stroke: experience from the Perth Community Stroke Study, 1989–1990. *Med J Aust*. 1993;158:80–84.
  38. Czlonkowska A, Ryglewicz D, Weissbein T, Baranska-Gieruszczak M, Hier DB. A prospective community-based study of stroke in Warsaw, Poland. *Stroke*. 1994;25:547–551.
  39. Vibo R, Kõrv J, Roose M. The Third Stroke Registry in Tartu, Estonia: decline of stroke incidence and 28-day case-fatality rate since 1991. *Stroke*. 2005;36:2544–2548.
  40. Jiang B, Wang WZ, Chen H, Hong Z, Yang QD, Wu SP, Du XL, Bao QJ. Incidence and trends of stroke and its subtypes in China: results from three large cities. *Stroke*. 2006;37:63–68.
  41. Lauria G, Gentile M, Fassetta G, Casetta I, Agnoli F, Andreotta G, Barp C, Caneve G, Cavallaro A, Cielo R, Mongillo D, Mosca M, Olivieri P. Incidence and prognosis of stroke in the Belluno province, Italy. First-year results of a community-based study. *Stroke*. 1995;26:1787–1793.
  42. Vemmos KN, Bots ML, Tsiouris PK, Zis VP, Grobbee DE, Stranjalis GS, Stamatelopoulos S. Stroke incidence and case fatality in southern Greece: the Arcadia stroke registry. *Stroke*. 1999;30:363–370.
  43. Carolei A, Marini C, Di Napoli M, Di Gianfilippo G, Santalucia P, Baldassarre M, De Matteis G, di Orio F. High stroke incidence in the prospective community-based L'Aquila registry (1994–1998). First year's results. *Stroke*. 1997;28:2500–2506.
  44. Du X, Sourbutts J, Cruickshank K, Summers A, Roberts N, Walton E, Holmes S. A community based stroke register in a high risk area for stroke in north west England. *J Epidemiol Community Health*. 1997;51:472–478.
  45. Kolominsky-Rabas PL, Sarti C, Heuschmann PU, Graf C, Siemonsen S, Neundoerfer B, Katalinic A, Lang E, Gassmann KG, von Stockert TR. A prospective community-based study of stroke in Germany—the Erlangen Stroke Project (ESPro): incidence and case fatality at 1, 3, and 12 months. *Stroke*. 1998;29:2501–2506.
  46. Ellekjer H, Holmen J, Indredavik B, Terént A. Epidemiology of stroke in Innherred, Norway, 1994 to 1996. Incidence and 30-day case-fatality rate. *Stroke*. 1997;28:2180–2184.
  47. Stewart JA, Dundas R, Howard RS, Rudd AG, Wolfe CD. Ethnic differences in incidence of stroke: prospective study with stroke register. *BMJ*. 1999;318:967–971.
  48. Di Carlo A, Inzitari D, Galati F, Baldereschi M, Giunta V, Grillo G, Furchi A, Manno V, Naso F, Vecchio A, Consoli D. A prospective community-based study of stroke in Southern Italy: the Vibo Valentia incidence of stroke study (VISS). Methodology, incidence and case fatality at 28 days, 3 and 12 months. *Cerebrovasc Dis*. 2003;16:410–417.
  49. Smadja D, Cabre P, May F, Fanon JL, Rene-Corail P, Riocreux C, Charpentier JC, Fournier P, Saint-Vil M, Ketterle J. ERMANCIA: Epidemiology of Stroke in Martinique, French West Indies: Part I: methodology, incidence, and 30-day case fatality rate. *Stroke*. 2001;32:2741–2747.
  50. Correia M, Silva MR, Matos I, Magalhães R, Lopes JC, Ferro JM, Silva MC. Prospective community-based study of stroke in Northern Portugal: incidence and case fatality in rural and urban populations. *Stroke*. 2004;35:2048–2053.
  51. Syme PD, Byrne AW, Chen R, Devenny R, Forbes JF. Community-based stroke incidence in a Scottish population: the Scottish Borders Stroke Study. *Stroke*. 2005;36:1837–1843.
  52. Appelros P, Nydevik I, Seiger Å, Terént A. High incidence rates of stroke in Örebro, Sweden: Further support for regional incidence differences within Scandinavia. *Cerebrovasc Dis*. 2002;14:161–168.
  53. Mihálka L, Smolanka V, Bulecza B, Mulesa S, Bereczki D. A population study of stroke in West Ukraine: incidence, stroke services, and 30-day case fatality. *Stroke*. 2001;32:2227–2231.
  54. Musolino R, La Spina P, Serra S, Postorino P, Calabró S, Savica R, Salemi G, Gallitto G. First-ever stroke incidence and 30-day case fatality in the Sicilian Aeolian archipelago, Italy. *Stroke*. 2005;36:2738–2741.
  55. Tsiskaridze A, Djibuti M, van Melle G, Lomidze G, Apridonidze S, Gaurashvili I, Piechowski-Józwiak B, Shakarishvili R, Bogousslavsky J. Stroke incidence and 30-day case-fatality in a suburb of Tbilisi: results of the first prospective population-based study in Georgia. *Stroke*. 2004;35:2523–2528.
  56. Hallström B, Jönsson AC, Nerbrand C, Norrving B, Lindgren A. Stroke incidence and survival in the beginning of the 21st century in Southern Sweden. Comparisons with the late 20th century and projections into the future. *Stroke*. 2008;39:10–15.
  57. Corbin DO, Poddar V, Hennis A, Gaskin A, Rambarat C, Wilks R, Wolfe CD, Fraser HS. Incidence and case fatality rates of first-ever stroke in a black Caribbean population: the Barbados Register of Strokes. *Stroke*. 2004;35:1254–1258.
  58. Lavados PM, Sacks C, Prina L, Escobar A, Tossi C, Araya F, Feuerhake W, Galvez M, Salinas R, Alvarez G. Incidence, 30-day case-fatality rate, and prognosis of stroke in Iquique, Chile: a 2-year community-based prospective study (PISCIS project). *Lancet*. 2005;365:2206–2215.
  59. Minelli C, Fu Fen L, Camara Minelli DP. Stroke incidence, prognosis, 30-day, and 1-year case fatality rates in Matão, Brazil: a population-based prospective study. *Stroke*. 2007;38:2906–2911.
  60. Sarti C, Tuomilehto J, Sivenius J, Kaarsalo E, Narva EV, Salmi K, Torppa J, Salomaa V. Declining trends in incidence, case-fatality and mortality of stroke in three geographic areas of Finland during 1983–1989. Results from the FINMONICA stroke register. *J Clin Epidemiol*. 1994;47:1259–1269.
  61. Sarti C, Stegmayr B, Tolonen H, Mahonen M, Tuomilehto J, Asplund K. Are changes in mortality from stroke caused by changes in stroke event rates or case fatality? Results from the WHO MONICA Project. *Stroke*. 2003;34:1833–1840.
  62. Ueshima H. Explanation for the Japanese paradox: prevention of increase in coronary heart disease and reduction in stroke. *J Atheroscler Thromb*. 2007;14:278–286.
  63. Rautio A, Eliasson M, Stegmayr B. Favorable trends in the incidence and outcome in stroke in nondiabetic and diabetic subjects: findings from the Northern Sweden MONICA Stroke Registry in 1985 to 2003. *Stroke*. 2008;39:3137–3144.
  64. Truelsen T, Prescott E, Grønbaek M, Schnohr P, Boysen G. Trends in stroke incidence: The Copenhagen City Heart Study. *Stroke*. 1997;28:1903–1907.
  65. Mayo NE, Nadeau L, Daskalopoulou SS, Cote R. The evolution of stroke in Quebec: a 15-year perspective. *Neurology*. 2007;68:1122–1127.
  66. Islam MS, Anderson CS, Hankey GJ, Hardie K, Carter K, Broadhurst R, Jamrozik K. Trends in incidence and outcome of stroke in Perth, Western Australia during 1989 to 2001: the Perth Community Stroke Study. *Stroke*. 2008;39:776–782.
  67. Terént A. Trends in stroke incidence and 10-year survival in Söderhamn, Sweden, 1975–2001. *Stroke*. 2003;34:1353–1358.
  68. Benatru I, Rouaud O, Durier J, Contegal F, Couvreur G, Bejot Y, Osseby GV, Ben Salem D, Ricolfi F, Moreau T, Giroud M. Stable stroke incidence rates but improved case-fatality in Dijon, France, from 1985 to 2004. *Stroke*. 2006;37:1674–1679.
  69. Bonita R, Solomon N, Broad JB. Prevalence of stroke and stroke-related disability: estimates from the Auckland stroke studies. *Stroke*. 1997;28:1898–1902.
  70. Reggion A, Rocca WA, Patti F, Grigoletto F, Meneghini F, Morgante L, Savettieri G, Salemi G, Cappello S, Di Perri R. Prevalence of stroke: a door-to-door survey in three Sicilian municipalities. Sicilian Neuroepidemiologic Study (SNES) Group. *Neuroepidemiology*. 1996;15:92–102.
  71. Mittelmark MB, Psaty BM, Rautaharju PM, Fried LP, Borhani NO, Tracy RP, Gardin JM, O'Leary DH. Prevalence of cardiovascular diseases among older adults: the Cardiovascular Health Study. *Am J Epidemiol*. 1993;137:311–317.



72. Geddes JM, Fear J, Tennant A, Pickering A, Hillman M, Chamberlain MA. Prevalence of self reported stroke in a population in northern England. *J Epidemiol Community Health*. 1996;50:140–143.
73. Melcon CM, Melcon MO. Prevalence of stroke in an Argentine community. *Neuroepidemiology*. 2006;27:81–88.
74. Prencipe M, Ferretti C, Casini AR, Santini M, Giubilei F, Culasso F. Stroke, disability, and dementia: results of a population survey. *Stroke*. 1997;28:531–536.
75. O'Mahony PG, Thomson RG, Dobson R, Rodgers H, James OF. The prevalence of stroke and associated disability. *J Public Health Med*. 1999;21:166–171.
76. Orlandi G, Gelli A, Fanucchi S, Tognoni G, Acerbi G, Murri L. Prevalence of stroke and transient ischaemic attack in the elderly population of an Italian rural community. *Eur J Epidemiol*. 2003;18:879–882.
77. Lin HC, Lin YJ, Liu TC, Chen CS, Chiu WT. Urbanization and stroke prevalence in Taiwan: analysis of a nationwide survey. *J Urban Health*. 2007;84:604–614.
78. Venketasubramanian N, Tan LC, Sahadevan S, Chin JJ, Krishnamoorthy ES, Hong CY, Saw SM. Prevalence of stroke among Chinese, Malay, and Indian Singaporeans: a community-based tri-racial cross-sectional survey. *Stroke*. 2005;36:551–556.
79. Giroud M, Milan C, Beuriat P, Gras P, Essayagh E, Arveux P, Dumas R. Incidence and survival rates during a two-year period of intracerebral and subarachnoid haemorrhages, cortical infarcts, lacunes and transient ischaemic attacks. The Stroke Registry of Dijon: 1985–1989. *Int J Epidemiol*. 1991;20:892–899.
80. Giroud M, Gras P, Chadan N, Beuriat P, Milan C, Arveux P, Dumas R. Cerebral haemorrhage in a French prospective population study. *J Neurol Neurosurg Psychiatry*. 1991;54:595–598.
81. Khan FA, Engström G, Jerntorp I, Pessah-Rasmussen H, Janzon L. Seasonal Patterns of Incidence and Case Fatality of Stroke in Malmö, Sweden: The STROMA Study. *Neuroepidemiology*. 2005;24:26–31.
82. Anderson CS, Jamrozik KD, Burvill PW, Chakera TM, Johnson GA, Stewart-Wynne EG. Determining the incidence of different subtypes of stroke: results from the Perth Community Stroke Study, 1989–1990. *Med J Aust*. 1993;158:85–89.
83. Thrift AG, Dewey HM, Macdonnell RA, McNeil JJ, Donnan GA. Incidence of the major stroke subtypes: initial findings from the North East Melbourne stroke incidence study (NEMESIS). *Stroke*. 2001;32:1732–1738.
84. Feigin V, Carter K, Hackett M, Barber PA, McNaughton H, Dyall L, Chen MH, Anderson C. Ethnic disparities in incidence of stroke subtypes: Auckland Regional Community Stroke Study, 2002–2003. *Lancet Neurol*. 2006;5:130–139.
85. Petty GW, Brown RD Jr, Whisnant JP, Sicks JD, O'Fallon WM, Wiebers DO. Ischemic stroke subtypes: a population-based study of incidence and risk factors. *Stroke*. 1999;30:2513–2516.
86. Arboix A, Oliveres M, García-Eroles L, Maragall C, Massons J, Targa C. Acute cerebrovascular disease in women. *Eur Neurol*. 2001;45:199–205.
87. Vemmos KN, Takis CE, Georgilis K, Zakopoulos NA, Lekakis JP, Papamichael CM, Zis VP, Stamatelopoulos S. The Athens stroke registry: results of a five-year hospital-based study. *Cerebrovasc Dis*. 2000;10:133–141.
88. Kitamura A, Nakagawa Y, Sato M, Iso H, Sato S, Imano H, Kiyama M, Okada T, Okada H, Iida M, Shimamoto T. Proportions of stroke subtypes among men and women  $\geq 40$  years of age in an urban Japanese city in 1992, 1997, and 2002. *Stroke*. 2006;37:1374–1378.
89. Kolominsky-Rabas PL, Weber M, Gefeller O, Neundorfer B, Heuschmann PU. Epidemiology of ischemic stroke subtypes according to TOAST criteria: incidence, recurrence, and long-term survival in ischemic stroke subtypes: a population-based study. *Stroke*. 2001;32:2735–2740.
90. Roquer J, Campello AR, Gomis M. Sex differences in first-ever acute stroke. *Stroke*. 2003;34:1581–1585.
91. Saposnik G, Gonzalez L, Lepera S, Luraschi A, Sica RE, Caplan LR, Rey RC. Southern Buenos Aires stroke project. *Acta Neurol Scand*. 2001;104:130–135.
92. Saposnik G, Caplan LR, Gonzalez LA, Baird A, Dashe J, Luraschi A, Llinas R, Lepera S, Linfante I, Chaves C, Kanis K, Sica RE, Rey RC. Differences in stroke subtypes among natives and caucasians in Boston and Buenos Aires. *Stroke*. 2000;31:2385–2389.
93. Grau AJ, Weimar C, Bugge F, Heinrich A, Goertler M, Neumaier S, Glahn J, Brandt T, Hacke W, Diener HC. Risk factors, outcome, and treatment in subtypes of ischemic stroke: the German stroke data bank. *Stroke*. 2001;32:2559–2566.
94. Uchino K, Risser JM, Smith MA, Moye LA, Morgenstern LB. Ischemic stroke subtypes among Mexican Americans and non-Hispanic whites: the BASIC Project. *Neurology*. 2004;63:574–576.
95. Jørgensen HS, Weber U, Nakayama H, Kammersgaard LP, Olsen TS. Differences in risk factor distribution, initial stroke severity, and outcome in men and women. The Copenhagen Stroke Study. *Cerebrovasc Dis*. 1999;9(Suppl 1):19.
96. Andersen MN, Andersen KK, Kammersgaard LP, Olsen TS. Sex differences in stroke survival: 10-year follow-up of the Copenhagen Stroke Study cohort. *J Stroke Cerebrovasc Dis*. 2005;14:215–220.
97. Appelros P, Nydevik I, Seiger Å, Terént A. Predictors of severe stroke: influence of pre-existing dementia and cardiac disorders. *Stroke*. 2002;33:2357–2362.
98. Barrett KM, Brott TG, Brown RD Jr, Frankel MR, Worrall BB, Silliman SL, Case LD, Rich SS, Meschia JF. Sex differences in stroke severity, symptoms, and deficits after first-ever ischemic stroke. *J Stroke Cerebrovasc Dis*. 2007;16:34–39.
99. Lyden P, Brott T, Tilley B, Welch KM, Mascha EJ, Levine S, Haley EC, Grotta J, Marler J. Improved reliability of the NIH Stroke Scale using video training. NINDS TPA Stroke Study Group. *Stroke*. 1994;25:2220–2226.
100. Appelros P, Terént A. Characteristics of the NIH Stroke Scale: Results from a population-based stroke cohort at baseline and after one year. *Cerebrovasc Dis*. 2004;17:21–27.
101. Bonita R, Anderson CS, Broad JB, Jamrozik KD, Stewart-Wynne EG, Anderson NE. Stroke incidence and case fatality in Australasia: a comparison of the Auckland and Perth population-based stroke registers. *Stroke*. 1994;25:552–557.
102. Ashok PP, Radhakrishnan K, Sridharan R, el-Mangoush MA. Incidence and pattern of cerebrovascular diseases in Benghazi, Libya. *J Neurol Neurosurg Psychiatry*. 1986;49:519–523.
103. Bejot Y, Rouaud O, Durier J, Caillier M, Marie C, Freysz M, Yeguiayan JM, Chantegret A, Osseby G, Moreau T, Giroud M. Decrease in the stroke case fatality rates in a French population-based twenty-year study: a comparison between men and women. *Cerebrovasc Dis*. 2007;24:439–444.
104. Sacco RL. Stroke risk factors: an overview. In: Norris JW, Hachinski V, eds. *Stroke Prevention*. New York: Oxford University Press;2001:17–42.
105. Statistics Sweden. *Population Statistics*. Stockholm: Statistiska Centralbyrån; 2007. Available from: <http://www.scb.se/>. Accessed October 8, 2007.
106. Touzé E, Rothwell PM. Sex differences in heritability of ischemic stroke: a systematic review and meta-analysis. *Stroke*. 2008;39:16–23.
107. Krause DN, Duckles SP, Pelligrino DA. Influence of sex steroid hormones on cerebrovascular function. *J Appl Physiol*. 2006;101:1252–1261.
108. de Leciñana MA, Egido JA, Fernández C, Martínez-Vila E, Santos S, Morales A, Martínez E, Pareja A, Álvarez-Sabin J, Casado I. Risk of ischemic stroke and lifetime estrogen exposure. *Neurology*. 2007;68:33–38.
109. Murphy SJ, McCullough LD, Smith JM. Stroke in the female: role of biological sex and estrogen. *ILAR J*. 2004;45:147–159.
110. Rossouw JE, Anderson GL, Prentice RL, LaCroix AZ, Kooperberg C, Stefanick ML, Jackson RD, Beresford SA, Howard BV, Johnson KC, Kotchen JM, Ockene J. Risks and benefits of estrogen plus progestin in healthy postmenopausal women: principal results From the Women's Health Initiative randomized controlled trial. *JAMA*. 2002;288:321–333.
111. Goldstein LB, Adams R, Alberts MJ, Appel LJ, Brass LM, Bushnell CD, Culebras A, Degra TJ, Gorelick PB, Guyton JR, Hart RG, Howard G, Kelly-Hayes M, Nixon JV, Sacco RL. Primary prevention of ischemic stroke: A guideline from the American Heart Association/American Stroke Association Stroke Council. *Stroke*. 2006;37:1583–1633.
112. Khoury S, Yarows SA, O'Brien TK, Sowers JR. Ambulatory blood pressure monitoring in a nonacademic setting: effects of age and sex. *Am J Hypertens*. 1992;5:616–623.
113. Wiinberg N, Høegholm A, Christensen HR, Bang LE, Mikkelsen KL, Nielsen PE, Svendsen TL, Kampmann JP, Madsen NH, Bentzon MW. 24-h ambulatory blood pressure in 352 normal Danish subjects, related to age and gender. *Am J Hypertens*. 1995;8:978–986.
114. Holroyd-Leduc JM, Kapral MK, Austin PC, Tu JV. Sex differences and similarities in the management and outcome of stroke patients. *Stroke*. 2000;31:1833–1837.
115. Fang MC, Singer DE, Chang Y, Hylek EM, Henault LE, Jensvold NG, Go AS. Gender differences in the risk of ischemic stroke and peripheral



- embolism in atrial fibrillation: the AnTicoagulation and Risk factors In Atrial fibrillation (ATRIA) study. *Circulation*. 2005;112:1687–1691.
116. Worrall BB, Johnston KC, Kongable G, Hung E, Richardson D, Gorelick PB. Stroke risk factor profiles in African American women: an interim report from the African-American Antiplatelet Stroke Prevention Study. *Stroke*. 2002;33:913–919.
  117. Glader EL, Stegmayr B, Norrving B, Terént A, Hulter-Åsberg K, Wester PO, Asplund K. Sex differences in management and outcome after stroke: a Swedish national perspective. *Stroke*. 2003;34:1970–1975.
  118. Ingall T, Asplund K, Mähönen M, Bonita R. A multinational comparison of subarachnoid hemorrhage epidemiology in the WHO MONICA stroke study. *Stroke*. 2000;31:1054–1061.
  119. Mhurchu CN, Anderson C, Jamrozik K, Hankey G, Dunbabin D. Hormonal factors and risk of aneurysmal subarachnoid hemorrhage: an international population-based, case-control study. *Stroke*. 2001;32:606–612.
  120. Phillips SJ, Whisnant JP, O'Fallon WM, Frye RL. Prevalence of cardiovascular disease and diabetes mellitus in residents of Rochester, Minnesota. *Mayo Clin Proc*. 1990;65:344–359.
  121. Stroke Prevention in Atrial Fibrillation Investigators. Risk factors for thromboembolism during aspirin therapy in patients with atrial fibrillation: the Stroke Prevention in Atrial Fibrillation Study. *J Stroke Cerebrovasc Dis*. 1995;5:147–157.
  122. Stewart S, Hart CL, Hole DJ, McMurray JJ. A population-based study of the long-term risks associated with atrial fibrillation: 20-year follow-up of the Renfrew/Paisley study. *Am J Med*. 2002;113:359–364.
  123. Wang TJ, Massaro JM, Levy D, Vasan RS, Wolf PA, D'Agostino RB, Larson MG, Kannel WB, Benjamin EJ. A risk score for predicting stroke or death in individuals with new-onset atrial fibrillation in the community: the Framingham Heart Study. *JAMA*. 2003;290:1049–1056.
  124. Friberg J, Scharling H, Gadsbøll N, Truelsen T, Jensen GB. Comparison of the impact of atrial fibrillation on the risk of stroke and cardiovascular death in women versus men (The Copenhagen City Heart Study). *Am J Cardiol*. 2004;94:889–894.
  125. Atrial Fibrillation Investigators. Risk factors for stroke and efficacy of antithrombotic therapy in atrial fibrillation: analysis of pooled data from five randomized controlled trials. *Arch Intern Med*. 1994;154:1449–1457.
  126. Kaarisalo MM, Immonen-Raiha P, Marttila RJ, Salomaa V, Kaarsalo E, Salmi K, Sarti C, Sivenius J, Torppa J, Tuomilehto J. Atrial fibrillation and stroke. Mortality and causes of death after the first acute ischemic stroke. *Stroke*. 1997;28:311–315.
  127. Lip GY, Watson T, Shantsila E. Anticoagulation for stroke prevention in atrial fibrillation: is gender important? *Eur Heart J*. 2006;27:1893–1894.
  128. Counsell C, Dennis M. Systematic review of prognostic models in patients with acute stroke. *Cerebrovasc Dis*. 2001;12:159–170.