

Self-critical rumination - A network approach

1. Introduction

Rumination is a repetitive thinking pattern about the causes, symptoms, circumstances and consequences of negative mood states and distress (Nolen-Hoeksema, 1991; E. R. Watkins, 2008). Rumination could maintain and exacerbate existing psychopathology (Butler & Nolen-Hoeksema, 1994; Nolen-Hoeksema et al., 1999; Whisman et al., 2020). Depressive rumination can also contribute to the onset or recurrence of several symptoms and disorders (Aldao & Nolen-Hoeksema, 2012; Lyubomirsky et al., 2015; Lyubomirsky & Nolen-Hoeksema, 1993; Nolen-Hoeksema et al., 2008; Ruscio et al., 2015; E. R. Watkins & Roberts, 2020; Whisman et al., 2020). As a trans-diagnostic characteristic (Nolen-Hoeksema & Watkins, 2011), rumination was found to predict the onset of major depressive disorder across one year in adolescent (Broderick & Korteland, 2004; Wilkinson et al., 2013) and adult samples (Nolen-Hoeksema et al., 2008), and was involved in generalized anxiety disorder (Nolen-Hoeksema, 2000) and PTSD (Moulds et al., 2020). Rumination predicted depression and anxiety (McLaughlin et al., 2014), bipolar disorder (Kovács et al., 2020), borderline personality disorder (Selby et al., 2008), social phobia (Abbott & Rapee, 2004), or psychosis (Hartley et al., 2013).

Rumination can lead to negative mood states and thoughts mutually amplifying and prolonging each other (E. R. Watkins & Roberts, 2020). However, consequences and effects of rumination are not specific to clinical samples, and can be identified in the background of the ruminative processes of healthy people (Ciesla & Roberts, 2007; Moulds et al., 2007). As such, rumination could interfere with effective problem-solving by reducing self-confidence (E. Watkins & Moulds, 2005); reduce willingness to engage in pleasant activities (Lyubomirsky et al., 2015); lead to elevated level of uncertainty (Ward et al., 2003), avoidance or personal difficulties (Hayes et al., 2004; Yasinski et al., 2019). Rumination is often operationalized as depressive rumination and consequently measured by the *Response Style Questionnaire* **RSQ**, or its shorter form the *Ruminative Response Scale*, **RRS** (Nolen-Hoeksema, 1991; Treynor, 2003). Two sub-factors within the **RRS**: the *brooding* and *reflective pondering* (or *reflection*) were identified (Treynor, 2003). Brooding is a maladaptive, passive dwelling on negative events and associated affects, while reflection is a more adaptive (or less pathological) form of thinking, a non-evaluative introspection towards one's mood. Several empirical findings support the adaptive-maladaptive distinction of brooding and reflection. Brooding alone increased the risk of substance use among people with history of illicit drug use (Memedovic et al., 2019), and with attentional disengagement difficulties, while controlling for depression and anxiety (Allard & Yaroslavsky, 2019). Apart from brooding and reflection, many other rumination types or facets were described in the last decades. People (especially with social phobia) tend to ruminate about their mood after a social event or performance (Abbott & Rapee, 2004). Individuals can dwell upon infuriating memories or on anger itself (Bushman et al., 2005; Pont et al., 2017). Repetitive thinking can target sadness as well (Peled & Moretti, 2009).

These results indicate that not just the style of thinking, but the content of these ruminative thoughts has special relevance (Smart et al., 2015). More recently, research has started to investigate the role that rumination plays in maintaining high levels of self-criticism (Kolubinski et al., 2017; Smart et al., 2015). Self-criticism is a form of negative self-evaluation with one harshly and hostilely judging oneself, especially in the context of mistakes, failures, unreach goals or unmet standards (Rose & Rimes, 2018; Shahar et al., 2014). Self-criticism was found to be a vulnerability factor in many forms psychopathology such as depressive and anxiety disorders, binge eating or non-suicidal self injury (Glassman et al., 2007; Rose & Rimes, 2018; Shahar et al., 2014). Rumination was found to be a significant mediator between self-criticism and depression across 2.5 years (Spasojević & Alloy, 2001), and between self-criticism and suicidal ideation over 3 months (O'Connor & Noyce, 2008). Thus, we can assume that rumination is one of those processes by which these negative self-devaluating thoughts are sustained and connected to psychopathology. Investigating rumination in the context of self-critical thoughts is paramount, especially in light of the results of a recent

meta-analysis which indicated that the harsh and persistent forms of self-criticism did not change through short-term interventions (Werner et al., 2019).

Self-critical rumination significantly differs from depressive rumination, mostly by the focus of thoughts (Smart et al., 2015). While in self-critical rumination, individuals concentrate on their lack of self-worth or those aspects of the self of which they are ashamed of, in depressive rumination one’s attention is directed towards the symptoms of depression (Nolen-Hoeksema, 2000). Indeed, in a recent study, self-critical rumination was strongly related to shame and predicted sensitivity to stress after a perceived failure (Milia et al., 2020). Positive meta-cognitions in self-critical rumination (Kolubinski et al., 2017) likely has less importance than in depressive rumination (Papageorgiou & Wells, 2001). One can assume that in depressive rumination the positive, while in self-critical rumination the negative meta-cognitions (e.g. “*I will get depressed if I don’t stop reviewing my self-critical thoughts*”) are those processes which maintain the ruminative process (Kolubinski et al., 2017). The Self-Critical Rumination Scale (**SCRS**, Smart et al. (2015)) showed incremental validity over other rumination scales (including the widely used **RRS** and **RSQ**) in predicting bipolar personality features and general distress. However, in the same study, **SCRS** and **RRS** were highly correlated, indicating that those individuals who ruminate in a more general or broader way are also likely to ruminate on numerous topics, including self-devaluating thoughts and vice versa. Previous work also supported the notion that individuals can show different forms of repetitive thought and they can ruminate and worry simultaneously, or they are able to passively brooding on something and adaptively reflect on other topics (E. R. Watkins, 2008).

Because of its multifaceted structure and the question of boundaries between the different subtypes and more importantly, patterns of rumination, it could be beneficial to investigate them in a network perspective. Modeling rumination as activation in a system of interacting components, such as a network would allow us to shift away from the distinctive labeling of brooding, reflection or self-critical aspects of rumination and would allow a more detailed, pattern-like view on the psychology of repetitive thoughts (see also at Bernstein et al., 2019). In this network approach, items are not assumed to have a common, underlying cause, nor can we expect that resulting networks fully map and cover the whole construct of trait rumination. Psychological networks such the ones presented below rather can be used to model components (items, and in our case scales) in dynamic interaction and to test past distinctions and classifications of rumination in a data-driven way. Contrary to latent variable models, network analysis does not seek for a common cause of co-occurrence of indicators (e.g. different items of a questionnaire) but shift the focus toward the relationships and interactions between variables, items or other constructs of interest (Epskamp, Kruis, et al., 2017). As opposed to *formative* models where observed scores define an attribute, or *reflective* models where a latent attribute is causing observables; in this alternative approach, psychological attributes (such as rumination) are conceptualized as *network* of directly related observables (Schmittmann et al., 2013). *Psychological networks* have their own characteristics which differentiate them from other kinds of networks of graph theory (Epskamp, Borsboom, et al. (2017a); Epskamp & Fried (2018); Newman (2010)). Vertices (also called nodes) usually represent some psychological measures or items of psychological surveys or questionnaires, while the edges of the network reflect the corresponding estimates of association derived from the data. Thus in a network model, the construct of rumination could be interpreted as a system in which some aspects relate or even stimulate each other, while other aspects have negative or inhibiting effects (Costantini et al., 2015). With this data-driven approach we can test how self-critical and depressive ruminative processes or the different items of the ruminative questionnaires cluster and relate to each other and we can gain important insights which could remain hidden by relying only on latent variable models (Bernstein et al., 2019). Having two rumination-related questionnaires, the **Ruminative Response Style (RRS)** and the **Self-critical Rumination Scale (SCRS)**, we assumed the following hypotheses:

- Network analysis will reveal their respective internal structure previously established in the literature (such as *RRS* having a two-scale structure: *reflection* and *brooding*; while *SCRS* will show a single-scale structure);
- on the other hand, we hypothesized that in their common network, items overlapping in scope will change these clear outlines. According to our hypothesis, items of the *RRS* scale *brooding* will have non-trivial association with the self-critical aspects of rumination;
- we also assumed a combined rumination network, representing the generic (more trait-like) and the

self-critical aspects of rumination will exhibit a three-component structure.

Previously, network modeling was successfully applied to investigate depressive rumination by testing the original 22-item version of **RRS** (Bernstein et al., 2019), but to the best of our knowledge no other rumination questionnaires were investigated, especially in terms of self-critical thoughts. We provide a glossary of the main network analysis related concepts in Table 8. of the Appendix.

2. Methods

2.1 Participants and procedure

Students from several Hungarian high schools participated in the study on a voluntary basis. The first stage of the recruitment process was to contact the heads of the schools to acquire institutional consent. Afterwards, classes were visited asking the students to participate. Since, the participants were under-aged, parents as well as students were issued with written informed consent letters. Following their consent, students were asked to fill out paper-pencil questionnaires in their own classroom within one class session. Participation in the study was anonymous and voluntary. Data collection was supervised by trained undergraduate students of psychology. No teaching staff were present. The total sample of the study comprises data from 851 participants (aged 14-19 years, females= 502 (59%), males=349 (41%) collected via convenience sampling method. The mean age was 16.1 (SD=1.1) years. The protocol was designed in accordance with the Declaration of Helsinki and the study was ethically approved by the Institutional Review Board (note: the name of the university is masked for review process).

2.2 Measures

The *Self-Critical Rumination Scale* **SCRS**; (Smart et al., 2015) was used to assess those thoughts that criticize the self for perceived failures, mistakes or bad habits in a prolonged and repetitive manner. The SCRS is a self-reported, one factor scale that consists of 10 items. Participants were asked to rate the statements (i.e. “*My attention is often focused on aspects of myself that I’m ashamed of*”; “*I criticize myself a lot for how I act around other people*”) on a 4-point Likert scale from 1 (not at all) to 4 (very much). The scale showed high internal consistency in the original version ($\alpha=0.92$). We obtained a satisfactory estimate ($\alpha=0.89$).

The short form of *Ruminative Response Scale* **RRS**; (Treyner, 2003) was used to measure mood dependent rumination, which consists of 10 items and quantifies ruminative thoughts via two dimensions (each composed of 5 items). The *brooding* subscale of **RRS** measures the maladaptive form or ruminative thinking (i.e. “*What am I doing to deserve this?*”), while the *reflective pondering* (or reflection) factor measures a more adaptive form of self-reflecting thinking (i.e. “*I write down what I am thinking and analyze it*”). Participants are asked to indicate on a 4-point Likert scale how often the items apply to themselves when they feel down, sad or depressed (1=almost never; 4=almost always). Internal consistency was found to be satisfactory for RRS ($\alpha=0.75$), as well as for the *reflection* ($\alpha=0.70$) and *brooding* ($\alpha=0.65$) scales. Besides the **SCRS** and **RRS** basic socio-demographic information (e.g. age, gender, parents’ level of education) was recorded.

2.3. Network modeling, descriptives and communities

In the current study, we constructed a dense, undirected network of rumination-related psychological constructs. The vertices represent the items of the *RRS* and *SCRS*, while edges are penalized partial correlations using LASSO (Least Absolute Shrinkage and Selection Operator) penalization (Friedman et al., 2007; Tibshirani, 1996).

We used individual and combined networks of *RRS* and *SCRS* with respective penalized correlation matrices using the *EBICglasso* function of the **qgraph** package (Epskamp et al., 2012). See our Git repository for other parameter specifications. In order to shrink the edges, achieve high specificity, and get a less dense network, we let the function to exclude any elements of the underlying precision matrix below a threshold value (*threshold* parameter set to TRUE).

The LASSO uses $L1$ (absolute) regularization (Friedman et al., 2007; Tibshirani, 1996). In the current study, the method resulted in a partial correlation network where edges represent direct associations between pairs of items, while the influence of all other items in the network are kept controlled. The regularization by $L1$ penalty lets trivially small partial correlations to be set to zero (McNally et al., 2017), thus ensure a *sparse* network which are more interpretable (Epskamp & Fried, 2018).

The tuning parameter of *gamma* in the graphical LASSO algorithm has been set to 0.5 value in accordance with (Bernstein et al., 2019; McNally et al., 2017). The final networks were chosen by the lowest values of the *extended Bayesian Information Criterion* (EBIC).

We applied the *goldbricker* function from the R package *networktools* (Jones, 2017) in order to search for pairs of items in the *RRS*, *SCRS* and their combined networks that were highly correlated ($r > 0.75$) and exhibited the same pattern with other items. After applying the LASSO regularization, no further dimensionality reduction was necessary based upon the *goldbricker* evaluation.

A comparative overview of the three networks was produced with their descriptives (degree distribution) and their centrality measures (strength, closeness, betweenness; Kolaczyk (2020)), and their One- and Two-Step Expected Influence (EI) measures to understand the cumulative importance of vertices (Bernstein et al., 2019; Donald J. Robinaugh, 2016; Heeren et al., 2018). In order to further elaborate the question of vertex importance, we identified *bridge nodes* using the functions provided by the *networktools* R package (Jones, 2017) to identify bridge vertices, in other words, connectors between sub-graphs (see an example at Bernstein et al., 2019). We estimated both one-step and two-step bridge expected influence measures which reflect the sum of edge weights connecting one vertex to other groups on vertices (or communities); and also express a vertex’s secondary influence on the rest of the network, respectively. Higher influence scores indicate that a certain vertex is more likely to active neighbouring vertices or communities. In order to estimate the Expected Influence metrics, we used the *expectedInf* function of the *networktools* R package. Also, to get an understanding of communities of vertices and to see if vertices can be grouped into neighbourhoods corresponding to the previously established sub-scale structure of *RRS* and *SCRS*; and to investigate their separation in their combined network, we used the community detection methods available in the *igraph* R package (Kolaczyk, 2020). When detecting possible communities of vertices, we followed guidelines by (Bernstein et al., 2019; Kolaczyk, 2020; Reichardt & Bornholdt, 2006). In order to mitigate the effect of choice of community detection algorithms, we performed more available algorithms from the *igraph* package multiple times ($B=1000$) in weighted and unweighted settings, too.

2.4 Internal Structure with an exponential random graph model

In order to measure the impact of of one item belonging to a questionnaire (*RRS*, *SCRS*) and their respective scales (*brooding*, *reflection* and *self-critical*) on the combined network structure, we constructed an *exponential random graph model* (ERGM) (Hunter, 2007). In our model, *endogenous* predictors (related to the nature of the network structure) included the **number of edges** in the network (*edges* member), and a generalization of triadic structures based on alternating sums of *k-triangles* (Goodreau et al., 2009; Hunter, 2007) to involve some higher-order global network structures. This predictor served as a reflection on **clustering** in the form of completed triangles of vertices. As estimator, we used *GWESP* (*geometrically weighted edgewise shared partner distribution*) (see Goodreau et al., 2009). *GWESP* has the advantage of taking a parametric form of the shared common vertex count distribution on each edge, giving each connecting vertex a declining positive probability of two selected vertices being connected. In order to capture *exogenous* effects (i.e., vertex attributes), we added *questionnaire* and *sub-scale* as predictors, especially testing for *homophily* (assuming that items belonging to the same questionnaire and its scale will have higher probability of being connected by an edge).

2.5 Association Network Inference

In order to understand the nature of the edges in the combined association network, we conceptualized the combined network as an *association network*, associations given by the correlation coefficients between responses to items. Our aim was to discern real edges from non-edges (when two items are not correlated) and derive the mixture null distribution in a data-driven fashion. The R package **fdrtool** (Strimmer, 2008a; Strimmer, 2008b) allowed us to model the probability density function and estimate η_0 (the fraction of true null hypotheses); and $(1-\eta_0)$ for the true alternative hypotheses. The mixture model for some statistic S thus takes the following form (where in our analysis, S is the partial correlation coefficient).

$$f(S) = \eta_0 f_0(S) + (1 - \eta_0) f_1(S)$$

To apply adjustment due to the number of performed tests, we controlled p-value by the *False Discovery Rate* (*FDR*; the rate of false rejections against the number of rejections).

2.6. Latent Network Modeling

We constructed an *eigenmodel* to further explore the structure of the combined graph of *RRS* and *SCRS* (Hoff et al., 2002; Kolaczyk, 2020; methodology following Mair, 2018) and how well the network structure predicts edges. Our expectation was that sub-scales will have non-trivial impact on the resulting eigenvalues. In our two-part analysis, first we used eigenvalue decomposition in order to use the first two eigenvectors reflective to our scale- and sub-scale structures of *RRS* and *SCRS*. In the second part, we defined three conditions of our *eigenmodel*. In the first condition we used the original *eigenmodel* (discussed above) without any additional regression components, defining the adjacency matrix only as a function of the pairwise latent effects summarized in the following matrix.

$$Y_{ij} = f(u_i \Lambda u_j)$$

In the second and third conditions, we defined optional regression components in addition to the pairwise latent effects, where $\mathbf{x}_{ij}$ is a pair-specific predictor vector. This extends our model to the following equation.

$$Y_{ij} = f(\beta' x_{ij} + u_i \Lambda u_j)$$

In the second condition, we cross-validated an extended *eigenmodel* with **Questionnaire** (*RRS* or *SCRS*) serving as grouping factor of vertices (the model captured the pairwise connections and Questionnaire playing an additional role in the edge formation process). In the same manner, the third condition tested a model with the **Scales** of *brooding*, *reflection* and *self-critical* serving as pair-wise covariate predictor effects.

2.7. Power analysis, network accuracy and stability

Accuracy and stability have an important role in estimating network parameters when samples might vary in characteristics and size (Epskamp, Borsboom, et al., 2017a, 2017b). One way of estimating network accuracy is to produce a 95% confidence interval of edge-weights with some variant of bootstrap. In the present paper, we chose the non-parametric bootstrap estimate for interpretability. Network stability is quantified with the *Correlation Stability* (*CS*-coefficient): an estimate of the maximum proportion of cases that can be dropped (with 95% certainty) to retain a correlation with the original centrality higher than some fixed threshold value (in the present paper, 0.7). Simulation studies (Epskamp, Borsboom, et al., 2017a) lead us in our expectation as the *CS*-coefficient should be between above 0.25, preferably higher than 0.5.

3. Results

3.1 Descriptives, network descriptives and community detection

Pearson correlation coefficients and Cronbach’s α s of the scales, means and standard deviations (for the total sample, and for females and males separately) are presented in Table 1. below. The data being skewed and homoscedasticity being hurt between gender groups, we estimated 20% trimmed Cohen’s d coefficients. With medium-to-high effect sizes, Cliff’s delta suggests group differences for female participants having higher scores on **RRS**, **SCRS** and RRS sub-scales (with at least 0.577 or higher probability). As expected (Smart et al., 2015), significant positive correlations were found between SCRS and RRS total scores (and both sub-scales), indicating that self-critical rumination somewhat differs from depressive mood related rumination, but people who are likely to ruminative self-critically are also likely to ruminate about their mood.

Table 1: Sample and Scale Descriptives

	1.	2.	3.	4.
Correlations:				
1. RRS				
2. RRS - brooding	0.85			
3. RRS - reflection	0.82	0.40		
4. SCRS	0.64	0.44	0.63	
Female Mean (SD)	10.71 (3.08)	10.80 (2.93)	25.19 (7.15)	10.88 (2.77)
Male Mean (SD)	9.42 (2.80)	9.53 (2.76)	21.23 (6.71)	9.65 (2.72)
Cronbach’s α	0.75	0.65	0.70	0.89
Cohen’s d (20% trimmed)	0.61	0.47	0.54	0.53
Cliff’s delta	P(f>m)=0.64	P(f>m)=0.58	P(f>m)=0.59	P(f>m)=0.62

Note. *RRS= Ruminative Response Scale. SCRS= Self-Critical Rumination Scale. SD=standard deviation. P(f>m)=probability of difference of female scores over male. All correlations are significant at $p<.001$.*

The undirected, weighted, associational networks of *RRS*, *SCRS* and their combined structure can be seen below (see Figure 1). All three networks proved to be connected (any vertex is reachable from every other). Comparing diameters to mean distances, we see densely connected graphs (for the *RRS* network, its diameter is 0.467 compared to its mean distance of 0.207; the *SCRS* and the combined networks show similar ratio patterns). On the plot, vertices represent the corresponding items of each questionnaire with labels showing item number, sub-scale and a label for each item. Sub-scales are presented in a colour-coded fashion as the outer circle of each vertex. Edges are the **LASSO** *LI*-penalized partial correlation coefficients whose value is represented as shade and width of the arcs.

The *RRS* (order of 10, size of 17) and *SCRS* (order of 10, size of 16) networks are densely connected networks. Analyzing their structure, while *SCRS* exhibits no *articulation point* [cut vertex whose removal would disconnect the network; Kolaczyk (2020)]. The vertex of the ‘problem’ (“What am I doing to deserve this?”) item of *RRS* acts as an **articulation point**, as it would cut off the ‘deserve’ (“Why do I have problems other people don’t have?”) vertex from the rest of the network. Articulation points are marked with * on Figure 1. Besides **size** and **order**, on Figure 1. we also indicated the **Clustering coefficient** (*transitivity*), the **diameter** and the average vertex-to-vertex path lengths (**mean distance**).

The combined network (order: 20, size: 39) has no articulation points. The previously identified articulation point of the *RRS* network (*problem* vertex) is no longer a cut-off vertex when all rumination data is encompassed in one graph.

In the *RRS* network two Reflection items proved to be have highly positive influence on their neighbourhoods, these are the ‘Go away by yourself and think about why you feel this way’- [EI1 (Expected Influence) = 0.91; EI2 (Two-Step Expected Influence) = 1.17; Br.EI1 (Bridge Expected Influence) = 0.22; Br.EI2 (Two-Step Bridge Expected Influence) = 0.24; TPR (True Positive Rate) = 1.00] and the ‘Analyze your personality to try to understand why you are depressed’- [EI1 = 1.11; EI2 = 0.95; Br.EI1 = 1.94; Br.EI2 = 1.77; TPR =


0.22] items. The item ‘Think ‘Why can’t I handle things better?’- [EI1 = 0.75; EI2 = 0.68; Br.EI1 = 1.18; Br.EI2 = 1.19; TPR = 0.96] was the most influential item of the *RRS* network. The items with the most negative, prohibiting influence of the *RRS* graph were ‘Think ‘What am I doing to deserve this?’- [EI1 = -1.57; EI2 = -1.60; Br.EI1 = -1.15; Br.EI2 = -1.37; TPR = 1.00]; ‘Analyze recent events to try to understand why you are depressed’- [EI1 = -0.82; EI2 = -0.70; Br.EI1 = -1.00; Br.EI2 = -0.69; TPR = 0.22] and ‘Write down what you are thinking and analyze it’- [EI1 = -1.29; EI2 = -1.25; Br.EI1 = -0.76; Br.EI2 = -1.02; TPR = 0.99]. These items stand out in their characteristic that their two-step influence metrics (influence on the rest of the graph also taken into account) are negative.

In the *SCRS* graph highly positive influence were contributed to items such as ‘I spend a lot of time thinking about how ashamed I am of some of my personal habits.’- [EI1 = 2.12; EI2 = 2.09; Br.EI1 = 2.32; Br.EI2 = 2.07; TPR = 1.00] and ‘I wish I spent less time criticizing myself.’- [EI1 = 1.51; EI2 = 1.21; Br.EI1 = 0.42; Br.EI2 = 0.22; TPR = 1.00]. The highest negative influence items of *SCRS* were ‘I can’t stop thinking about how I should have acted differently in certain situations.’- [EI1 = -0.77; EI2 = -0.73; Br.EI1 = -0.96; Br.EI2 = -1.14; TPR = 1.00] and ‘I spend a lot of time wishing I were different.’- [EI1 = -0.75; EI2 = -0.82; Br.EI1 = -0.63; Br.EI2 = -0.60; TPR = 1.00].


In the combined *RRS* - *SCRS* graph, the item of ‘Go away by yourself and think about why you feel this way’- [EI1 = 0.88; EI2 = 0.71; Br.EI1 = -0.51; Br.EI2 = -0.57; TPR = 1.00] not only proved to be highly positively influential, but also increased its Betweenness centrality, meaning it started to act as a connecting point between sub-domains of the network. The rest of the most positively influential nodes of this network are from *SCRS* such as ‘I can’t stop thinking about how I should have acted differently in certain situations.’- [EI1 = 1.61; EI2 = 1.47; Br.EI1 = 0.18; Br.EI2 = 0.12; TPR = 0.92], ‘I spend a lot of time thinking about how ashamed I am of some of my personal habits.’- [EI1 = 1.47; EI2 = 1.70; Br.EI1 = 0.80; Br.EI2 = 0.89; TPR = 0.92], ‘I criticize myself a lot for how I act around other people.’- [EI1 = 1.06; EI2 = 1.15; Br.EI1 = 0.26; Br.EI2 = 0.39; TPR = 0.92] and ‘I often worry about all of the mistakes I have made.’- [EI1 = 1.32; EI2 = 1.16; Br.EI1 = 3.19; Br.EI2 = 3.02; TPR = 0.92]. With regard to the negative influence of network activation in the combined graph, the pattern is dominated by *RRS* items such as ‘Think ‘What am I doing to deserve this?’- [EI1 = -1.82; EI2 = -1.80; Br.EI1 = -1.00; Br.EI2 = -0.91; TPR = 0.96], ‘Think ‘Why do I always react this way?’- [EI1 = -1.23; EI2 = -1.33; Br.EI1 = 0.47; Br.EI2 = 0.23; TPR = 0.90], ‘Write down what you are thinking and analyze it’- [EI1 = -1.12; EI2 = -1.11; Br.EI1 = -1.00; Br.EI2 = -1.16; TPR = 1.00], ‘Think about a recent situation, wishing it had gone better’- [EI1 = -0.81; EI2 = -0.78; Br.EI1 = 0.18; Br.EI2 = 0.37; TPR = 0.96] and ‘Think ‘Why can’t I handle things better?’- [EI1 = -1.08; EI2 = -1.23; Br.EI1 = -0.31; Br.EI2 = -0.37; TPR = 0.95]. Note the item of **write** increased its Betweenness compared to the standalone *RRS* graph. Among the *SCRS* items, the ‘I wish I spent less time criticizing myself.’- [EI1 = -0.94; EI2 = -0.65; Br.EI1 = -0.27; Br.EI2 = -0.43; TPR = 0.92] was the most negatively influential on the common network.

In order to analyze their respective internal structures, we performed community detection using the *igraph* package in R. On Figure 1., the inner node point of each vertex provides colour coding for our community detection results (the presented shades correspond to the True Positive Rates of each sub-scale; in other words, the average match between the detected community and the original sub-scale [ranging between 0 and 1]).


RRS undirected graph
Order=10; Size=17; Clustering Coefficient=0.444; Diameter=0.467; Mean distance=0.207



SCRS undirected graph
Order=10; Size=16; Clustering Coefficient=0.125; Diameter=0.868; Mean distance=0.461



RRS & SCRS undirected graph
Order=20; Size=39; Clustering Coefficient=0.206; Diameter=0.989; Mean distance=0.391



a., TPR is average True Positive Rate with 1000 times reruns of community detection
b., * marks Articulation Points (cut vertices; when such vertices are removed disconnect the graph)
c., Order is # of vertices; Size is # of edges
d., Edge width reflect edge weight (penalized part. corr.)
e., Dashed edge line indicates the 95% CI of accuracy estimate ranges below 0.0

Figure 1: Tableau view of three graphs of rumination

As the choice of algorithm has non-trivial impact on the identified community structure, after having selected the “*optimal*”, the “*spinglass*”, the “*walktrap*” and (to also include hierarchical clustering) the “*fast greedy*” methods, we tested each 1000 times on the three networks. We also provided an edge weight vector for community detection in the form of the *LASSO*-penalized partial correlation coefficients (or alternatively, omitted weights). Presented TPR values are therefore averages of estimates from all these conditions.

The **RRS** two-scale structure could be recreated with all four algorithms, the inclusion of the weight vector not having effect on sub-scale membership. For the sub-scale of *brooding* the average **true positive rate** (TPR) was 0.8; for *reflection* 0.743. In case of the *reflection* sub-scale, lower average TPR becomes tangible with the inspection of items 2 (“*analyze*”; “*Analyze recent events to try to understand why you are depressed*”) and 9 (“*person*”; “*Analyze your personality to try to understand why you are depressed*”) whose low TPR indicates that depending in the choice of community detection they have a chance of forming their own community or being classified as members of the *brooding* scale, rather than being part of *reflection*. The one-scale structure of the *SCRS* has been reestablished.

The combined rumination network shows a clear expected structure, with having high TPR for each sub-scale (on average, *brooding* (0.9), *reflection* (1) and *self-critical* (0.8)). Commenting on the stability of the network community structure, the *RRS reflection* sub-scale exhibits strong TPR rates, and the weakness of the two items from their respective stand-alone network for items 2 and 9 are no longer present in the combined graph. The *RRS brooding* scale has good separation in terms of TPR, except for its item 3 (*react*) having <0.9 TPR (“*Why do I always react this way?*”). The one-scale *SCRS* has also a separable community with high (>0.925) TPR. The combined network has large TPR of 0.9; while keeping only the *spinglass* algorithm (as used by Bernstein et al., 2019), we get high average 1.0 TPR for the combined graph.

Please refer to our vertex summary table in the Appendix (Table 4.) for actual descriptives, including expected influence, bridge node related statistics and average True Positive Rates for each vertex. Where applicable, we used the standardized scores.

3.2 Internal Structure with an exponential random graph model

Our ERGM model was built with *Markov chain Monte Carlo* (MCMC) maximum likelihood estimation (1000 iterations) to obtain model parameters by generating samples from the space of possible networks allowed by the data.

Table 2: ERGM model fit

Term	Estimate	Std. Error	MCMC Error %	Statistic	p	p of EB
# of edges	-2.24	0.61	0	-3.70	0.00	0.10
GWESP [triangle]	-0.37	0.16	0	-2.32	0.02	0.41
Questionnaire (main effect)	-0.04	0.41	0	-0.10	0.92	0.49
Questionnaire (homophily effect)	-0.82	1.24	0	-0.67	0.51	0.30
Scale (homophily effect)	4.18	1.34	0	3.13	0.00	0.99

Note. MCMC=Monte Carlo Markov Chain; GWESP=Geometrically Weighted Edgewise Shared Partner Distribution; p of EB=probability of Edge Building; model built with Markov chain Monte Carlo Maximum Likelihood Estimation

An ANOVA analysis on the overall model indicates (with df=5, residual deviance=143.38, $p < 0.001$) that variables involved in the model explain the network connectivity in a highly non-trivial fashion. The detailed analysis reveals that *endogenous* effects such as the overall density of the network (number of *edges*) change the probability of the formation of an edge by 0.096), and at $p < 0.05$ level the closed triangle formulation indicates a non-trivial impact on clustering (connectivity probability changed by 0.408). As expected, *exogenous* effects, such as two items belonging to the same sub-scale have a significant positive impact on connectivity (sub-scale changes edge probability by 0.985, keeping the rest of the network fixed). Questionnaires themselves have no impact on edge formation (‘Questionnaire main effect’). In terms of goodness-of-fit, observed

frequencies of degree levels differ in non-significant manner from *MCMC*-simulated expectations. In two cases of vertex degree ($V_d=3$ and $V_d=6$), the observed frequencies are above the simulated average ($6 >$ the expected avg. of 4.49 for $V_d=3$; and $4 >$ the expected avg. of 1.94 for $V_d=6$), but differences are not significant.

Regarding the expected edgewise shared partners, the only deviation from expected levels is for 1 partner, where the observed frequency is 17 against 13.28. Additional MCMC diagnostics revealed good fit of the model, as the observed graph does not differ from the *MCMC* simulated graphs.

3.3 Association Network Inference

Results show low η_0 (0.051) indicating large mixing rate for the alternative hypothesis of edges having non-zero weights. The histogram of absolute correlation coefficients (see Figure 3. in Appendix) has a centered shape in contrast to the null component. The density-based *FDR* drops once empirical correlations reach at least $\rho=0.1$ (see the lowest section of Figure 3.).

3.4 Latent Network Modeling

Our next model was built with the *eigenmodel_mcmc* function of the R package **eigenmodel**. The function uses *Markov chain Monte Carlo* (MCMC) techniques to complete a model specification. By applying eigenvalue decomposition, we obtained the first two eigenvectors (see Figure 2) on the combined network ($S=10000$, number of iterations).

Our visual insight from Figure 2. is that the first eigenvector serves as a separator between *RRS* and *SCRS* having a posterior mean of $e\bar{v}_1=1.344$. The second eigenvector seems to separate between the upper half of the two-dimensional space of *SCRS* and the *brooding* scale, and the lower half being the *reflection* scale ($e\bar{v}_2=2.332$).

In order to test the goodness-of-fit of the eigenmodel, using k -fold cross-validation ($k=5$), we refitted the model to a subset of the original data, using the complement 1/5 fraction for predicting the elements of the adjacency matrix of our original network. We used three different models in this part as discussed above (Section 2.6).

Summarizing the results from the *eigenmodels*, using the original combined network's adjacency matrix as a reference, we estimated the *Area Under the Curve* (AUC) statistics for each model. Compared to the latent model ($AUC_{latent}=0.348$), the inclusion of the questionnaires as a grouping factor resulted in a better predictive model ($AUC_{latent+questionnaire}=0.637$). The final model involving the scale-level vertex characteristics has the best goodness-of-fit ($AUC_{latent+scale}=0.725$). See Figure 4. in the Appendix for the corresponding ROC curves.

3.5 Power analysis, network accuracy and stability

We used the *bootnet* R package to have an estimate of edge-weight accuracy with *non-parametric bootstrap* (5000 times sampling). The resulting **accuracy** estimates have been visually incorporated in Figure 1 in the following manner. Edge shade reflects the lower bound of the 95% confidence interval (CI) for each edge, while edges are presented with dashed line for cases of the CI including 0.0 edge weights. For such cases, we cannot be sure if their weight persist given different samples and varying sample characteristics. Darker shaded, continuous edges represent stable edges according to the bootstrap estimate. Edges such as between the *SCRS(5)* and *SCRS(7)* items indicate moderate penalized partial correlation between items in the *SCRS* network ($\rho=0.352$), but the edge is presented with dashed line and has lighter shade indicating a lower stability. Examples for accurate and strong edges are *RRS(4)* (*alone1*; “Go away by yourself and think about why you feel this way”) and *RRS(10)* (*alone2*; “Go someplace alone to think about your feelings”). The edge weight for these items in their stand-alone *RRS* network is 0.572 (estimated edge-weight accuracy is with average 0.511 with lower and upper CI bounds of 0.454 - 0.567). These items exhibit the same pattern in the combined network with estimated edge-weight accuracy with average weight of 0.493; lower and upper CI bounds of 0.436 - 0.549.




Figure 2: Vertices of the combined graph by the first two eigenvalues

In the SCRS network we can track a *path* (a graph movement [walk] without repeated edges or vertices) between SCRS 6, 5, 9 and 10 items (please refer to Table 5. for edge-level weight and accuracy estimates of SCRS).

Network **stability** estimates for our three networks were carried out by using the *bootnet* function (results in Table 3. below). We performed a *case dropping bootstrap* to simulate scenarios where fewer and fewer cases of the original sample would have been available and quantified the stability of the centrality metrics with the *Correlation Stability* (CS) coefficient. Bootstrap was performed with B=5000 samples.

Accuracy and stability statistics have been summarized for RRS, SCRS and their combined graphs in respective tables of Appendix (see Table 5., 6., and 7.).

As shown at (Epskamp, Borsboom, et al., 2017a), the *Correlation Stability* coefficient is not acceptable below 0.25, and preferably stays above 0.5.

Table 3: Correlation Stability coefficient estimates

Metric	RRS	RRS:Brooding	RRS:Reflection	SCRS	RRS & SCRS
Betweenness	0.01	0.26	0.10	0.07	0.04
Closeness	0.17	0.39	0.33	0.37	0.01
Edge Strength	0.28	0.56	0.75	0.28	0.52
Expected Influence	0.36	0.56	0.82	0.27	0.52
Edge Weight	0.70	0.58	0.85	0.72	0.66

We found satisfactory correlation stability of the Betweenness statistic of the Brooding (RRS) stand-alone network ($CS_{Brooding}=0$) as Brooding has retraceable communities, and as we already pointed out above, had the only Articulation Point of the RRS network. Brooding, Reflection and the SCRS networks have acceptable correlation statistics for the closeness statistic ($CS_{Brooding}=0$, $CS_{Reflection}=0$ and $CS_{SCRS}=0$). The Edge Strength (the sum of edge weight, in the present paper the partial correlation coefficients) and the Expected Influence statistics have good correlation stabilities on all networks; specifically, the Reflection stand-alone network exhibits high estimates for Edge Strength ($CS_{Reflection}=1$) and Expected Influence ($CS_{Reflection}=1$) which might be driven by the high correlation between items 4. and 10. of RRS. The Edge Weight shows highly satisfactory patterns on all networks (“*Edge Weight*” row of Table 3.) which is the expected stability of our penalized partial correlation coefficients. For the combined network of RRS and SCRS, we see stable Edge Strength, Edge Weight and Expected Influence stability estimates, whereas Betweenness ($CS_{combined}=0$) and Closeness ($CS_{combined}=0$) are unsatisfactory. These findings suggest that the combined network tolerates very few sample loss and sample variance when it comes to the Betweenness centrality and the Closeness centrality measures, as all networks are densely connected and fewer items can stand out with higher local importance such as being between sub-networks.

4. Discussion

In the present paper, we examined the separation of two facets of rumination (a depressive dimension, lying along the *brooding-reflection* axis; and the *self-critical* aspects of rumination) using a network approach. The **Self-Critical Rumination Scale** proved to have a one-scale structure. In its respective network, the **Ruminative Response Style Questionnaire** exhibits a two-scale structure. Our community detection estimates suggest though that the reflective aspects of rumination such as analyzing one’s self and recent events for better emotional insights (items 2 and 9 of *RRS*) have more in common with brooding, or have a tendency to emerge as a third aspect. In analyzing the factor structure of *RRS* on samples from Argentina and the United States, the error variances of these two items were found to be correlated and thus ended up being excluded from the solution (Arana & Rice, 2017). The association of these items with depression, and therefore their elimination from any analysis related to Reflection have been suggested and practiced (Armey et al., 2009; Lee & Kim, 2014; Treynor, 2003). With concurring results on these items, the hypothesis can be phrased that “*the distinction between reflection and brooding is blurred in depressed individuals*,” meaning the confounding effect of these items with depression (Whitmer & Gotlib, 2011). The same authors also suggest that “*given that these three items measure a tendency to self-isolate with the intention to ruminate, we call this factor ‘intentional rumination’*.” On a sample of Dutch students, researchers excluded these two items for reasons of their “*reference to negative feelings*,” and the resulting three-item Reflection scale was named as “*reflective self-regulation*” (Brose et al., 2020). Although, in a recent study on samples of Peruvian students, the separation of the “analyze” items was not present (Valencia & Paredes-Angeles, 2022). A clear two-factor structure of *RRS* was revealed on Columbian samples (Ruiz, 2017) or on Japanese sample (Hasegawa, 2013). These findings on two- or three-factor structures of *RRS* support the notion of our findings that the items of ‘intentional rumination’ are sensitive to sampling characteristics, demographics, and as we have seen in our combined network, to covariates or additional activation paths taken into account. Items #2 and #9 of *RRS* on their own graph emerged as a third component, presented a low *true positive rate* of 0.22 each, but this pattern dissolved in a combined graph with the self-critical aspects of rumination accounted for.

Separation between scales in the **combined graph** is underlined by our *eigenmodel*, as in the two-dimensional space of the first two eigenvectors, expected grouping of *RRS* and *SCRS* items can be achieved. Items #2 and #9 of *RRS* are well separated from the bulk of the rest of the *RRS* items, supporting our above conclusion of them being a part of reflection or found a new component. Brooding has more associations with the self-critical aspects of rumination than reflection, which is highlighted by our eigenvector analysis as well as the sparse, unstable and weak edges with which reflection connects to brooding and self-critical rumination in their combined network. Our graphs suggest and support the notion of not referring to a psychological attribute as a sum score, but rather take into account the underlying activation pattern (Bernstein et al., 2019). A similar approach was used by (Williams et al., 2021), where researchers combined items of “*reflecting pondering*” into one node.

We have gathered evidence of scale playing an important role in the edge-building process. According to our exponential random graph model, the internal scales of the combined network indeed have a non-trivial impact on associations between items, whereas the effect of questionnaire dissolved, no longer having an impact on the probability of edge formation. This suggests that all three scales are effective in assessing a separate facet of rumination when used together. As shown, associations within the combined network only have low chances of false discovery.

Results indicate that network associations are accurately aligned to the internal factor structure, while some associations are more sensitive to sampling characteristics and sampling variation. Edge strengths and edge weights have also high stability and relatively high resistance to sample loss, whereas betweenness is remarkably sensitive to case-dropping.

Our findings suggest that the brooding and reflective aspects of depressive rumination and the self-critical rumination are three distinct, unique areas of the multifaceted construct of rumination. A network approach identified them as not merging with one-another, but establishing a stable and sensitive network of items.

Future steps include analyzing such data from a person-level graph perspective, where vertices are individuals, and edges can be some distance measure between vertices based upon demographics and survey data. Also, if keeping our analysis on the item-level, as in the present paper, we would extend our methodology to directed

acyclic graphs (DAGs) which would allow causal assumptions to support our understanding various facets of rumination.

Appendix

Table 4: Network descriptive statistics for RRS, SCRS and their combined networks

Graph	Item	Scale	Label	Deg.	Str.	Bet.	Clo.	EI1	EI2	Br.Str.	Br.Bet.	Br.Cl.	Br.EI1	Br.EI2	TPR
RRS	RRS_1	brooding	deserve	1	0.14	0	0.33	-1.57	-1.60	-1.15	-0.89	-1.60	-1.15	-1.37	1.00
	RRS_2	reflection	analyze	3	0.34	0	0.78	-0.82	-0.70	-1.00	-0.89	-0.09	-1.00	-0.69	0.22
	RRS_3	brooding	react	3	0.65	0	0.41	0.28	0.40	0.46	-0.08	0.78	0.46	0.72	0.96
	RRS_4	reflection	alone1	4	0.82	0	0.72	0.91	1.17	0.22	0.32	0.71	0.22	0.24	1.00
	RRS_5	reflection	write	2	0.21	0	0.57	-1.29	-1.25	-0.76	-0.89	-1.43	-0.76	-1.02	0.99
	RRS_6	brooding	wish	2	0.38	0	0.51	-0.69	-0.70	-0.63	-0.89	-0.77	-0.63	-0.65	1.00
	RRS_7	brooding	problem	4	0.67	8	0.51	0.36	0.04	0.22	1.33	0.12	0.22	0.06	1.00
	RRS_8	brooding	handle	7	0.78	26	0.81	0.75	0.68	1.18	0.32	0.47	1.18	1.19	0.96
	RRS_9	reflection	person	5	0.88	16	0.68	1.11	0.95	1.94	1.94	1.55	1.94	1.77	0.22
	RRS_10	reflection	alone2	3	0.83	0	0.47	0.95	1.01	-0.49	-0.28	0.25	-0.49	-0.25	1.00
SCRS	SCRS_1	self-critical	aspects	2	0.77	0	0.19	-0.31	0.29	-0.16	0.53	0.84	-0.16	0.71	1.00
	SCRS_2	self-critical	said	3	0.59	3	0.30	-0.70	-0.75	-0.31	-0.74	-0.88	-0.31	-0.46	1.00
	SCRS_3	self-critical	thoughts	4	0.94	6	0.29	0.04	-0.27	0.51	-0.32	-0.44	0.51	0.06	1.00
	SCRS_4	self-critical	situation	2	0.56	0	0.27	-0.77	-0.73	-0.96	-0.74	-1.17	-0.96	-1.14	1.00
	SCRS_5	self-critical	habits	7	1.92	15	0.34	2.12	2.09	2.32	2.21	1.63	2.32	2.07	1.00
	SCRS_6	self-critical	act around	2	0.74	0	0.19	-0.37	0.16	-0.11	-0.74	0.96	-0.11	0.17	1.00
	SCRS_8	self-critical	mistakes	5	1.63	6	0.27	1.51	1.21	0.42	0.95	0.52	0.42	0.22	1.00
	SCRS_9	self-critical	different	3	0.80	3	0.25	-0.25	-0.11	0.22	0.32	0.37	0.22	0.42	1.00
	SCRS_7	self-critical	time	2	0.57	0	0.21	-0.75	-0.82	-0.63	-0.74	-0.50	-0.63	-0.60	1.00
	SCRS_10	self-critical	productive	2	0.67	0	0.19	-0.52	-1.09	-1.30	-0.74	-1.32	-1.30	-1.44	1.00
Combined	RRS_1	brooding	deserve	2	0.31	0	0.11	-1.82	-1.80	-1.00	-1.05	-0.91	-1.00	-0.91	0.96
	RRS_2	reflection	analyze	3	0.57	2	0.13	-0.74	-0.79	-0.17	-0.35	-0.86	-0.17	-0.48	1.00
	RRS_3	brooding	react	3	0.45	15	0.15	-1.23	-1.33	0.47	-0.08	0.23	0.47	0.23	0.90
	RRS_4	reflection	alone1	4	0.94	23	0.17	0.88	0.71	-0.51	0.45	-1.15	-0.51	-0.57	1.00
	RRS_5	reflection	write	2	0.48	11	0.09	-1.12	-1.11	-1.00	-0.88	-1.45	-1.00	-1.16	1.00
	RRS_6	brooding	wish	3	0.55	3	0.12	-0.81	-0.78	0.18	-0.52	0.55	0.18	0.37	0.96
	RRS_7	brooding	problem	5	0.80	26	0.16	0.28	0.06	1.50	0.81	1.11	1.50	1.49	0.96
	RRS_8	brooding	handle	3	0.49	7	0.14	-1.08	-1.23	-0.31	-0.35	-0.20	-0.31	-0.37	0.95
	RRS_9	reflection	person	4	0.75	31	0.16	0.05	-0.21	-0.38	1.07	-0.66	-0.38	-0.41	1.00
	SCRS_1	self-critical	aspects	4	0.69	5	0.14	-0.19	-0.01	-1.00	-1.05	-0.83	-1.00	-0.94	0.92
	SCRS_2	self-critical	said	4	0.86	0	0.13	0.51	0.53	-1.00	-0.70	-0.58	-1.00	-0.90	0.92
	SCRS_3	self-critical	thoughts	5	0.94	6	0.14	0.87	0.90	-0.01	-0.52	0.34	-0.01	0.07	0.92
	SCRS_4	self-critical	situation	6	1.11	28	0.18	1.61	1.47	0.18	0.19	0.31	0.18	0.12	0.92

Graph	Item	Scale	Label	Deg.	Str.	Bet.	Clo.	EI1	EI2	Br.Str.	Br.Bet.	Br.Cl.	Br.EI1	Br.EI2	TPR
	SCRS_5	self-critical	habits	6	1.08	12	0.17	1.47	1.70	0.80	1.34	1.26	0.80	0.89	0.92
	SCRS_6	self-critical	act around	6	0.99	51	0.18	1.06	1.15	0.26	2.22	0.83	0.26	0.39	0.92
	SCRS_7	self-critical	time	4	0.76	9	0.14	0.11	0.34	0.10	-0.52	0.72	0.10	0.07	0.92
	SCRS_8	self-critical	mistakes	3	0.52	2	0.14	-0.94	-0.65	-0.27	-0.70	-0.37	-0.27	-0.43	0.92
	SCRS_9	self-critical	different	6	1.05	38	0.18	1.32	1.16	3.19	2.04	2.18	3.19	3.02	0.92
	RRS_10	reflection	alone2	2	0.76	0	0.07	0.11	0.13	-1.00	-0.88	-1.48	-1.00	-1.10	1.00
	SCRS_10	self-critical	productive	3	0.66	1	0.12	-0.35	-0.23	-0.01	-0.52	0.93	-0.01	0.63	0.92

Deg.=Degree; Str.=Strength; Bet.=Betweenness; Clo.=Closeness; EI1= One-step Expected Influence; EI2= Two-step Expected Influence; Br.Str.=Bridge Strength; Br.Bet.=Bridge Betweenness; Br.Cl.=Bridge Closeness; Br.EI1=One-step Bridge Expected Influence; Br.EI2=Two-step Bridge Expected Influence; TPR=average True Positive Rate






Figure 3: False Discovery Rate (FDR) for the combined network

Table 5: Edge Accuracy and Stability Statistics for the RRS network

Vx.1	Vx.2	Avg. Acc.	L.B.Acc.	U.B.Acc.	Miss.:%:0	Miss.:%:0.2	Miss.:%:0.4	Miss.:%:0.6	Miss.:%:0.7
RRS_1	RRS_7	0.20	0.13	0.27	0.20	0.20	0.19	0.18	0.12
RRS_2	RRS_4	0.15	0.08	0.22	0.16	0.15	0.14	0.12	0.08
RRS_2	RRS_8	0.00	-0.01	0.01	0.00	0.00	0.00	0.00	0.00
RRS_2	RRS_9	0.25	0.19	0.32	0.25	0.25	0.26	0.24	0.22
RRS_3	RRS_7	0.06	-0.07	0.18	0.01	0.01	0.02	0.02	0.02
RRS_3	RRS_8	0.23	0.16	0.30	0.23	0.23	0.23	0.22	0.17
RRS_3	RRS_9	0.09	-0.04	0.22	0.10	0.07	0.05	0.04	0.02
RRS_4	RRS_10	0.51	0.45	0.57	0.51	0.51	0.51	0.51	0.50
RRS_4	RRS_8	0.06	-0.05	0.17	0.06	0.04	0.02	0.02	0.02
RRS_4	RRS_9	0.09	-0.03	0.20	0.10	0.08	0.05	0.05	0.03
RRS_5	RRS_10	0.11	0.00	0.22	0.12	0.11	0.07	0.06	0.03
RRS_5	RRS_8	0.00	-0.04	0.04	0.00	0.00	0.00	0.00	0.00
RRS_6	RRS_7	0.20	0.13	0.27	0.20	0.20	0.19	0.17	0.10
RRS_6	RRS_8	0.09	-0.04	0.22	0.11	0.07	0.05	0.03	0.02
RRS_7	RRS_9	0.00	-0.01	0.01	0.00	0.00	0.00	0.00	0.00
RRS_8	RRS_10	0.00	-0.03	0.04	0.00	0.00	0.00	0.00	0.00
RRS_8	RRS_9	0.09	-0.03	0.22	0.11	0.09	0.05	0.04	0.03

Vx.1=1st Vertex the edge attaches to; Vx.2=2nd Vertex the edge attaches to; Avg.Acc.=Average Edge Weight by Bootstrap; L.B.Acc.=Lower Bound of 95% CI for bootstrapped network accuracy; U.P.Acc.=Upper Bound of 95% CI for bootstrapped network accuracy; Miss.%=Estimated average stability at the given missing rate of sample size

Table 6: Edge Accuracy and Stability Statistics for the SCRS network

Vx.1	Vx.2	Avg. Acc.	L.B.Acc.	U.B.Acc.	Miss.:%:0	Miss.:%:0.2	Miss.:%:0.4	Miss.:%:0.6	Miss.:%:0.7
SCRS_1	SCRS_5	0.16	0.09	0.23	NA	NA	NA	NA	NA
SCRS_1	SCRS_8	0.04	-0.06	0.13	NA	NA	NA	NA	NA
SCRS_2	SCRS_3	0.03	-0.06	0.11	NA	NA	NA	NA	NA
SCRS_2	SCRS_5	0.00	-0.03	0.03	NA	NA	NA	NA	NA
SCRS_2	SCRS_8	0.18	0.11	0.25	NA	NA	NA	NA	NA
SCRS_3	SCRS_10	0.18	0.11	0.24	NA	NA	NA	NA	NA
SCRS_3	SCRS_5	0.11	0.03	0.19	NA	NA	NA	NA	NA
SCRS_3	SCRS_7	0.25	0.18	0.32	NA	NA	NA	NA	NA
SCRS_4	SCRS_5	0.11	0.02	0.20	NA	NA	NA	NA	NA
SCRS_4	SCRS_8	0.23	0.16	0.29	NA	NA	NA	NA	NA
SCRS_5	SCRS_6	0.29	0.23	0.36	NA	NA	NA	NA	NA
SCRS_5	SCRS_7	0.06	-0.05	0.16	NA	NA	NA	NA	NA
SCRS_5	SCRS_9	0.20	0.13	0.27	NA	NA	NA	NA	NA
SCRS_6	SCRS_8	0.09	-0.02	0.19	NA	NA	NA	NA	NA
SCRS_8	SCRS_9	0.12	0.03	0.20	NA	NA	NA	NA	NA
SCRS_9	SCRS_10	0.30	0.23	0.36	NA	NA	NA	NA	NA

Vx.1=1st Vertex the edge attaches to; Vx.2=2nd Vertex the edge attaches to; Avg.Acc.=Average Edge Weight by Bootstrap; L.B.Acc.=Lower Bound of 95% CI for bootstrapped network accuracy; U.P.Acc.=Upper Bound of 95% CI for bootstrapped network accuracy; Miss.%=Estimated average stability at the given missing rate of sample size

Table 7: Edge Accuracy and Stability Statistics for the RRS and SCRS network

Vx.1	Vx.2	Avg. Acc.	L.B.Acc.	U.B.Acc.	Miss.:%:0	Miss.:%:0.2	Miss.:%:0.3	Miss.:%:0.6	Miss.:%:0.7
RRS_1	RRS_6	0.06	-0.09	0.22	0.00	0.01	0.01	0.00	0.01
RRS_1	RRS_7	0.13	-0.03	0.29	0.16	0.11	0.07	0.03	0.01
RRS_2	RRS_3	0.06	-0.09	0.21	0.00	0.01	0.01	0.01	0.01
RRS_2	RRS_4	0.13	0.01	0.26	0.15	0.13	0.10	0.04	0.02
RRS_2	RRS_9	0.24	0.17	0.30	0.24	0.24	0.24	0.19	0.11
RRS_3	RRS_8	0.16	0.04	0.28	0.17	0.16	0.13	0.06	0.02
RRS_3	SCRS_6	0.03	-0.08	0.14	0.00	0.00	0.00	0.00	0.00
RRS_4	RRS_10	0.49	0.44	0.55	0.49	0.49	0.49	0.49	0.46
RRS_4	RRS_9	0.06	-0.08	0.19	0.01	0.02	0.01	0.02	0.01
RRS_4	SCRS_9	0.02	-0.07	0.10	0.00	0.00	0.00	0.00	0.00
RRS_5	RRS_10	0.06	-0.08	0.21	0.01	0.01	0.01	0.00	0.00
RRS_5	RRS_9	0.19	0.09	0.29	0.20	0.19	0.17	0.08	0.03
RRS_6	RRS_7	0.14	0.00	0.28	0.16	0.13	0.09	0.04	0.02
RRS_6	SCRS_4	0.17	0.08	0.26	0.17	0.17	0.15	0.09	0.05
RRS_7	RRS_8	0.06	-0.08	0.21	0.00	0.01	0.01	0.01	0.00
RRS_7	SCRS_5	0.07	-0.06	0.21	0.03	0.03	0.03	0.02	0.01
RRS_7	SCRS_9	0.12	-0.01	0.25	0.14	0.11	0.07	0.05	0.04
RRS_8	SCRS_10	0.12	-0.01	0.25	0.15	0.12	0.09	0.04	0.03
RRS_9	SCRS_6	0.02	-0.07	0.12	0.00	0.00	0.00	0.00	0.00
SCRS_1	SCRS_2	0.24	0.17	0.30	0.24	0.24	0.24	0.23	0.18
SCRS_1	SCRS_3	0.14	0.05	0.23	0.15	0.15	0.13	0.09	0.05
SCRS_1	SCRS_5	0.15	0.06	0.24	0.16	0.15	0.14	0.11	0.06
SCRS_1	SCRS_7	0.05	-0.07	0.17	0.01	0.01	0.02	0.01	0.01
SCRS_2	SCRS_4	0.24	0.18	0.31	0.24	0.24	0.24	0.24	0.19
SCRS_2	SCRS_6	0.07	-0.05	0.19	0.07	0.04	0.03	0.01	0.02
SCRS_2	SCRS_8	0.16	0.08	0.24	0.16	0.16	0.15	0.11	0.07
SCRS_3	SCRS_10	0.15	0.05	0.24	0.16	0.16	0.14	0.10	0.06
SCRS_3	SCRS_4	0.17	0.10	0.25	0.17	0.17	0.17	0.14	0.09
SCRS_3	SCRS_5	0.08	-0.04	0.20	0.09	0.06	0.05	0.03	0.02
SCRS_3	SCRS_7	0.23	0.17	0.30	0.24	0.24	0.23	0.23	0.16
SCRS_4	SCRS_5	0.11	0.00	0.22	0.12	0.11	0.09	0.06	0.04
SCRS_4	SCRS_6	0.03	-0.07	0.14	0.00	0.01	0.01	0.01	0.01
SCRS_4	SCRS_8	0.20	0.13	0.27	0.20	0.20	0.20	0.18	0.13
SCRS_5	SCRS_6	0.29	0.22	0.36	0.29	0.29	0.29	0.28	0.25
SCRS_5	SCRS_9	0.17	0.10	0.25	0.17	0.17	0.17	0.15	0.10

Vx.1	Vx.2	Avg. Acc.	L.B.Acc.	U.B.Acc.	Miss.:%:0	Miss.:%:0.2	Miss.:%:0.3	Miss.:%:0.6	Miss.:%:0.7
SCRS_6	SCRS_7	0.17	0.09	0.25	0.17	0.17	0.17	0.12	0.08
SCRS_7	SCRS_9	0.14	0.03	0.24	0.14	0.14	0.12	0.09	0.06
SCRS_8	SCRS_9	0.09	-0.03	0.21	0.12	0.09	0.06	0.03	0.02
SCRS_9	SCRS_10	0.26	0.20	0.33	0.27	0.27	0.27	0.27	0.21

Vx.1=1st Vertex the edge attaches to; Vx.2=2nd Vertex the edge attaches to; Avg.Acc.=Average Edge Weight by Bootstrap; L.B.Acc.=Lower Bound of 95% CI for bootstrapped network accuracy; U.P.Acc.=Upper Bound of 95% CI for bootstrapped network accuracy; Statistic=type of network characteristic; Miss.%=Estimated average stability at the given missing rate of sample size




Figure 4: ROC curves of the combined RRS-SCRS network

Table 8: Glossary of network analysis relevant terms mentioned in the article

Term	Definition
Articulation point	A measure of vertex influence: $V(i)$ vertex of G graph is an Articulation Point, if removed from the graph, would segment, cut or disintegrate the graph; therefore is considered locally important.
Association network	A network where neighbouring vertices when connected are assumed to being associated; in other words, edges represent some measure of correlation or other kind of association; this makes them different from energy or information flow networks, for instance.
Community Detection	A group of algorithmic methods to find clusters of nodes whose community can be described by systematic variance encoded in data represented in edges and node characteristics, and therefore are non-trivial. In the current paper, we used four selected algorithms to estimate how well they identify the original subscale-structure of the RRS and SCRS questionnaires.
Cut vertex	see ‘Articulation point’
Edge	A link (E , arc) between nodes of a network. Undirected if direction of the connection (e.g. flow of energy or information) is not essential to the network; directed (uni- or bidirectional) otherwise. Can be weighted by some measure of association, frequency or other attribute.
Exponential Random Graph Models	ERGM; a family of models to the analogy of generalized linear models adopted to network modeling. They allow the construction, fitting and comparison of network models. Their nature is structural, incorporating different configurations of a graph (sets of possible edges). ERGMs have numerous extensions and variations.
Graph	see ‘Network’

Term	Definition
Markov Chain Monte Carlo	Monte Carlo methods are a collection of techniques which approximate a quantity (mean, variance, some other measure of location, for instance) by generating random values of a random variable. In the present paper, MCMC was used to get the ERGM model parameters; and also in the fitting of the eigenmodel for the latent network modeling.
Network	Graph $G = (V, E)$, a construct of V vertices and E edges.
Network, Adjacency matrix	Given v number of vertices, the adjacency matrix is a $v \times v$ matrix, where any $[i, j]$ cell's value reflects with 0 or 1 whether a selected pair of vertices are connected or not
Network, Clustering coefficient	Transitivity, a descriptive measure to reflect the degree of connectedness [or process of clustering] in a graph by giving the ratio of connected triangles (triads of vertices) versus the total number of triangles in the graph.
Network, Diameter	The value of the longest distance (traceable route along edges and incident nodes) in a graph.
Network, Order	Descriptive; order of a network is the number of vertices.
Network, Size	Descriptive; size of a network is the total number of edges.
Network, Vertex, Betweenness	Descriptive of vertex centrality; quantifies how much a given vertex is between other pairs of vertices, thus has a connecting power.
Network, Vertex, Closeness	Descriptive of vertex centrality; assumes that important vertices are closer to others; its measure is inverse of total distances to other vertices.
Network, Vertex, Degree	Descriptive; the degree of a vertex is the number of edges incident on the given vertex. Vertex degrees are summarized as Degree Distribution of a graph.
Network, Vertex, Strength	Descriptive of vertex centrality; equals the sum of edge weights incident upon the given vertex, reflecting vertex centrality (importance). Vertex Strength Distribution is a histogram of vertex strength estimates of an entire graph.

Term	Definition
Node	Nodes (V,vertices) are representation of constructs, latent variables, subjects or any level of attributes in a network. Like edges, they can inherently bear any attributes.
Penalization	In regression, penalization introduces additional penalty to the sum of squared errors the model is trying to minimize. This way, by allowing a small increase in bias, the model is expected to exhibit a lower level of variance. The graphical LASSO method, used in the current paper, uses an absolute fraction of coefficientnts as penalties. With the applied penalty, insignificant or weak associations are expected to shrink towards 0.

References

- Abbott, M. J., & Rapee, R. M. 2004. Post-event rumination and negative self-appraisal in social phobia before and after treatment. *Journal of Abnormal Psychology*, 113(1), 136–144. <https://doi.org/10.1037/0021-843x.113.1.136>
- Aldao, A., & Nolen-Hoeksema, S. 2012. When are adaptive strategies most predictive of psychopathology? *Journal of Abnormal Psychology*, 121(1), 276–281. <https://doi.org/10.1037/a0023598>
- Allard, E. S., & Yaroslavsky, I. 2019. Attentional disengagement deficits predict brooding, but not reflection, over a one-year period. *Frontiers in Psychology*, 10. <https://doi.org/10.3389/fpsyg.2019.02282>
- Arana, F. G., & Rice, K. G. 2017. Cross-cultural validity of the ruminative responses scale in argentina and the united states. *Assessment*, 27(2), 309–320. <https://doi.org/10.1177/1073191117729204>
- Armey, M. F., Fresco, D. M., Moore, M. T., Mennin, D. S., Turk, C. L., Heimberg, R. G., Kecmanovic, J., & Alloy, L. B. 2009. Brooding and pondering. *Assessment*, 16(4), 315–327. <https://doi.org/10.1177/1073191109340388>
- Bernstein, E. E., Heeren, A., & McNally, R. J. 2019. Reexamining trait rumination as a system of repetitive negative thoughts: A network analysis. *Journal of Behavior Therapy and Experimental Psychiatry*, 63, 21–27. <https://doi.org/10.1016/j.jbtep.2018.12.005>
- Broderick, P. C., & Korteland, C. 2004. A prospective study of rumination and depression in early adolescence. *Clinical Child Psychology and Psychiatry*, 9(3), 383–394. <https://doi.org/10.1177/1359104504043920>
- Brose, A., Raedt, R. D., & Vanderhasselt, M.-A. 2020. Eight items of the ruminative response scale are sufficient to measure weekly within-person variation in rumination. *Current Psychology*, 41(7), 4210–4218. <https://doi.org/10.1007/s12144-020-00913-y>
- Bushman, B. J., Bonacci, A. M., Pedersen, W. C., Vasquez, E. A., & Miller, N. 2005. Chewing on it can chew you up: Effects of rumination on triggered displaced aggression. *Journal of Personality and Social Psychology*, 88(6), 969–983. <https://doi.org/10.1037/0022-3514.88.6.969>
- Butler, L. D., & Nolen-Hoeksema, S. 1994. Gender differences in responses to depressed mood in a college sample. *Sex Roles*, 30(5-6), 331–346. <https://doi.org/10.1007/bf01420597>
- Ciesla, J. A., & Roberts, J. E. 2007. Rumination, negative cognition, and their interactive effects on depressed mood. *Emotion*, 7(3), 555–565. <https://doi.org/10.1037/1528-3542.7.3.555>
- Costantini, G., Epskamp, S., Borsboom, D., Perugini, M., Möttus, R., Waldorp, L. J., & Cramer, A. O. J. 2015. State of the aRt personality research: A tutorial on network analysis of personality data in r. *Journal of Research in Personality*, 54, 13–29. <https://doi.org/10.1016/j.jrp.2014.07.003>
- Donald J. Robinaugh, R. J. M., Alexander J. Millner. 2016. Identifying highly influential nodes in the complicated grief network. *J Abnorm Psychol.*, 125(6). <https://doi.org/10.1037/abn0000181>
- Epskamp, S., Borsboom, D., & Fried, E. I. 2017a. Estimating psychological networks and their accuracy: A tutorial paper. *Behavior Research Methods*, 50(1), 195–212. <https://doi.org/10.3758/s13428-017-0862-1>
- Epskamp, S., Borsboom, D., & Fried, E. I. 2017b. *Estimating psychological networks and their accuracy: Supplementary materials*.
- Epskamp, S., Cramer, A. O. J., Waldorp, L. J., Schmittmann, V. D., & Borsboom, D. 2012. Qgraph: Network visualizations of relationships in psychometric data. *Journal of Statistical Software*, 48(4). <https://doi.org/10.18637/jss.v048.i04>
- Epskamp, S., & Fried, E. I. 2018. A tutorial on regularized partial correlation networks. *Psychological Methods*, 23(4), 617–634. <https://doi.org/10.1037/met0000167>
- Epskamp, S., Kruis, J., & Marsman, M. 2017. Estimating psychopathological networks: Be careful what you wish for. *PLOS ONE*, 12(6), e0179891. <https://doi.org/10.1371/journal.pone.0179891>
- Friedman, J., Hastie, T., & Tibshirani, R. 2007. Sparse inverse covariance estimation with the graphical lasso. *Biostatistics*, 9(3), 432–441. <https://doi.org/10.1093/biostatistics/kxm045>
- Glassman, L. H., Weierich, M. R., Hooley, J. M., Deliberto, T. L., & Nock, M. K. 2007. Child maltreatment, non-suicidal self-injury, and the mediating role of self-criticism. *Behaviour Research and Therapy*, 45(10), 2483–2490. <https://doi.org/10.1016/j.brat.2007.04.002>
- Goodreau, S. M., Kitts, J. A., & Morris, M. 2009. Birds of a feather, or friend of a friend? Using exponential random graph models to investigate adolescent social networks. *Demography*, 46(1), 103–125. <https://doi.org/10.1353/dem.0.0045>
- Hartley, S., Haddock, G., Sa, D. V. e, Emsley, R., & Barrowclough, C. 2013. An experience sampling study

- of worry and rumination in psychosis. *Psychological Medicine*, 44(8), 1605–1614. <https://doi.org/10.1017/s0033291713002080>
- Hasegawa, A. 2013. TRANSLATION AND INITIAL VALIDATION OF THE JAPANESE VERSION OF THE RUMINATIVE RESPONSES SCALE¹. *Psychological Reports*, 130718111108005. <https://doi.org/10.2466/02.08.pr0.112.3>
- Hayes, S. C., Strosahl, K., Wilson, K. G., Bissett, R. T., Pistorello, J., Toarmino, D., Polusny, M. A., Dykstra, T. A., Batten, S. V., Bergan, J., Stewart, S. H., Zvolensky, M. J., Eifert, G. H., Bond, F. W., Forsyth, J. P., Karekla, M., & McCurry, S. M. 2004. Measuring experiential avoidance: A preliminary test of a working model. *The Psychological Record*, 54(4), 553–578. <https://doi.org/10.1007/bf03395492>
- Heeren, A., Jones, P. J., & McNally, R. J. 2018. Mapping network connectivity among symptoms of social anxiety and comorbid depression in people with social anxiety disorder. *Journal of Affective Disorders*, 228, 75–82. <https://doi.org/10.1016/j.jad.2017.12.003>
- Hoff, P. D., Raftery, A. E., & Handcock, M. S. 2002. Latent space approaches to social network analysis. *Journal of the American Statistical Association*, 97(460), 1090–1098. <https://doi.org/10.1198/016214502388618906>
- Hunter, D. R. 2007. Curved exponential family models for social networks. *Social Networks*, 29(2), 216–230. <https://doi.org/10.1016/j.socnet.2006.08.005>
- Jones, P. 2017. *Networktools: Assorted and tools for identifying and important nodes and in networks*.
- Kolaczyk, G. C. E. D. 2020. *Statistical analysis of network data with r*. Springer-Verlag GmbH. https://www.ebook.de/de/product/39334469/gabor_csardi_eric_d_kolaczyk_statistical_analysis_of_network_data_with_r.html
- Kolubinski, D. C., Nikčević, A. V., Lawrence, J. A., & Spada, M. M. 2017. The metacognitions about self-critical rumination questionnaire. *Journal of Affective Disorders*, 220, 129–138. <https://doi.org/10.1016/j.jad.2017.06.002>
- Kovács, L. N., Takacs, Z. K., Tóth, Z., Simon, E., Schmelowszky, Ágoston, & Kökönyei, G. 2020. Rumination in major depressive and bipolar disorder – a meta-analysis. *Journal of Affective Disorders*, 276, 1131–1141. <https://doi.org/10.1016/j.jad.2020.07.131>
- Lee, S., & Kim, W. 2014. Cross-cultural adaptation, reliability, and validity of the revised korean version of ruminative response scale. *Psychiatry Investigation*, 11(1), 59. <https://doi.org/10.4306/pi.2014.11.1.59>
- Lyubomirsky, S., Layous, K., Chancellor, J., & Nelson, S. K. 2015. Thinking about rumination: The scholarly contributions and intellectual legacy of susan nolen-hoeksema. *Annual Review of Clinical Psychology*, 11(1), 1–22. <https://doi.org/10.1146/annurev-clinpsy-032814-112733>
- Lyubomirsky, S., & Nolen-Hoeksema, S. 1993. Self-perpetuating properties of dysphoric rumination. *Journal of Personality and Social Psychology*, 65(2), 339–349. <https://doi.org/10.1037/0022-3514.65.2.339>
- Mair, P. 2018. *Modern psychometrics with r*. Springer International Publishing. https://www.ebook.de/de/product/33047896/patrick_mair_modern_psychometrics_with_r.html
- McLaughlin, K. A., Aldao, A., Wisco, B. E., & Hilt, L. M. 2014. Rumination as a transdiagnostic factor underlying transitions between internalizing symptoms and aggressive behavior in early adolescents. *Journal of Abnormal Psychology*, 123(1), 13–23. <https://doi.org/10.1037/a0035358>
- McNally, R. J., Mair, P., Mugno, B. L., & Riemann, B. C. 2017. Co-morbid obsessive-compulsive disorder and depression: A bayesian network approach. *Psychological Medicine*, 47(7), 1204–1214. <https://doi.org/https://doi.org/10.1017/S0033291716003287>
- Memedovic, S., Slade, T., Ross, J., Darke, S., Mills, K. L., Marel, C., Burns, L., Lynskey, M., & Teesson, M. 2019. Rumination and problematic substance use among individuals with a long-term history of illicit drug use. *Drug and Alcohol Dependence*, 203, 44–50. <https://doi.org/10.1016/j.drugalcdep.2019.05.028>
- Milia, C., Kolubinski, D. C., & Spada, M. M. 2020. The effects of self-critical rumination on shame and stress: An experimental study. *Behavioural and Cognitive Psychotherapy*, 49(3), 272–286. <https://doi.org/10.1017/s1352465820000727>
- Moulds, M. L., Bisby, M. A., Wild, J., & Bryant, R. A. 2020. Rumination in posttraumatic stress disorder: A systematic review. *Clinical Psychology Review*, 82, 101910. <https://doi.org/10.1016/j.cpr.2020.101910>
- Moulds, M. L., Kandris, E., Starr, S., & Wong, A. C. M. 2007. The relationship between rumination, avoidance and depression in a non-clinical sample. *Behaviour Research and Therapy*, 45(2), 251–261. <https://doi.org/10.1016/j.brat.2006.03.003>
- Newman, M. E. J. 2010. *Networks - an introduction*. Oxford University Press.

- Nolen-Hoeksema, S. 2000. The role of rumination in depressive disorders and mixed anxiety/depressive symptoms. *Journal of Abnormal Psychology*, 109(3), 504–511. <https://doi.org/10.1037/0021-843x.109.3.504>
- Nolen-Hoeksema, S. 1991. Responses to depression and their effects on the duration of depressive episodes. *Journal of Abnormal Psychology*, 100(4), 569–582. <https://doi.org/10.1037/0021-843x.100.4.569>
- Nolen-Hoeksema, S., Larson, J., & Grayson, C. 1999. Explaining the gender difference in depressive symptoms. *Journal of Personality and Social Psychology*, 77(5), 1061–1072. <https://doi.org/10.1037/0022-3514.77.5.1061>
- Nolen-Hoeksema, S., & Watkins, E. R. 2011. A heuristic for developing transdiagnostic models of psychopathology. *Perspectives on Psychological Science*, 6(6), 589–609. <https://doi.org/10.1177/1745691611419672>
- Nolen-Hoeksema, S., Wisco, B. E., & Lyubomirsky, S. 2008. Rethinking rumination. *Perspectives on Psychological Science*, 3(5), 400–424. <https://doi.org/10.1111/j.1745-6924.2008.00088.x>
- O'Connor, R. C., & Noyce, R. 2008. Personality and cognitive processes: Self-criticism and different types of rumination as predictors of suicidal ideation. *Behaviour Research and Therapy*, 46(3), 392–401. <https://doi.org/10.1016/j.brat.2008.01.007>
- Papageorgiou, C., & Wells, A. 2001. Metacognitive beliefs about rumination in recurrent major depression. *Cognitive and Behavioral Practice*, 8(2), 160–164. [https://doi.org/10.1016/s1077-7229\(01\)80021-3](https://doi.org/10.1016/s1077-7229(01)80021-3)
- Peled, M., & Moretti, M. M. 2009. Ruminating on rumination: Are rumination on anger and sadness differentially related to aggression and depressed mood? *Journal of Psychopathology and Behavioral Assessment*, 32(1), 108–117. <https://doi.org/10.1007/s10862-009-9136-2>
- Pont, A. du, Rhee, S. H., Corley, R. P., Hewitt, J. K., & Friedman, N. P. 2017. Rumination and psychopathology: Are anger and depressive rumination differentially associated with internalizing and externalizing psychopathology? *Clinical Psychological Science*, 6(1), 18–31. <https://doi.org/10.1177/2167702617720747>
- Reichardt, J., & Bornholdt, S. 2006. Statistical mechanics of community detection. *Physical Review*, 74(1).
- Rose, A. V., & Rimes, K. A. 2018. Self-criticism self-report measures: Systematic review. *Psychology and Psychotherapy: Theory, Research and Practice*, 91(4), 450–489. <https://doi.org/10.1111/papt.12171>
- Ruiz, S. F., F. J. 2017. Psychometric properties and factor structure of the ruminative responses scale-short form in colombia. *International Journal of Psychology & Psychological Therapy*, 17(2), 199–208.
- Ruscio, A. M., Gentes, E. L., Jones, J. D., Hallion, L. S., Coleman, E. S., & Swendsen, J. 2015. Rumination predicts heightened responding to stressful life events in major depressive disorder and generalized anxiety disorder. *Journal of Abnormal Psychology*, 124(1), 17–26. <https://doi.org/10.1037/abn0000025>
- Schmittmann, V. D., Cramer, A. O. J., Waldorp, L. J., Epskamp, S., Kievit, R. A., & Borsboom, D. 2013. Deconstructing the construct: A network perspective on psychological phenomena. *New Ideas in Psychology*, 31(1), 43–53. <https://doi.org/10.1016/j.newideapsych.2011.02.007>
- Selby, E. A., Anestis, M. D., & Joiner, T. E. 2008. Understanding the relationship between emotional and behavioral dysregulation: Emotional cascades. *Behaviour Research and Therapy*, 46(5), 593–611. <https://doi.org/10.1016/j.brat.2008.02.002>
- Shahar, B., Doron, G., & Szepeswol, O. 2014. Childhood maltreatment, shame-proneness and self-criticism in social anxiety disorder: A sequential mediational model. *Clinical Psychology & Psychotherapy*, 22(6), 570–579. <https://doi.org/10.1002/cpp.1918>
- Smart, L. M., Peters, J. R., & Baer, R. A. 2015. Development and validation of a measure of self-critical rumination. *Assessment*, 23(3), 321–332. <https://doi.org/10.1177/1073191115573300>
- Spasojević, J., & Alloy, L. B. 2001. Rumination as a common mechanism relating depressive risk factors to depression. *Emotion*, 1(1), 25–37. <https://doi.org/10.1037/1528-3542.1.1.25>
- Strimmer, K. 2008a. Fdrtool: A versatile r package for estimating local and tail area-based false discovery rates. *Bioinformatics*, 24(12), 1461–1462. <https://doi.org/10.1093/bioinformatics/btn209>
- Strimmer, K. 2008b. A unified approach to false discovery rate estimation. *BMC Bioinformatics*, 9(1). <https://doi.org/10.1186/1471-2105-9-303>
- Tibshirani, R. 1996. Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society: Series B (Methodological)*, 58(1), 267–288. <https://doi.org/10.1111/j.2517-6161.1996.tb02080.x>
- Treynor, W. 2003. *Cognitive Therapy and Research*, 27(3), 247–259. <https://doi.org/10.1023/a:1023910315561>

- Valencia, P. D., & Paredes-Angeles, R. 2022. Revisiting the factor structure of the ruminative response scale: A bifactor approach. *Trends in Psychology*. <https://doi.org/10.1007/s43076-022-00214-2>
- Ward, A., Lyubomirsky, S., Sousa, L., & Nolen-Hoeksema, S. 2003. Can't quite commit: Rumination and uncertainty. *Personality and Social Psychology Bulletin*, 29(1), 96–107. <https://doi.org/10.1177/0146167202238375>
- Watkins, E. R. 2008. Constructive and unconstructive repetitive thought. *Psychological Bulletin*, 134(2), 163–206. <https://doi.org/10.1037/0033-2909.134.2.163>
- Watkins, E. R., & Roberts, H. 2020. Reflecting on rumination: Consequences, causes, mechanisms and treatment of rumination. *Behaviour Research and Therapy*, 127, 103573. <https://doi.org/10.1016/j.brat.2020.103573>
- Watkins, E., & Moulds, M. 2005. Distinct modes of ruminative self-focus: Impact of abstract versus concrete rumination on problem solving in depression. *Emotion*, 5(3), 319–328. <https://doi.org/10.1037/1528-3542.5.3.319>
- Werner, A. M., Tibubos, A. N., Rohrmann, S., & Reiss, N. 2019. The clinical trait self-criticism and its relation to psychopathology: A systematic review – update. *Journal of Affective Disorders*, 246, 530–547. <https://doi.org/10.1016/j.jad.2018.12.069>
- Whisman, M. A., Pont, A. du, & Butterworth, P. 2020. Longitudinal associations between rumination and depressive symptoms in a probability sample of adults. *Journal of Affective Disorders*, 260, 680–686. <https://doi.org/10.1016/j.jad.2019.09.035>
- Whitmer, A., & Gotlib, I. H. 2011. Brooding and reflection reconsidered: A factor analytic examination of rumination in currently depressed, formerly depressed, and never depressed individuals. *Cognitive Therapy and Research*, 35(2), 99–107. <https://doi.org/10.1007/s10608-011-9361-3>
- Wilkinson, P. O., Croudace, T. J., & Goodyer, I. M. 2013. Rumination, anxiety, depressive symptoms and subsequent depression in adolescents at risk for psychopathology: A longitudinal cohort study. *BMC Psychiatry*, 13(1). <https://doi.org/10.1186/1471-244x-13-250>
- Williams, Z. J., McKenney, E. E., & Gotham, K. O. 2021. Investigating the structure of trait rumination in autistic adults: A network analysis. *Autism*, 25(7), 2048–2063. <https://doi.org/10.1177/13623613211012855>
- Yasinski, C., Hayes, A. M., Ready, C. B., Abel, A., Görg, N., & Kuyken, W. 2019. Processes of change in cognitive behavioral therapy for treatment-resistant depression: Psychological flexibility, rumination, avoidance, and emotional processing. *Psychotherapy Research*, 30(8), 983–997. <https://doi.org/10.1080/10503307.2019.1699972>