




Kentucky Children's Hospital Pediatric Diabetic Ketoacidosis Guideline




Initial Evaluation—Frequently Performed in PED



PICU DKA Management and Disposition Guideline

Updated June 2021



Age	≤ 5 years	6-11 years	≥ 12 years
Total Daily Dose	0.5 units/kg/day	0.7 units/kg/day	1 unit/kg/day
Insulin glargine (Lantus)	0.25 units/kg	0.35 units/kg	0.5 units/kg
Meal-based CHO Coverage (1 unit of insulin per amount of CHO)	500 (wt in kg*0.5)	500 (wt in kg*0.7)	500 (wt in kg)
Insulin Sensitivity Factor (correction of high blood glucose over a certain target)	1500 (wt in kg*0.5)	1500 (wt in kg*0.7)	1500 (wt in kg)
	Use 1/2 unit increments Use insulin lispro (Humalog)	Use 1/2 unit increments Use insulin lispro (Humalog)	Use 1 unit increments Use insulin lispro (Humalog)
Can discuss special considerations with on-call Endocrinologist: children with almost normal Hgb A1c levels or mild DKA (would need lower doses) severely obese adolescents (would need higher doses)			

KCH DKA Guidelines: Management of Acute Cerebral Edema and Intracranial Hypertension

Updated June 2021

