

Department of Pediatrics

Division of Metabolism

University of Kentucky
Kentucky Clinic, J420
Lexington, Kentucky 40536-0284
Phone: 859-323-0396
Fax: 859-323-8275
www.ukhealthcare.uky.edu

**Protocol for patients with a risk for Hypoglycemia and
Metabolic Decompensation Due to Mitochondrial Disorders,
Fatty Acid Oxidation Problems and Related Abnormalities.**

This Protocol applies to patients who must refrain from oral nourishment or liquids for an operative procedure or diagnostic study.

- A. The critical time of fasting will vary by the age, size and diagnosis of the patient.
 - 1. Infants should not be fasting for more than two hours.
 - 2. Young children should not be fasting for more than four hours.
 - 3. For longer fasting children should be on IV replacement fluids of D 10/ ½ normal saline to run at 1 ¼ maintenance and continued until fully recovered. There should be a loading dose of L-Carnitine 100mg/kg body weight run in IV over 60 minutes. There should be Carnitine replacement 50 mg/kg IV every 6 hours until recovered. Potassium should be added to the IV solution after 6 hours.
- B. The Protocol should be continued until the patient is awake and taking oral fluids/ nourishment.

If any questions, please call the

Genetics and Metabolism Department at Kentucky Children's Hospital.

Phone: (859) 323-0396 Fax: (859) 257-1888