

|                                                                                                                          |                         |
|--------------------------------------------------------------------------------------------------------------------------|-------------------------|
| <b>University of Kentucky / UK HealthCare</b><br><b>Emergency Department</b><br><b>Policy and Procedure</b>              | <b>Policy # ED08-46</b> |
| <b>Title/Description:</b> Intranasal Fentanyl Guideline                                                                  |                         |
| <b>Purpose:</b> To provide guidelines for administration of intranasal Fentanyl in children in the Emergency Department. |                         |

[Policy](#)

[Procedure](#)

[Persons and Sites Affected](#)

[Policies Replaced](#)

[Effective Date](#)

[Review/Revision Dates](#)

**Policy**

This policy outlines procedures for administration of Fentanyl for pain management for children in the Emergency Department.

**Procedure**

*Indications*

- Isolated orthopedic injuries in children
- Other procedures as deemed appropriate by the ED provider

*Contraindications and Exclusion Criteria*

- Allergy or sensitivity to the medication
- Difficult airway
- Nasal obstruction or trauma
- Patients < 6 months of age
- Relative contraindications: asthma and controlled seizure disorder- consult attending prior to use

*Ordering and Dosing*

RNs may order intranasal Fentanyl if none of the contraindications or exclusion criteria noted above are present.

|               | <b>Fentanyl</b>                                                                                          |
|---------------|----------------------------------------------------------------------------------------------------------|
| Onset         | 2-5 minutes                                                                                              |
| Duration      | 30 – 60 minutes                                                                                          |
| Concentration | 50 mcg/mL                                                                                                |
| Dosing*       | 1.5 – 2 mcg/kg (0.1mL of overfill is needed to account for the deadspace in the atomizer)                |
| Monitoring    | Continuous pulse oximetry until return to baseline.<br>Assess vital signs 5 minutes post administration. |

\*divided between each nostril, max volume per nostril is 1 mL

### *Supplies*

The mucosal atomizer device (MAD) nasal adaptor will be used with a syringe. This atomizer delivers a metered dose of drug into the nares.

### *Documentation*

Follow standard ED documentation guidelines for analgesia (See Policy [ED08-26 Pain Management Guidelines](#)).

### *Adverse Events*

Monitor for respiratory depression and paradoxical reactions.

### **References**

Saunders, M., Adelgais, K., and Nelson, D. (2010). Use of intranasal fentanyl for the relief of pediatric orthopedic trauma pain. *Academic Emergency Medicine*, 17:11, 1155-1161.

|                                                                                                                                                                                                                                                                                                                                  |                                                |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------|
| <b>Persons and Sites Affected</b><br><input type="checkbox"/> Enterprise <input checked="" type="checkbox"/> Chandler <input checked="" type="checkbox"/> Good Samaritan <input type="checkbox"/> Kentucky Children's <input type="checkbox"/> Ambulatory <input checked="" type="checkbox"/> Department<br>Emergency Department |                                                |
| <b>Policies Replaced</b><br><input type="checkbox"/> Chandler HP <input type="checkbox"/> Good Samaritan <input type="checkbox"/> Kentucky Children's CH<br><input type="checkbox"/> Ambulatory KC <input type="checkbox"/> Other                                                                                                |                                                |
| <b>Effective Date:</b> 1/13                                                                                                                                                                                                                                                                                                      | <b>Review/Revision Dates:</b> 2/15, 3/18, 6/20 |
| <b>Approval by and date:</b><br><br>Signature <br>Name Patricia Kunz Howard, PhD, RN, CEN, CPEN, NE-BC, FAEN, FAAN<br>Date <u>6/23/2020</u><br>Review Team Leader                                                                             |                                                |