

Pediatric Massive transfusion Protocol

Inclusion Criteria

1. Age <16 yo
AND
2. Weight ≤50kg

Pediatric Trauma ABCD >3

- (1 point for each of the following)
1. Penetrating mechanism
 2. Positive FAST
 3. Elevated Pediatric-adjusted Shock Index (SIPA)*
 3. Lactate > 3.5
 4. Base Deficit > -8.8

OR

Attending Discretion

Pediatric CAT+ score, defined as use of >20ml/kg of total blood products within 1 hour

OR

*SIPA: Maximum HR/Minimal Systolic BP

4-6 years: Normal SIPA **1.22**

7-12 years: Normal SIPA **1.00**

13-16 years: Normal SIPA **0.90**

Activate PEDS MTP

- Contact Blood Bank (35401 or 35403) to "activate pediatric MTP"
- Give proper age and biologic sex of patient (Peds MTP ≤ 15 yrs)
- Surgeon must designate and place MTP order
- Send runner for cooler

Send blood bank specimen directly to Blood Bank (Tube #103 or #630)

Consider TXA

Children > 12yo: 1gm IV over 10 min followed by 1gm over 8 hours
Children <12yo: 15mg/kg (max 1gm) over 10 min followed by 2mg/kg/hr x 8 hours

Cooler #1: 4 RBC/ 4FFP/ 1 6pk Pheresis plt+,^

Bleeding Controlled?

No

Yes

Cooler #2: 4 RBC/ 4FFP/ 1 6pk Pheresis plt +,^

Bleeding Controlled?

No

Yes

Notify Blood Bank: Stop MTP

Cooler #3: 4 RBC/ 4FFP+

Bleeding Controlled?

No

Yes

Cooler #4: 4 RBC/ 4FFP/ 1 6pk Pheresis plt

Provider Info for Controlled Resuscitation Pack

< 10kg – 1 unit PRBCs/1FFP

10-20kg – 2 units PRBCs/2FFP/ (1/3) PLT*

20-50kg – 3 units PRBCs/3FFP/(1/2) PLT*

Volumes weight based ml 1:1:1 (pheresis PLT = 6pk)

*PCC can be considered if plasma shortage

+Consider calcium replacement after 20cc/kg or 2 units pRBC

^Consider cryo with coolers 2 & 3 (call if needed)

•<10kg- 1 unit (ml)

•10-20kg – 2 units (ml)

•20-50kg- 3 units (ml)