

TREATMENT AND EMERGENCY RECOMMENDATIONS FOR

ORNITHINE TRANSCARBAMYLASE DEFICIENCY (OTC DEFICIENCY)

PATIENT: _____ **MRN:** _____ **DOB:** _____

**The above patient is followed by our service for
Ornithine Transcarbamylase Deficiency (OTC Deficiency)**

Definition: OTC Deficiency is a sex-linked inherited disorder of the urea cycle function that is often lethal in infant males, has occasional expression in carrier females which may cause severe symptoms including death. OTC is the most frequent Urea Cycle Disorder with a frequency of 1/14,000.

Signs of Decompensation: Disorientation, lethargy, vomiting, irrational mood swings, headache, encephalopathy, elevated ammonia, coma and cerebral edema.

Treatments: Avoidance of high protein foods, supplementation with Arginine and/or Citrulline to support the blocked Urea Cycle and the use of ammonia scavengers (Sodium Benzoate or Sodium Phenylbutyrate).

Treatment During Metabolic Decompensation:

1. Reverse catabolism: IV D₁₀ with ½ NS at 1.5 times maintenance
2. IV Arginine HCl: 100 mg/kg IV over 1-2 hours, then 200 mg/kg IV per 24 hours
3. Ammonia scavengers
4. Obtain a STAT Ammonia and BMP (electrolytes and glucose)
5. **Notify the metabolic specialist immediately (after business hours this number allows you to speak with the geneticist on call:**

Kentucky Childrens Hospital
Division of Genetics and Metabolism

R. Stephen Amato, MD, PhD

859-323-0396

Kentucky Clinic

740 S. Limestone Street – J420

Lexington, KY 40536-0284

(After Hours: Please call 1-800-888-5533)



Department of Pediatrics / Division of Genetics & Metabolism / Kentucky Clinic
740 S. Limestone Street – J420 / Lexington, KY 40536-0284
Phone: 859-323-0396 / Fax: 859-257-1888