

- ☐ University of Kentucky A.B. Chandler Hospital
- ☐ UK HealthCare Good Samaritan Hospital
- ☐ UK HealthCare Ambulatory Services
- ☐ UK Dental and Oral Health Clinics

Pediatric Alteplase INCLUSION / EXCLUSION CRITERIA

(Patient Label Here)

Pediatric Alteplase INCLUSION / EXCLUSION CRITERIA

****Select box only if applicable****

Eligibility Criteria for IV Alteplase:	
☐ Diagnosis of ischemic stroke causing measurable neurological deficit ☐ Computed tomography (CT) rules out hemorrhage or non-stroke cause of deficit ☐ 10 years old or older ☐ Time to treatment less than 3 hours of confirmed time Last Known Well, or 4.5 hours with additional warning	
Contraindications IV Alteplase: ☐ Current intracranial hemorrhage ☐ Suspicion of subarachnoid hemorrhage ☐ Intra-axial intracranial neoplasm ☐ History of intracranial / intraspinal surgery within 3 months ☐ Severe head trauma within 3 months ☐ Gastrointestinal malignancy or active internal bleeding within 21 days ☐ Bleeding diathesis ☐ Platelets less than 100,000/mm³ ☐ International Normalized Ratio (INR) greater than 1.7 ☐ Activated partial thromboplastin time greater than 40 seconds ☐ Prothrombin time greater than 15 seconds ☐ Warfarin use with INR greater than 1.7 ☐ Treatment dose of low molecular weight heparin within 24 hours (<i>not prophylactic dose</i>) ☐ Direct thrombin inhibitors or direct factor Xa inhibitors (<i>unless not taken within 48 hours or absence of therapeutic effect on appropriate screening tests</i>) ☐ Sustained BP greater than 95 percentile for the patient's age/sex/BP parameter (Please refer to chart for BP parameter values that are specific for the patient's age/sex) ☐ Acute bacterial endocarditis ☐ Aortic Arch dissection ☐ Suspected moyamoya / fat, air or bone marrow embolism	Warnings / Use Clinical Judgment IV Alteplase: ☐ History of intracranial hemorrhage ☐ History of ischemic stroke within 3 months <i>(considerations: size, location and timing of prior stroke)</i> ☐ Extensive regions of clear hypoattenuation on initial CT ☐ Unruptured / unsecured AVM ☐ >10 mm unruptured / unsecured aneurysm ☐ Major surgery within 14 days ☐ Major trauma within 14 days ☐ Arterial puncture of noncompressible vessel within 7 days ☐ History of gastrointestinal or genitourinary hemorrhage ☐ Malignancy with life expectancy less than one year ☐ History bleeding diathesis ☐ Hemorrhagic ophthalmic condition ☐ Acute pericarditis ☐ History of myocardial infarction involving left anterior myocardium within 3 months ☐ Left sided heart thrombus ☐ Pregnancy
Additional Warning for 3 - 4.5 hour window IV Alteplase: 3 hour window contraindications and warnings continue to apply, PLUS: Warnings / Use Clinical Judgment: ☐ NIH Stroke Scale greater than 25	
☐ No contraindications for IV Alteplase administration ☐ Warning(s) considered	
IV Alteplase does <i>not</i> preclude mechanical retrieval, which can be considered out to 24 hours from last known well	
☐ Emergent Large Vessel Occlusion (ELVO) considered	
Maximum dose = 90 mg; Calculation: = 0.9 mg/kg; 10% over one minute IV push and remainder of calculated dose over one hour IV infusion; IV normal saline flush infused at the same rate immediately following IV Alteplase, to clear the line	

Physician Signature: _____
12/17/17

Date & Time: _____

- ☐ University of Kentucky A.B. Chandler Hospital
☐ UK HealthCare Good Samaritan Hospital
☐ UK HealthCare Ambulatory Services
☐ UK Dental and Oral Health Clinics

Mechanical Thrombectomy Intervention Ischemic Stroke Protocol Eligibility Checklist

(Patient Label Here)

Intervention Ischemic Stroke Protocol Eligibility Checklist		
Inclusion for mechanical thrombectomy : call for consult	Yes	No
NIHSS greater than or equal to 2		
Less than 24 hours from Last Known Well time (Mechanical thrombolysis time may be extended up to 24 hours for posterior circulation strokes)		
Emergent Large Vessel Occlusion seen on CTA		
Contact the neuro - interventionalist for any possible large vessel occlusion no more than 15 minutes from completion of CTA Time neuro - interventionalist consulted: _____ Dr. Fraser : pager - 330-3414 Dr. Sheikhi: pager- 330-0711 Dr. Pahwa: pager- 259-5237 Dr. Dornbos: pager- 330- 3122 Angio suite : 323 - 0348		
IV Alteplase Dosing and Treatment Standardized intra-VENOUS Alteplase dose for stroke: -- 0.9 mg/kg(maximum total dose -90 mg) -- 10% of total dose given over one minute IV push and remainder of calculated dose over one hour IV infusion -- All excess Alteplase)needs to be removed from vial prior to dose preparation to prevent patient from receiving excess Alteplase -- This needs to be followed by an IV Normal Saline fluid drip at the same rate to ensure total IV Alteplase dose -- administration Treatment for BP greater than 95th percentile for patient age/sex (Please reference chart in Appendix A on page 3 for BP parameters that are specific for age/sex) : - Hydralazine IV:0.1-0.2mg/kg every 4-6 hours (max 20 mg/dose) - Labetalol IV: intermittent bolus doses of 0.2-1mg/kg/dose (max 20mg/dose) During the first 24 hours post Alteplase administration the following should be avoided: -- BP greater than 95th percentile for patient age/sex (Please reference chart for BP parameters that are specific for age/sex) -- Anticoagulants, Antiplatelet agents, Antithrombotics -- Invasive procedures, invasive lines, catheters or tubes unless deemed medically necessary -- A physician order will be placed for the appropriate bedrest time/mobility status based on patients NIH, neurological status and hemodynamically stability. Monitoring: ** IF any adverse reaction - FIRST stop Alteplase and notify physician ** -Enhanced neurological checks, NIHSS, and vital signs as ordered *Vital signs and Enhanced Neuro checks: baseline prior to Alteplase bolus, q15 mins x 2 hrs, q30 mins x 6 hrs, q1 hr x 16 hrs, q2 hrs x 24 hrs, then q4 hrs x 24 hrs, q8 hrs and prn with any acute neurological decline until discharge *NIHSS: baseline prior to Alteplase bolus, repeat at 30 min (during infusion), at 60 min (end of infusion), then q2 hrs x 48 hrs, q4 hrs x 24 hrs, q12 hrs and prn with acute neurological decline until discharge -- Signs of bleeding -- Neurological deterioration (change of 2+ points on NIHSS) -- Complaints of new headache or nausea -- Allergic reactions (angioedema)		
Physician's Signature _____	Printed Name _____	Pager _____ / Date/Time _____

- ☐ University of Kentucky A.B. Chandler Hospital
- ☐ UK HealthCare Good Samaritan Hospital
- ☐ UK HealthCare Ambulatory Services
- ☐ UK Dental and Oral Health Clinics

Pediatric Alteplase
INCLUSION / EXCLUSION

(Patient Label Here)

Appendix A:

Blood pressure parameters:

Please maintain BP between the 50th and 95th percentiles

Boys			Girls		
Age	SBP	DBP	Age	SBP	DBP
1	80-106	34-58	1	83-107	38-60
2	84-110	39-63	2	85-109	43-65
3	86-113	44-67	3	86-110	47-69
4	88-115	47-71	4	88-112	50-72
5	90-116	50-74	5	89-113	52-74
6	91-117	53-76	6	91-115	54-76
7	92-119	55-78	7	93-116	55-77
8	94-120	56-80	8	95-118	57-78
9	95-121	57-81	9	96-120	58-79
10	97-123	58-82	10	98-122	59-80
11	99-125	59-82	11	100-124	60-81
12	101-127	59-83	12	102-126	61-82
13	104-130	60-83	13	104-128	62-83
14	106-132	60-84	14	106-129	63-84
15	109-135	61-85	15	107-131	64-85