

Molecular Tumor Board Recommendation
August 7, 2024

Dear Dr. Patel:

The Markey Cancer Center Molecular Tumor Board met on August 6, 2024 and reviewed your patient, Rosa Martinez Alejandro (110710532; carcinoma of the sigmoid colon). After reviewing the genomic report (Tempus xT, Tempus IHC, Tempus xR collected February 2, 2024, Tempus xF collected March 28, 2024) and clinical characteristics of the case, the MTB recommends consideration of the following:

1. If progression, consider off-label olaparib targeting the ATM mutation (E28*). This is a category 3 evidence level recommendation.
2. If progression, consider off-label erdafitinib targeting the FGFR3-DSP rearrangement. This is a category 3 evidence level recommendation. Please note, in the single arm, phase 2 RAGNAR study, 217 patients with solid tumors (with FGFR alterations) were treated with erdafitinib, which produced an ORR of 30% and a disease control rate of 74%. Of note, the 8 colorectal cancer patients in this study had little or no response which is difficult to interpret given the small sample size (PMID: 37541273). The MTB favors targeting the ATM mutation first.
 - Reference: Pant S, Schuler M, Iyer G, et al. Erdafitinib in patients with advanced solid tumours with FGFR alterations (RAGNAR): an international, single-arm, phase 2 study. *Lancet Oncol.* 2023;24(8):925-935. doi:10.1016/S1470-2045(23)00275-9
3. At time of relapse, consider repeating NGS testing on a new tumor sample or liquid biopsy given the relatively low VAF of the ATM mutation.

Thank-you for submitting this case to the Markey Cancer Center MTB.

Signature: Derek B. Allison, MD Digitally signed by Derek B. Allison, MD
Date: 2024.08.14 12:22:13 -04'00' Date: _____

Signature: Justine Pickarski Digitally signed by Justine Pickarski
Date: 2024.08.08 08:44:06 -04'00' Date: _____

Signature: Rachel W. Miller, MD Digitally signed by Rachel W. Miller, MD
Date: 2024.08.13 18:49:02 -04'00' Date: _____

Level of Evidence	Interpretation	Supporting Evidence
1	FDA-labeled indication or NCCN-recommended; substantial evidence supporting use (standard of care)	Meta-analysis, RCTs, etc. OncoKB level 1
2	Off-label use; substantial evidence supporting use	Phase II studies and above OncoKB level 2 and above
3	Off-label use; low to moderate evidence supporting use; MTB recommends, clinical trial preferred	Phase I, case studies and reports OncoKB level 3 and below
4	Off-label; no evidence supporting use; enrollment in clinical trial recommended; report experience to MTB if used outside clinical trial	Little to no evidence (pre-clinical, animal/cellular models), mechanistically plausible