

## Patient

**Name:** WHITLEY, AMY J  
**Date of Birth:** 07/04/1955  
**Sex:** Female  
**Case Number:** TN25-173573  
**Diagnosis:** Mucinous adenocarcinoma

## Specimen Information

**Primary Tumor Site:** Lower lobe, lung  
**Specimen Site:** Lower lobe, lung  
**Specimen ID:** C25-03977-B1  
**Specimen Collected:** 02-Apr-2025  
**Test Report Date:** 11-May-2025

## Ordered By

**Janeesh Sekkath-Veedu, MD**  
**UK HealthCare - Albert B Chandler**  
**Hospital - Markey Cancer Center**  
 800 Rose Street Whitney Hendrickson,  
 Building Room 145  
 Lexington, KY 40536 USA  
 (859) 257-4488

## Results with Therapy Associations

| BIOMARKER | METHOD | ANALYTE   | RESULT                          | THERAPY ASSOCIATION |                                                                             | BIOMARKER LEVEL*   |
|-----------|--------|-----------|---------------------------------|---------------------|-----------------------------------------------------------------------------|--------------------|
| MET       | IHC    | Protein   | Positive   3+, 35%              | BENEFIT             | telisotuzumab vedotin                                                       | Level 3            |
|           | Seq    | RNA-Tumor | Variant Transcript Not Detected | LACK OF BENEFIT     | capmatinib, tepotinib<br>crizotinib                                         | Level 2<br>Level 2 |
| ALK       | IHC    | Protein   | Negative   0                    | LACK OF BENEFIT     | alectinib, ceritinib, crizotinib, lorlatinib<br>brigatinib                  | Level 1<br>Level 2 |
|           | Seq    | RNA-Tumor | Fusion Not Detected             |                     | alectinib, brigatinib, ceritinib, crizotinib, lorlatinib                    | Level 2            |
| BRAF      | Seq    | DNA-Tumor | Mutation Not Detected           | LACK OF BENEFIT     | dabrafenib, dabrafenib + trametinib, encorafenib + binimetinib, vemurafenib | Level 2            |
| EGFR      | Seq    | DNA-Tumor | Mutation Not Detected           | LACK OF BENEFIT     | afatinib, dacomitinib, erlotinib, gefitinib, osimertinib                    | Level 2            |
| RET       | Seq    | RNA-Tumor | Fusion Not Detected             | LACK OF BENEFIT     | pralsetinib, selpercatinib                                                  | Level 2            |
| ROS1      | Seq    | RNA-Tumor | Fusion Not Detected             | LACK OF BENEFIT     | ceritinib, crizotinib, entrectinib, lorlatinib, repotrectinib               | Level 2            |

\* Biomarker reporting classification - Level 1 – MI Cancer Seek<sup>™</sup> or other companion diagnostic (CDx) performed as part of professional services; Level 2 – Strong evidence of clinical significance or is endorsed by standard clinical guidelines; Level 3 – Potential clinical significance. Bolded benefit therapies, if present, highlight the most clinically significant findings.

The selection of any, all, or none of the matched therapies resides solely with the discretion of the treating physician. Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking into consideration all available information concerning the patient's condition, the FDA prescribing information for any therapeutic, and in accordance with the applicable standard of care. Whether or not a particular patient will benefit from a selected therapy is based on many factors and can vary significantly. All trademarks and registered trademarks are the property of their respective owners.

## Important Note

This content is provided as part of the professional services. For MI Cancer Seek™ results, see CDx Associated Findings.

The MI GPSai (Genomic Probability Score) is an algorithm that uses a tumor's molecular signature to provide a differential diagnosis as to a tumor's origin and in this case yields several possible sites of origin for this tumor. All therapeutic associations in this report are based on a primary site/lineage of LUNG per the submitted requisition (probability 73%). Interpreting GPS results requires correlation with all available clinical, radiographic, and pathologic information. If a change in the tumor type is desired, which may affect the therapeutic associations on the report, please contact Caris Life Sciences and an amended report will be issued.

This patient's sample is positive for expression of MET by immunohistochemistry (Ventana, SP44) and is associated with telisotuzumab vedotin, pending FDA approval (NCT04928846). Please see [clinicaltrials.gov](https://clinicaltrials.gov) for additional trials which may require different thresholds for MET positivity (e.g. NCT05652868, NCT03175224, NCT03797391, NCT06083857, NCT05845671).

For information on Caris GPSai™, please refer to relevant section of the report (Additional Results Page 1).

## Cancer-Type Relevant Biomarkers

| Biomarker               | Method     | Analyte          | Result                                    |
|-------------------------|------------|------------------|-------------------------------------------|
| <b>KRAS</b>             | <b>Seq</b> | <b>DNA-Tumor</b> | <b>Pathogenic Variant Exon 2   p.G12D</b> |
| MSI                     | Seq        | DNA-Tumor        | Stable                                    |
| NTRK1/2/3               | Seq        | RNA-Tumor        | Fusion Not Detected                       |
| Tumor Mutational Burden | Seq        | DNA-Tumor        | Low, 2 mut/Mb                             |
| ALK                     | Seq        | DNA-Tumor        | Mutation Not Detected                     |
| BRAF                    | Seq        | RNA-Tumor        | Fusion Not Detected                       |
| ERBB2 (Her2/Neu)        | CNA-Seq    | DNA-Tumor        | Amplification Not Detected                |
|                         | IHC        | Protein          | Negative   Score 1+                       |
|                         | Seq        | DNA-Tumor        | Mutation Not Detected                     |
| FGFR3                   | Seq        | RNA-Tumor        | Fusion Not Detected                       |
| KEAP1                   | Seq        | DNA-Tumor        | Mutation Not Detected                     |
| MET                     | CNA-Seq    | DNA-Tumor        | Amplification Not Detected                |
|                         | Seq        | DNA-Tumor        | Mutation Not Detected                     |

| Biomarker     | Method  | Analyte   | Result                                             |
|---------------|---------|-----------|----------------------------------------------------|
| MTAP          | CNA-Seq | DNA-Tumor | Deletion Not Detected                              |
| NFE2L2        | Seq     | DNA-Tumor | Variant of Uncertain Significance Exon 5   p.Y576H |
| NRG1          | Seq     | RNA-Tumor | Fusion Not Detected                                |
| PD-L1 (22c3)  | IHC     | Protein   | Negative, TPS: 0%                                  |
| PD-L1 (28-8)  | IHC     | Protein   | Negative   0%                                      |
| PD-L1 (SP142) | IHC     | Protein   | Negative, IC: 5%<br>Negative, TC: 0%               |
| PD-L1 (SP263) | IHC     | Protein   | Negative, TC: 0%                                   |
| RB1           | Seq     | DNA-Tumor | Mutation Not Detected                              |
| RET           | Seq     | DNA-Tumor | Mutation Not Detected                              |
| STK11         | Seq     | DNA-Tumor | Mutation Not Detected                              |
| TP53          | Seq     | DNA-Tumor | Mutation Not Detected                              |

**PATIENT:** WHITLEY, AMY J (04-JUL-1955)

**TN25-173573**

**PHYSICIAN:** JANEESH SEKKATH-VEEDU, MD

## Genomic Signatures

| Biomarker                            | Method | Analyte   | Result                                                                               |
|--------------------------------------|--------|-----------|--------------------------------------------------------------------------------------|
| Microsatellite Instability (MSI)     | Seq    | DNA-Tumor | Stable                                                                               |
| Tumor Mutational Burden (TMB)        | Seq    | DNA-Tumor | <div> <div>Result: Low</div> <div> <div>2</div> <div>Low 10 High</div> </div> </div> |
| Genomic Loss of Heterozygosity (LOH) | Seq    | DNA-Tumor | Low - 2% of tested genomic segments exhibited LOH (assay threshold is $\geq 16\%$ )  |

## Genes Tested with Pathogenic or Likely Pathogenic Alterations

| Gene | Method | Analyte   | Variant Interpretation    | Protein Alteration | Exon | DNA Alteration | Variant Frequency % |
|------|--------|-----------|---------------------------|--------------------|------|----------------|---------------------|
| CDH1 | Seq    | DNA-Tumor | Likely Pathogenic Variant | p.D400N            | 9    | c.1198G>A      | 13                  |
| KRAS | Seq    | DNA-Tumor | Pathogenic Variant        | p.G12D             | 2    | c.35G>A        | 18                  |

Unclassified alterations for DNA and RNA sequencing can be found in the MI Portal.  
Formal nucleotide nomenclature and gene reference sequences can be found in the Appendix of this report.

## Genes Tested with Variants of Uncertain Significance

| Gene   | Method | Analyte   | Variant Interpretation            | Protein Alteration | Exon | DNA Alteration | Variant Frequency % |
|--------|--------|-----------|-----------------------------------|--------------------|------|----------------|---------------------|
| NFE2L2 | Seq    | DNA-Tumor | Variant of Uncertain Significance | p.Y576H            | 5    | c.1726T>C      | 50                  |

Additional Variants of Uncertain Significance can be found in the MI Portal.

## Predicted Human Leukocyte Antigen (HLA) Genotype Results

The predicted HLA genotype results are summarized in the table below. Please note that HLA typing via tumor tissue sequencing is not a substitute for histocompatibility testing performed using peripheral blood. It is recommended to confirm the patient's HLA type with an appropriate assay.

| Gene        | Method | Analyte   | Genotype         |
|-------------|--------|-----------|------------------|
| MHC CLASS I |        |           |                  |
| HLA-A       | Seq    | DNA-Tumor | A*25:01, A*32:01 |
| HLA-B       | Seq    | DNA-Tumor | B*49:01, B*51:01 |
| HLA-C       | Seq    | DNA-Tumor | C*01:02, C*07:01 |

HLA genotypes with only one allele are either homozygous or have loss-of-heterozygosity at that position.

## Immunohistochemistry Results

| Biomarker        | Result              | Biomarker     | Result                               |
|------------------|---------------------|---------------|--------------------------------------|
| ALK              | Negative   0        | PD-L1 (28-8)  | Negative   0%                        |
| ERBB2 (Her2/Neu) | Negative   Score 1+ | PD-L1 (SP142) | Negative, IC: 5%<br>Negative, TC: 0% |
| MET              | Positive   3+, 35%  | PD-L1 (SP263) | Negative, TC: 0%                     |
| PD-L1 (22c3)     | Negative, TPS: 0%   |               |                                      |

## Genes Tested with Indeterminate Results by Tumor DNA Sequencing

|         |        |      |       |      |       |       |       |       |      |         |        |
|---------|--------|------|-------|------|-------|-------|-------|-------|------|---------|--------|
| ATP6AP2 | COL2A1 | CUL3 | DGCR8 | MDH2 | PLCB4 | PRDM6 | PRKD1 | RASA1 | REST | SMARCA2 | TRIM28 |
|---------|--------|------|-------|------|-------|-------|-------|-------|------|---------|--------|

Genes in this table were ruled indeterminate due to low coverage for some or all exons.

## Genes Tested with Intermediate CNA Results by Tumor DNA Sequencing

|          |       |       |      |        |  |  |  |  |  |  |  |
|----------|-------|-------|------|--------|--|--|--|--|--|--|--|
| APOBEC3B | CSF1R | FGFR2 | JAK2 | PDGFRB |  |  |  |  |  |  |  |
|----------|-------|-------|------|--------|--|--|--|--|--|--|--|

The results in this report were curated to represent biomarkers most relevant for the submitted cancer type. These include results important for therapeutic decision-making, as well as notable alterations in other biomarkers known to be involved in oncogenesis. Additional results, including genes with normal findings, variants of uncertain significance, or unclassified alterations can be found in the MI Portal at [miportal.carismolecularintelligence.com](https://miportal.carismolecularintelligence.com). If you do not have an MI Portal account, or need assistance accessing it, please contact Caris Customer Support at (888) 979-8669.

## Notes of Significance

### SEE APPENDIX FOR DETAILS

CDx reports are generated through automated bioinformatics processing of a pre-defined list of genes and variants. Pathogenic or likely pathogenic variants identified outside the FDA-approved list will not be included in the CDx report. In addition, downgrading of variants approved by board-eligible/board-certified geneticists and pathologists may have occurred through manual variant annotation carried out by Caris' professional services. To ensure optimal coverage, multiple sequencing runs may be performed for a given case. This may account for some discrepancies observed between CDx reports and professional services.

Clinical Trials Connector<sup>™</sup> opportunities based on biomarker expression: 72 Targeted Therapies. See page 6 for details.

**Please Note:** The MI GPSai (Genomic Probability Score) is an algorithm that uses a tumor's molecular signature to provide a differential diagnosis as to a tumor's origin and in this case yields several possible sites of origin for this tumor. All therapeutic associations in this report are based on a primary site/lineage of LUNG per the submitted requisition (probability 73%). Interpreting GPS results requires correlation with all available clinical, radiographic, and pathologic information. If a change in the tumor type is desired, which may affect the therapeutic associations on the report, please contact Caris Life Sciences and an amended report will be issued.

## Specimen Information

**Specimen ID:** C25-03977-B1

**Specimen Collected:** 04/02/2025

**Specimen Received:** 04/30/2025

**Testing Initiated:** 05/02/2025

**Test Ordered\*:** MI Profile<sup>™</sup> (MI Cancer Seek<sup>™</sup> + IHCs and Other Tests by Tumor Type)

\* If the submitted specimen is inadequate, only a subset of the ordered testing may be reported.

**Gross Description:** 1 (A) Paraffin Block - Client ID(C25-03977-B1) from UK HealthCare - Albert B Chandler Hospital, Lexington, KY, with the corresponding pathology report labeled "C25-03977".

**Dissection Information:** Molecular testing of this specimen was performed after harvesting of targeted tissues with an approved manual microdissection technique. Candidate slides were examined under a microscope and areas containing tumor cells (and separately normal cells, when necessary for testing) were circled. A laboratory technician harvested targeted tissues for extraction from the marked areas using a dissection microscope.

## Clinical Trials Connector™

The Clinical Trials Connector lists agents that are matched to available clinical trials according to biomarker status. In some instances, older-generation agents may still be relevant in the context of new combination strategies and, therefore, will still appear on this report.

**Therapeutic agents listed below may or may not be currently FDA approved for the tumor type tested.**

Please see <https://clinicaltrials.gov/> for more information.

| TARGETED THERAPIES (72)    |           |        |           |                                                                                                 |
|----------------------------|-----------|--------|-----------|-------------------------------------------------------------------------------------------------|
| Drug Class                 | Biomarker | Method | Analyte   | Investigational Agent(s)                                                                        |
| Cancer vaccines (1)        | KRAS      | NGS    | DNA-Tumor | OSE2101                                                                                         |
| cMET-targeted therapy (39) | MET       | IHC    | Protein   | brigatinib, cabozantinib, capmatinib, crizotinib, savolitinib, telisotuzumab vedotin, tepotinib |
| Dual MEK/RAF inhibitor (3) | KRAS      | NGS    | DNA-Tumor | avutometinib                                                                                    |
| MEK inhibitors (29)        | KRAS      | NGS    | DNA-Tumor | binimetinib, cobimetinib, mirdametinib, trametinib                                              |

( ) = represents the total number of clinical trials identified by the Clinical Trials Connector for the provided drug class or table.

The Clinical Trials Connector may include trials that enroll patients with additional screening of molecular alterations. In some instances, only specific gene variants may be eligible.

## Disclaimer

Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking into consideration all available information concerning the patient's condition, prescribing information for any therapeutic, and in accordance with the applicable standard of care. Drug associations provided in this report do not guarantee that any particular agent will be effective for the treatment of any patient or for any particular condition. Caris Life Sciences<sup>®</sup> expressly disclaims and makes no representation or warranty whatsoever relating, directly or indirectly, to the performance of services, including any information provided and/or conclusions drawn from therapies that are included or omitted from this report. Whether or not a particular patient will benefit from a selected therapy is based on many factors and can vary significantly. The selection of therapy, if any, resides solely in the discretion of the treating physician and the tests should not be considered a companion diagnostic.

Caris MPI, Inc. d/b/a Caris Life Sciences is certified under the Clinical Laboratory Improvement Amendments (CLIA) as qualified to perform high complexity clinical laboratory testing, including all Caris molecular profiling assays. Individual assays that are available through Caris molecular profiling include both Laboratory Developed Tests (LDT) and U.S. Food and Drug Administration (FDA) approved or cleared tests. In addition, certain tests have been CE-marked as a general IVD under the In Vitro Diagnostic Directive (IVDD) 98/79/EC. Offered LDTs were developed and their performance characteristics determined by Caris. Certain tests have not been cleared or approved by the FDA. Caris LDTs are used for clinical purposes. They are not investigational or for research. Caris' CLIA certification number is located at the bottom of each page of this report.

The information presented in the Clinical Trials Connector<sup>™</sup> section of this report, if applicable, is compiled from sources believed to be reliable and current. However, the accuracy and completeness of the information provided herein cannot be guaranteed. The clinical trials information present in the biomarker description was compiled from [www.clinicaltrials.gov](http://www.clinicaltrials.gov). The contents are to be used only as a guide, and health care providers should employ their best comprehensive judgment in interpreting this information for a particular patient. Specific eligibility criteria for each clinical trial should be reviewed as additional inclusion criteria may apply.

All materials, documents, data, data software, information and/or inventions supplied to customers by or on behalf of Caris or created by either party relating to the services shall be and remain the sole and exclusive property of Caris. Customer shall not use or disclose the information provided by Caris through the services or related reports except in connection with the treatment of the patient for whom the services were ordered and shall not use such property for, or disseminate such property to, any third parties without expressed written consent from Caris. Customer shall deliver all such property to Caris immediately upon demand or upon Caris ceasing to provide the services. The technical and professional component of all testing was performed at the laboratory location displayed in the footer unless otherwise notated in the report.

Caris molecular testing is subject to Caris' intellectual property. Patent [www.CarisLifeSciences.com/ip](http://www.CarisLifeSciences.com/ip).

Professional Component Performed: -- Location 18, Irvine, CA. Caleb Cheng, MD, MBA, Lab Director.



Electronic Signature  
Caleb Cheng, MD, MBA  
11-May-2025



(01)00860008613325(21)H7KNLDSXF

MI Cancer Seek<sup>™</sup>

## Patient

**Name:** WHITLEY, AMY J  
**Date of Birth:** 07/04/1955  
**Sex:** Female  
**Case Number:** TN25-173573  
**CDx Report Generated:** 05/09/2025

## Specimen Information

**Diagnosis and Tumor Type:**  
Mucinous adenocarcinoma, Lower lobe, lung  
**Specimen Site:** Lower lobe, lung  
**Specimen ID:** C25-03977-B1  
**Specimen Type:** Formalin-fixed paraffin-embedded  
**Specimen Collected:** 04/02/2025

## Ordered By

**Janeesh Sekkath-Veedu, MD**  
UK HealthCare - Albert B Chandler  
Hospital - Markey Cancer Center

## CDx Associated Findings

### Genomic Findings Detected

### FDA-approved Therapeutic Options

**Findings did not yield any biomarker results with Companion Diagnostic (CDx) Claims.**

**EGFR** L858R

Not Detected

**EGFR** exon 19 deletions

Not Detected

**Microsatellite Instability (MSI)**

Not MSI-H

## Tumor Profiling Results

MI Cancer Seek is FDA-approved to provide tumor mutation profiling results for previously diagnosed oncology patients with solid tumors.

## Other Alterations and Biomarkers Identified

Results reported in this section are not prescriptive or conclusive for labeled use of any specific therapeutic product. Confirmation of tumor mutation status using an FDA-approved CDx test is needed for therapeutic use.

**Tumor Mutational Burden (TMB)**

Not Reportable (see limitations)

**CDH1**

D400N

**KRAS**

G12D



## MI Cancer Seek™

### Intended Use:

MI Cancer Seek is a next-generation sequencing (NGS) based in vitro diagnostic (IVD) device using total nucleic acid (TNA) isolated from formalin-fixed paraffin-embedded (FFPE) tumor tissue specimens for the detection of single nucleotide variants (SNVs) and insertions and deletions (indels) in 228 genes, microsatellite instability (MSI), tumor mutational burden (TMB) in patients with previously diagnosed solid tumors, and copy number amplification (CNA) in one gene in patients with breast cancer.

MI Cancer Seek is intended as a companion diagnostic to identify patients who may benefit from treatment with the targeted therapies listed in Table 1 below, in accordance with the approved therapeutic product labeling.

Additionally, MI Cancer Seek is intended to provide tumor mutational profiling to be used by qualified healthcare professionals in accordance with professional oncology guidelines for cancer patients with previously diagnosed solid malignant neoplasms. Genomic findings other than those listed in Table 1 are not prescriptive or conclusive for labeled use of any specific therapeutic product.

Table 1. MI Cancer Seek Companion Diagnostic Indications

| INDICATION                         | BIOMARKER                                                                                                                               | THERAPY                                                                    |
|------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------|
| Breast Cancer                      | <i>PIK3CA</i> (C420R, E542K, E545A, E545D [1635G>T only], E545G, E545K, Q546E, Q546R, H1047L, H1047R, H1047Y)                           | PIQRAY® (alpelisib)                                                        |
| Colorectal Cancer (CRC)            | <i>KRAS</i> wild-type (absence of mutations in exons 2, 3, and 4) and <i>NRAS</i> wild-type (absence of mutations in exons 2, 3, and 4) | VECTIBIX® (panitumumab)                                                    |
|                                    | <i>BRAF</i> V600E                                                                                                                       | BRAFTOVI® (encorafenib) in combination with ERBITUX® (cetuximab)           |
| Melanoma                           | <i>BRAF</i> V600E                                                                                                                       | BRAF Inhibitors approved by FDA*                                           |
|                                    | <i>BRAF</i> V600E or <i>BRAF</i> V600K                                                                                                  | MEKINIST® (trametinib) or BRAF/MEK Inhibitor Combinations approved by FDA* |
| Non-small cell lung cancer (NSCLC) | <i>EGFR</i> exon 19 deletions and exon 21 L858R alterations                                                                             | EGFR Tyrosine Kinase Inhibitors approved by FDA*                           |
| Solid Tumors                       | MSI-H                                                                                                                                   | KEYTRUDA® (pembrolizumab), JEMPERLI (dostarlimab-gxly)                     |
| Endometrial Carcinoma              | Not MSI-H                                                                                                                               | KEYTRUDA® (pembrolizumab) in combination with LENVIMA® (lenvatinib)        |

\*For the most current information about the therapeutic products in this group, go to:

[https://www.fda.gov/medical-devices/in-vitro-diagnostics/list-cleared-or-approved-companion-diagnostic-devices-in-vitro-and-imaging-tools#Group\\_Labeling](https://www.fda.gov/medical-devices/in-vitro-diagnostics/list-cleared-or-approved-companion-diagnostic-devices-in-vitro-and-imaging-tools#Group_Labeling)

MI Cancer Seek™ is a single-site assay performed at Caris Life Sciences, Phoenix, AZ.

### Contraindications:

There are no known contraindications.

### Warnings and Precautions:

Biopsy may pose a risk to the patient when archival tissue is not available for use with the assay. The patient's physician should determine whether the patient is a candidate for biopsy.

#### Limitations:

- For *in vitro* diagnostic use.
- CAUTION: Federal law restricts this device to sale by or on the order of a physician.
- The acceptable preparation method for MI Cancer Seek CDx specimens is FFPE. Other preparations have not been evaluated.
- The test is designed to report out somatic variants and is not intended to report germline variants.
- MI Cancer Seek requires a minimum tumor percentage of 20% for detection of alterations, with tumor content enrichment recommended for specimens with tumor percentage lower than 20%.
- Genomic findings other than those listed in the Companion Diagnostic Indications table are not prescriptive or conclusive for labeled use of any specific therapeutic product. Confirmation of tumor mutation status using an FDA-approved CDx test is needed for therapeutic use.
- A negative result does not rule out the presence of a mutation below the limits of detection of the assay.
- MI Cancer Seek is only approved for use with Caris Life Sciences pre-qualified Illumina NovaSeq 6000 instruments.
- The test is intended to be performed on specific serial number-controlled instruments by Caris Life Sciences.
- MI Cancer Seek does not report TMB for values lower than 3 mut/Mb as the accuracy of TMB values below 3 mut/Mb are unreliable.
- Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community.
- Based on the low positive percent agreement (PPA) in the accuracy study for ERBB2 copy number amplifications (CNAs) in breast cancer patients, this alteration may not be detected. Additional clinical investigation to confirm the presence of ERBB2 CNAs in the breast cancer patient's tumor with another FDA approved or cleared test is strongly recommended.
- Patients with breast cancer whose specimens have intermediate ERBB2 CNA status should be tested with another FDA approved or cleared test to ascertain ERBB2 CNA status in their tumor.
- Test may have reduced sensitivity and may yield false negative results in samples where necrotic tissue is >15%, melanin is >5%, fat cells are >10%.

#### Test Principle:

MI Cancer Seek is a single-site assay performed at Caris Life Sciences located at 4610 South 44th Place, Phoenix, AZ 85040. The test includes reagents, software, and procedures for testing of total nucleic acid (TNA) from formalin fixed paraffin embedded (FFPE) tumor tissue. The test uses a custom bait panel to measure all coding regions of the exome to detect and report SNVs and indels within 228 genes outlined in Table 2 across solid tumors and amplifications in ERBB2 in patients with breast cancer only. The test also detects MSI (determined from 3,210 genes) and whole exome based TMB in patients with solid tumors.

**Table 2. MI Cancer Seek Reportable Gene List for SNVs and indels**

|               |               |               |               |                 |               |               |                |                |                |                 |
|---------------|---------------|---------------|---------------|-----------------|---------------|---------------|----------------|----------------|----------------|-----------------|
| <i>ABL1</i>   | <i>BCL9</i>   | <i>CDKN2A</i> | <i>ESR1</i>   | <i>FLT1</i>     | <i>JAK3</i>   | <i>MET</i>    | <i>NRAS</i>    | <i>PPP2R2A</i> | <i>RUNX1</i>   | <i>STK11</i>    |
| <i>ACVR1</i>  | <i>BCOR</i>   | <i>CHEK1</i>  | <i>EZH2</i>   | <i>FLT3</i>     | <i>KDM5C</i>  | <i>MITF</i>   | <i>NSD1</i>    | <i>PRDM1</i>   | <i>SDHA</i>    | <i>SUFU</i>     |
| <i>AIP</i>    | <i>BLM</i>    | <i>CHEK2</i>  | <i>FANCA</i>  | <i>FOXA1</i>    | <i>KDM6A</i>  | <i>MLH1</i>   | <i>NSD2</i>    | <i>PRKACA</i>  | <i>SDHAF2</i>  | <i>TCF7L2</i>   |
| <i>AKT1</i>   | <i>BMPR1A</i> | <i>CIC</i>    | <i>FANCB</i>  | <i>FOXL2</i>    | <i>KDR</i>    | <i>MLH3</i>   | <i>NTHL1</i>   | <i>PRKAR1A</i> | <i>SDHB</i>    | <i>TERT</i>     |
| <i>AKT2</i>   | <i>BRAF</i>   | <i>CREBBP</i> | <i>FANCC</i>  | <i>FUBP1</i>    | <i>KEAP1</i>  | <i>MPL</i>    | <i>NTRK1</i>   | <i>PRKDC</i>   | <i>SDHC</i>    | <i>TET2</i>     |
| <i>AKT3</i>   | <i>BRCA1</i>  | <i>CSF1R</i>  | <i>FANCD2</i> | <i>GATA3</i>    | <i>KIT</i>    | <i>MRE11</i>  | <i>NTRK2</i>   | <i>PTCH1</i>   | <i>SDHD</i>    | <i>TMEM127</i>  |
| <i>ALK</i>    | <i>BRCA2</i>  | <i>CTCF</i>   | <i>FANCE</i>  | <i>GNA11</i>    | <i>KLF4</i>   | <i>MSH2</i>   | <i>NTRK3</i>   | <i>PTEN</i>    | <i>SETD2</i>   | <i>TNFAIP3</i>  |
| <i>AMER1</i>  | <i>BRIP1</i>  | <i>CTNNA1</i> | <i>FANCF</i>  | <i>GNA13</i>    | <i>KMT2A</i>  | <i>MSH3</i>   | <i>PALB2</i>   | <i>PTPN11</i>  | <i>SF3B1</i>   | <i>TNFRSF14</i> |
| <i>APC</i>    | <i>BTB</i>    | <i>CTNNA1</i> | <i>FANCG</i>  | <i>GNAQ</i>     | <i>KMT2C</i>  | <i>MSH6</i>   | <i>PBRM1</i>   | <i>RAC1</i>    | <i>SMAD2</i>   | <i>TP53</i>     |
| <i>AR</i>     | <i>CALR</i>   | <i>CXCR4</i>  | <i>FANCI</i>  | <i>GNAS</i>     | <i>KMT2D</i>  | <i>MTOR</i>   | <i>PDGFRA</i>  | <i>RAD50</i>   | <i>SMAD4</i>   | <i>TRAF7</i>    |
| <i>ARAF</i>   | <i>CARD11</i> | <i>CYLD</i>   | <i>FANCL</i>  | <i>H3F3A</i>    | <i>KRAS</i>   | <i>MUTYH</i>  | <i>PDGFRB</i>  | <i>RAD51B</i>  | <i>SMARCA4</i> | <i>TSC1</i>     |
| <i>ARID1A</i> | <i>CBFB</i>   | <i>DDR2</i>   | <i>FANCM</i>  | <i>H3F3B</i>    | <i>LZTR1</i>  | <i>MYC</i>    | <i>PIK3CA</i>  | <i>RAD51C</i>  | <i>SMARCB1</i> | <i>TSC2</i>     |
| <i>ARID2</i>  | <i>CCND1</i>  | <i>DICER1</i> | <i>FAS</i>    | <i>HIST1H3B</i> | <i>MAP2K1</i> | <i>MYCN</i>   | <i>PIK3CB</i>  | <i>RAD51D</i>  | <i>SMARCE1</i> | <i>U2AF1</i>    |
| <i>ASXL1</i>  | <i>CCND2</i>  | <i>DNMT3A</i> | <i>FAT1</i>   | <i>HNF1A</i>    | <i>MAP2K2</i> | <i>MYD88</i>  | <i>PIK3R1</i>  | <i>RAD54L</i>  | <i>SMO</i>     | <i>VHL</i>      |
| <i>ATM</i>    | <i>CCND3</i>  | <i>EGFR</i>   | <i>FBXW7</i>  | <i>HOXB13</i>   | <i>MAP2K4</i> | <i>NBN</i>    | <i>PIK3R2</i>  | <i>RAF1</i>    | <i>SOCS1</i>   | <i>WRN</i>      |
| <i>ATRX</i>   | <i>CD79B</i>  | <i>EP300</i>  | <i>FGFR1</i>  | <i>HRAS</i>     | <i>MAP3K1</i> | <i>NF1</i>    | <i>PIM1</i>    | <i>RASA1</i>   | <i>SOS1</i>    | <i>WT1</i>      |
| <i>AXIN2</i>  | <i>CDC73</i>  | <i>EPHA2</i>  | <i>FGFR2</i>  | <i>IDH1</i>     | <i>MAPK1</i>  | <i>NF2</i>    | <i>PMS2</i>    | <i>RB1</i>     | <i>SPEN</i>    | <i>XPO1</i>     |
| <i>B2M</i>    | <i>CDH1</i>   | <i>ERBB2</i>  | <i>FGFR3</i>  | <i>IDH2</i>     | <i>MAX</i>    | <i>NFE2L2</i> | <i>POLD1</i>   | <i>RET</i>     | <i>SPOP</i>    | <i>XRCC1</i>    |
| <i>BAP1</i>   | <i>CDK12</i>  | <i>ERBB3</i>  | <i>FGFR4</i>  | <i>IRF4</i>     | <i>MED12</i>  | <i>NFKBIA</i> | <i>POLE</i>    | <i>RHOA</i>    | <i>SRC</i>     |                 |
| <i>BARD1</i>  | <i>CDK4</i>   | <i>ERBB4</i>  | <i>FH</i>     | <i>JAK1</i>     | <i>MEF2B</i>  | <i>NOTCH1</i> | <i>POT1</i>    | <i>RNF43</i>   | <i>STAG2</i>   |                 |
| <i>BCL2</i>   | <i>CDKN1B</i> | <i>ERCC2</i>  | <i>FLCN</i>   | <i>JAK2</i>     | <i>MEN1</i>   | <i>NPM1</i>   | <i>PPP2R1A</i> | <i>ROS1</i>    | <i>STAT3</i>   |                 |

**FDA Evidence Levels:**

Genomic findings other than those listed in the Intended Use are not prescriptive or conclusive for labeled use of any specific therapeutic product. Test results should be interpreted in the context of pathological evaluation of tumors, treatment history, clinical findings, and other laboratory data. The test report includes genomic findings reported in the following levels (Table 3).

Table 3. Classification Criteria for FDA Evidence Levels

| LEVEL   | CRITERIA                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
|---------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Level 1 | Biomarker is FDA-approved as a companion diagnostic as part of MI Cancer Seek™                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Level 2 | Cancer alterations that are well-established with strong clinical evidence that the clinician must know according to professional consensus guidelines in the specific tumor type.                                                                                                                                                                                                                                                                                                                                    |
| Level 3 | Cancer alterations with potential clinical significance, e.g., biomarkers deemed useful for directing patients to a clinical trial or simply for informational purposes.                                                                                                                                                                                                                                                                                                                                              |
|         | <div><div>i. Clinical data such as case reports, single or several case series, or Phase I/II clinical trial data that support the utility of specific biomarker alteration to direct a patient to clinical trials, or</div><div>ii. Pre-clinical and/or <i>in vitro</i> studies provide structural analysis of the mutation, fusion, or isoform to confirm pathogenicity (tumor-promoting), sensitivity, or resistance through colony forming assays, growth inhibition or drug sensitivity assays, etc.</div></div> |



The Caris GPSai (Caris Genomic Probability Score – Artificial Intelligence) is a cancer-type similarity assessment which compares the characteristics of a patient's tumor against other tumors in the Caris database. Caris GPSai analyzes a tumor's molecular signature and provides the prevalence of that signature in the Caris Life Sciences genomic and transcriptomic database across 90 distinct cancer categories.

## Most Likely Tumor Type: Non-Small Cell Lung Carcinoma

73%

The predicted probability of each cancer type that can be detected by the product are shown below. The total probabilities of the top level cancer type (blue) total 100%. The subtypes (indented) sum to the total probability predicted for the top level cancer type. Discrepancies between expected sums and displayed sums may be exhibited due to rounding.

|   |                                                                |
|---|----------------------------------------------------------------|
| 0 | <b>Adrenal Cortical Carcinoma</b>                              |
| 1 | <b>Bladder/Urinary Tract</b>                                   |
| 0 | Urothelial Carcinoma                                           |
| 1 | Bladder Adenocarcinoma                                         |
| 0 | <b>Bowel</b>                                                   |
| 0 | Appendiceal Adenocarcinoma                                     |
| 0 | Colorectal Adenocarcinoma                                      |
| 0 | Small Bowel Carcinoma                                          |
| 1 | <b>Breast</b>                                                  |
| 0 | Metaplastic Breast Cancer                                      |
| 0 | Breast Invasive Lobular Carcinoma                              |
| 1 | Breast Invasive Ductal Carcinoma                               |
| 0 | <b>CNS/Brain</b>                                               |
| 0 | Diffuse Glioma                                                 |
| 0 | Meningioma                                                     |
| 0 | <b>Cervix/Uterine Carcinoma</b>                                |
| 0 | Uterine Serous Carcinoma/Uterine Papillary Serous Carcinoma    |
| 0 | Uterine Carcinosarcoma/Uterine Malignant Mixed Mullerian Tumor |
| 0 | Uterine Clear Cell Carcinoma                                   |
| 0 | Cervical Adenocarcinoma                                        |
| 0 | Uterine Endometrioid Carcinoma                                 |
| 0 | <b>Cutaneous Squamous Cell Carcinoma</b>                       |
| 5 | <b>Esophagus/Stomach</b>                                       |
| 0 | Esophageal Squamous Cell Carcinoma                             |
| 5 | Esophagogastric Adenocarcinoma                                 |
| 2 | Stomach Adenocarcinoma                                         |
| 1 | Adenocarcinoma of the Gastroesophageal Junction                |
| 1 | Esophageal Adenocarcinoma                                      |
| 0 | <b>Germ Cell Tumor</b>                                         |
| 0 | <b>Hematological</b>                                           |
| 0 | <b>Hepatocellular Carcinoma</b>                                |
| 0 | <b>Kidney</b>                                                  |
| 0 | Renal Cell Carcinoma                                           |
| 0 | Chromophobe Renal Cell Carcinoma                               |
| 0 | Papillary Renal Cell Carcinoma                                 |
| 0 | Renal Clear Cell Carcinoma                                     |
| 0 | Wilms Tumor                                                    |
| 0 | <b>Melanoma</b>                                                |
| 0 | <b>Mesothelioma</b>                                            |
| 1 | <b>Neuroendocrine Neoplasm</b>                                 |
| 1 | Well/Moderately-Differentiated Neuroendocrine Tumor            |
| 0 | Poorly-Differentiated Neuroendocrine Carcinoma                 |
| 0 | Large Cell Neuroendocrine Carcinoma                            |
| 0 | Small Cell Neuroendocrine Carcinoma                            |
| 0 | Paraganglioma/Pheochromocytoma                                 |
| 0 | Merkel Cell Carcinoma                                          |

|    |                                                            |
|----|------------------------------------------------------------|
| 73 | <b>Non-Small Cell Lung Carcinoma</b>                       |
| 0  | Lung Squamous Cell Carcinoma                               |
| 73 | Lung Adenocarcinoma                                        |
| 0  | <b>Orogenital Squamous Cell Carcinoma</b>                  |
| 0  | <b>Ovarian Epithelial Tumor</b>                            |
| 0  | Ovarian Carcinosarcoma/Malignant Mixed Mesodermal Tumor    |
| 0  | Serous Ovarian/Fallopian Tube/Peritoneal                   |
| 0  | Low-Grade Serous Ovarian/Fallopian Tube/Peritoneal Cancer  |
| 0  | High-Grade Serous Ovarian/Fallopian Tube/Peritoneal Cancer |
| 0  | Mucinous Ovarian Cancer                                    |
| 0  | Clear Cell Ovarian Cancer                                  |
| 0  | Endometrioid Ovarian Cancer                                |
| 19 | <b>Pancreatobiliary</b>                                    |
| 0  | Gallbladder Cancer                                         |
| 15 | Pancreatic Adenocarcinoma                                  |
| 4  | Cholangiocarcinoma                                         |
| 0  | <b>Peripheral Nervous System</b>                           |
| 0  | Schwannoma                                                 |
| 0  | Neuroblastoma                                              |
| 0  | Malignant Peripheral Nerve Sheath Tumor                    |
| 0  | <b>Prostate Adenocarcinoma</b>                             |
| 0  | <b>Salivary Gland Tumor</b>                                |
| 0  | <b>Sex Cord Stromal Tumor</b>                              |
| 0  | Granulosa Cell Tumor                                       |
| 0  | Sertoli-Leydig Cell Tumor                                  |
| 0  | <b>Soft Tissue/Bone</b>                                    |
| 0  | Gastrointestinal Stromal Tumor                             |
| 0  | Leiomyosarcoma                                             |
| 0  | Angiosarcoma                                               |
| 0  | Rhabdomyosarcoma                                           |
| 0  | Synovial Sarcoma                                           |
| 0  | Osteosarcoma                                               |
| 0  | Liposarcoma                                                |
| 0  | Chondrosarcoma                                             |
| 0  | Endometrial Stromal Sarcoma                                |
| 0  | Ewing Sarcoma                                              |
| 0  | <b>Thymic Carcinoma</b>                                    |
| 0  | <b>Thyroid</b>                                             |
| 0  | Medullary Thyroid Cancer                                   |
| 0  | Hurthle Cell Thyroid Cancer                                |
| 0  | Well-Differentiated Thyroid Cancer                         |
| 0  | Follicular Thyroid Cancer                                  |
| 0  | Papillary Thyroid Cancer                                   |
| 0  | Anaplastic Thyroid Cancer                                  |

PATIENT: WHITLEY, AMY (04-JUL-1955)

TN25-173573

PHYSICIAN: JANEESH SEKKATH-VEEDU, MD

## Gene Expression

| Gene    | Percentile in Cancer Type | Gene    | Percentile in Cancer Type | Gene     | Percentile in Cancer Type |
|---------|---------------------------|---------|---------------------------|----------|---------------------------|
| ADORA2A | 78                        | FOLR1   | 56                        | NTRK3    | 55                        |
| ALK     | 41                        | GPC3    | 72                        | PDCD1    | 80                        |
| ASCL1   | 89                        | HGF     | 93                        | PDCD1LG2 | 57                        |
| ATM     | 73                        | HRAS    | 32                        | PIK3CA   | 82                        |
| AURKA   | 1                         | IGF1R   | 76                        | POU3F2   | 84                        |
| BRAF    | 24                        | KDM1A   | 14                        | PRAME    | 48                        |
| BRCA1   | 9                         | KDR     | 60                        | PTEN     | 62                        |
| BRCA2   | 14                        | KEAP1   | 21                        | RB1      | 64                        |
| BRD4    | 74                        | KRAS    | 17                        | RET      | 70                        |
| CCND1   | 48                        | LAG3    | 68                        | ROR1     | 77                        |
| CCND2   | 54                        | MAGEA4  | 61                        | ROR2     | 24                        |
| CCNE1   | 7                         | MDM2    | 44                        | ROS1     | 72                        |
| CD274   | 52                        | MET     | 74                        | SLFN11   | 44                        |
| CD276   | 34                        | MKI67   | 24                        | SRC      | 80                        |
| CDKN2A  | 22                        | MSLN    | 82                        | SSTR2    | 68                        |
| CEACAM5 | 100                       | MTAP    | 21                        | SSTR3    | 62                        |
| CLDN18  | 98                        | MTOR    | 29                        | SSTR5    | 85                        |
| CLDN6   | 32                        | MUC1    | 77                        | STK11    | 43                        |
| CTLA4   | 68                        | MUC16   | 93                        | TACSTD2  | 84                        |
| DLL3    | 72                        | MYC     | 62                        | TGFB1    | 28                        |
| EGFR    | 22                        | NECTIN4 | 47                        | TOP1     | 42                        |
| EPHA2   | 96                        | NEUROD1 | 69                        | TP53     | 29                        |
| ERBB2   | 78                        | NF1     | 42                        | TSC1     | 58                        |
| ERBB3   | 94                        | NRAS    | 67                        | TSC2     | 48                        |
| FGFR1   | 34                        | NRG1    | 64                        | VEGFA    | 44                        |
| FGFR2   | 76                        | NTRK1   | 68                        | XPO1     | 29                        |
| FGFR3   | 46                        | NTRK2   | 16                        | YAP1     | 40                        |

### Gene Expression of Selected Genes by Whole Transcriptome Sequencing (WTS) Methods:


Gene expression is derived from whole transcriptome sequencing. Relative expression of genes are calculated as normalized values using Transcripts per Million Molecules or TPM. TPM is presented as a percentile derived by comparison to a distribution of Caris' internal cohort of the tumor-type profiled. Selected genes reported in this section were chosen based on their tumor-type specific relevance for matching to clinical trials, or tumor type subclassification.

**PATIENT:** WHITLEY, AMY (04-JUL-1955)

**TN25-173573**

**PHYSICIAN:** JANEESH SEKKATH-VEEDU, MD

## Karyotype



### Karyotyping using Copy Number Analysis by Whole Exome Sequencing (WES) Methods:

Whole exome sequencing in combination with interrogation of single nucleotide polymorphisms (SNPs) tiled throughout the genome, allows for the identification and visualization of cytogenetic aberrations.

Somatic structural variants like whole or partial chromosome duplications or deletions, are important for cancer development and progression, and may identify clinically actionable alterations.

PATIENT: WHITLEY, AMY (04-JUL-1955)

TN25-173573

PHYSICIAN: JANEESH SEKKATH-VEEDU, MD

## Mutational Analysis by Next-Generation Sequencing (NGS)

| TUMOR MUTATIONAL BURDEN |        |
|-------------------------|--------|
| Mutations / Megabase    | Result |
| 2                       | Low    |

### TMB

Tumor Mutational Burden (TMB) is defined as the number of somatic non-synonymous mutations per million bases of sequenced DNA in a tumor sample. Tumors with high TMB may increase the number of neoantigens which is hypothesized to increase T-cell reactivity and potential for response to immune checkpoint inhibitors. TMB analysis was performed based on next generation sequencing analysis of genomic DNA isolated from a tumor sample.

| MICROSATELLITE INSTABILITY ANALYSIS |        |
|-------------------------------------|--------|
| Test                                | Result |
| MSI                                 | Stable |

### MSI

Microsatellite instability (MSI) status is a measure of the number of somatic mutations within short, repeated sequences of DNA (microsatellites). MSI-High status can indicate that the tumor has a defect in mismatch repair (MMR) abrogating the ability to correct mistakes during DNA replication. Tumors with MSI-high status may increase the number of neoantigens which is hypothesized to increase T-cell reactivity and potential for response to immune checkpoint inhibitors. Tumor-only microsatellite instability status by NGS (MSI-NGS) is measured by the direct analysis of known microsatellite regions sequenced in the CMI NGS panel.

| GENOMIC LOSS OF HETEROZYGOSITY       |                                                                                     |
|--------------------------------------|-------------------------------------------------------------------------------------|
| Test                                 | Result                                                                              |
| Genomic Loss of Heterozygosity (LOH) | Low - 2% of tested genomic segments exhibited LOH (assay threshold is $\geq 16\%$ ) |

### LOH

To calculate genomic loss-of-heterozygosity (LOH), the 22 autosomal chromosomes are split into 552 segments and the LOH of single nucleotide polymorphisms (SNPs) within each segment is calculated. Caris WES data consist of approximately 250k SNPs spread across the genome. SNP alleles with frequencies skewed towards 0 or 100% indicate LOH (heterozygous SNP alleles have a frequency of 50%). The final call of genomic LOH is based on the percentage of all 552 segments with observed LOH.

*Additional Next-Generation Sequencing results continued on the next page. >*

**PATIENT:** WHITLEY, AMY J (04-JUL-1955)

**TN25-173573**

**PHYSICIAN:** JANEESH SEKKATH-VEEDU, MD

## Mutational Analysis by Next-Generation Sequencing (NGS)

| Gene | Analyte   | Variant Interpretation    | Protein Alteration | Exon | DNA Alteration | Variant Frequency % | Transcript ID |
|------|-----------|---------------------------|--------------------|------|----------------|---------------------|---------------|
| CDH1 | DNA-Tumor | Likely Pathogenic Variant | p.D400N            | 9    | c.1198G>A      | 13                  | NM_004360.4   |

**Interpretation:** A mutation, D400N, was detected in the CDH1 (E-cadherin) gene. This mutation is presumed to be pathogenic because it affects a highly-conserved position in the E-cadherin protein, and this mutation along with other similar mutations have been reported in both mutation database and at Caris for several tumors commonly associated with loss of CDH1 function (diffuse gastric cancer, lobular breast cancer). Pathogenic germline mutations in the CDH1 gene are causal for hereditary diffuse gastric cancer.

This gene is a classical cadherin from the cadherin superfamily. The encoded protein is a calcium dependent cell-cell adhesion glycoprotein comprised of five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. The protein plays a major role in epithelial architecture, cell adhesion and cell invasion. Mutations in this gene are correlated with gastric, breast, colorectal, thyroid and ovarian cancer. Loss of function is thought to contribute to progression in cancer by increasing proliferation, invasion, and/or metastasis. The ectodomain of this protein mediates bacterial adhesion to mammalian cells and the cytoplasmic domain is required for internalization.

| Gene | Analyte   | Variant Interpretation | Protein Alteration | Exon | DNA Alteration | Variant Frequency % | Transcript ID |
|------|-----------|------------------------|--------------------|------|----------------|---------------------|---------------|
| KRAS | DNA-Tumor | Pathogenic Variant     | p.G12D             | 2    | c.35G>A        | 18                  | NM_004985.4   |

**Interpretation:** A pathogenic codon 12 mutation was detected in KRAS.

KRAS or V-Ki-ras2 Kirsten rat sarcoma viral oncogene homolog encodes a signaling intermediate involved in many signaling cascades including the EGFR pathway. KRAS somatic mutations have been found in pancreatic (57%), colon (35%), lung (16%), biliary tract (28%), and endometrial (15%) cancers. Several germline mutations of KRAS (V14I, T58I, and D153V amino acid substitutions) are associated with Noonan syndrome.

| Gene   | Analyte   | Variant Interpretation            | Protein Alteration | Exon | DNA Alteration | Variant Frequency % | Transcript ID |
|--------|-----------|-----------------------------------|--------------------|------|----------------|---------------------|---------------|
| NFE2L2 | DNA-Tumor | Variant of Uncertain Significance | p.Y576H            | 5    | c.1726T>C      | 50                  | NM_006164.4   |

**Interpretation:** A variant with no known clinical or functional significance was detected in NFE2L2.

This gene encodes a transcription factor which is a member of a small family of basic leucine zipper (bZIP) proteins. The encoded transcription factor regulates genes which contain antioxidant response elements (ARE) in their promoters; many of these genes encode proteins involved in response to injury and inflammation which includes the production of free radicals.

*Additional Next-Generation Sequencing results continued on the next page. >*

**PATIENT:** WHITLEY, AMY J (04-JUL-1955)

**TN25-173573**

**PHYSICIAN:** JANEESH SEKKATH-VEEDU, MD



## Mutational Analysis by Next-Generation Sequencing (NGS)

### GENES TESTED WITH INDETERMINATE\* RESULTS BY TUMOR DNA SEQUENCING

|         |       |       |       |       |         |
|---------|-------|-------|-------|-------|---------|
| ATP6AP2 | CUL3  | MDH2  | PRDM6 | RASA1 | SMARCA2 |
| COL2A1  | DGCR8 | PLCB4 | PRKD1 | REST  | TRIM28  |

\* Genes in this table were ruled indeterminate due to low coverage for some or all exons.

For a complete list of genes tested, visit [www.CarisMolecularIntelligence.com/profilemenu](http://www.CarisMolecularIntelligence.com/profilemenu).

### NGS Methods

Direct sequence analysis was performed on genomic DNA isolated from a micro-dissected tumor sample using Illumina NovaSeq 6000 sequencers. A hybrid pull-down panel of baits was used to enrich more than 700 clinically relevant genes along with > 20,000 other genes. Sequence data is analyzed using a customized bioinformatics pipeline to detect sequencing variants, copy number alterations (amplifications and deletions) indels and HLA genotypes. In addition, genomic signatures for tumor mutational burden (TMB), microsatellite instability (MSI), genomic loss-of-heterozygosity (LOH) or HRD-Genomic Scar Score (HRD-GSS), and homologous recombination deficiency (HRD) are reported when applicable. For a complete list of what is covered by the assay, and genes with partial coverage, please contact Caris Customer Support.

## Copy Number Alterations by Next-Generation Sequencing (NGS)

### GENES TESTED WITH INTERMEDIATE CNA RESULTS

|          |       |       |      |        |  |
|----------|-------|-------|------|--------|--|
| APOBEC3B | CSF1R | FGFR2 | JAK2 | PDGFRB |  |
|----------|-------|-------|------|--------|--|

### GENES WITH INDETERMINATE CNA RESULTS

|        |      |      |       |        |  |
|--------|------|------|-------|--------|--|
| CDKN1B | ETV6 | KRAS | RAD52 | ZNF384 |  |
|--------|------|------|-------|--------|--|

### CNA Methods

The copy number alteration (CNA) of each exon is determined by a calculation using the average sequencing depth of the sample along with the sequencing depth of each exon and comparing this calculated result to a pre-calibrated value. A complete list of genes for reporting copy number alterations, including amplifications and deletions, is available upon request.

## Gene Fusion and Transcript Variant Detection by RNA Sequencing

### Whole Transcriptome Sequencing (WTS) Methods

Gene fusion and variant transcript detection were performed on RNA isolated from a tumor sample using next generation sequencing. The assay also detects fusions occurring at known and novel breakpoints within genes. The genes included in this report represent the subset of genes associated with cancer. The complete list of unclassified alterations is available by request.

## Protein Expression by Immunohistochemistry (IHC)

| Biomarker | Patient Tumor                         |                  |          | Thresholds                                           |
|-----------|---------------------------------------|------------------|----------|------------------------------------------------------|
|           | Staining Intensity<br>(0, 1+, 2+, 3+) | Percent of cells | Result   | Conditions for a Positive Result:                    |
| ALK       | 0                                     | 100              | Negative | Intensity $\geq 3+$ and $\geq 1\%$ of cells stained  |
| MET       | 3 +                                   | 35               | Positive | Intensity $\geq 3+$ and $\geq 25\%$ of cells stained |

| PD-L1 TUMOR CELL STAINING |                                       |                  |          |                                                      |
|---------------------------|---------------------------------------|------------------|----------|------------------------------------------------------|
| Biomarker                 | Patient Tumor                         |                  |          | Thresholds                                           |
|                           | Staining Intensity<br>(0, 1+, 2+, 3+) | Percent of cells | Result   | Conditions for a Positive Result:                    |
| PD-L1 (28-8)              | 0                                     | 100%             | Negative | Intensity $\geq 1+$ and $\geq 1\%$ of cells stained  |
| PD-L1 (SP142)             | 0                                     | 100%             | Negative | Intensity $\geq 1+$ and $\geq 50\%$ of cells stained |
| PD-L1 (SP263)             | 0                                     | 100%             | Negative | Intensity $\geq 1+$ and $\geq 1\%$ of cells stained  |

PD-L1 (28-8): Scoring was based on percentage of viable tumor cells showing partial or complete membrane staining at any intensity.

PD-L1 (SP142): TC scoring was based on the presence of discernible PD-L1 membrane staining of any intensity in  $\geq 50\%$  of viable tumor cells.

PD-L1 (SP263): Scoring was based on percentage of viable tumor cells showing partial or complete membrane staining at any intensity.

| PD-L1 TUMOR PROPORTION SCORE (TPS) |          |     |                |
|------------------------------------|----------|-----|----------------|
| Biomarker                          | Result   | TPS | Threshold      |
| PD-L1 (22c3)                       | Negative | 0%  | TPS $\geq 1\%$ |

PD-L1 22c3: Scoring was based on the percentage of viable tumor cells showing partial or complete membrane staining. There are three categories of PD-L1 expression defined by the PD-L1 22c3 IHC pharmDx NSCLC interpretation guide: TPS  $< 1\%$  (negative), TPS  $\geq 1\%$  and TPS  $\geq 50\%$ .

| Her-2 IHC: Biopsy   |                                                 |
|---------------------|-------------------------------------------------|
| Final Score         | Threshold for Positive Result                   |
| Negative   Score 1+ | Intensity =3+ and $\geq 5$ tumor cells staining |

| PD-L1 IMMUNE CELL (IC) SCORE |          |    |             |
|------------------------------|----------|----|-------------|
| Biomarker                    | Result   | IC | Threshold   |
| PD-L1 (SP142)                | Negative | 5% | $\geq 10\%$ |

PD-L1 (SP142): IC scoring was based on discernible PD-L1 staining of any intensity in tumor-infiltrating immune cells covering  $\geq 10\%$  of tumor area occupied by tumor cells, associated intratumoral or contiguous peritumoral stroma.

Clones used: ALK (D5F3), ERBB2 (Her2/Neu) (4B5), MET (SP44) RxDx Assay, PD-L1 (22c3), PD-L1 (28-8), PD-L1 (SP142), PD-L1 (SP263).



Electronic Signature  
Caleb Cheng, MD, MBA  
05/11/2025

Additional IHC results continued on the next page. >

PATIENT: WHITLEY, AMY J (04-JUL-1955)

TN25-173573

PHYSICIAN: JANEESH SEKKATH-VEEDU, MD

## Protein Expression by Immunohistochemistry (IHC)

### IHC Methods

The Laboratory Developed Tests (LDT) immunohistochemistry (IHC) assays were developed and their performance characteristics determined by Caris Life Sciences. These tests have not been cleared or approved by the US Food and Drug Administration. The FDA has determined that such clearance or approval is not currently necessary. Interpretations of all immunohistochemistry (IHC) assays were performed manually or with the assistance of an AI-based image analysis tool by a board certified pathologist using a microscope and/or digital whole slide image(s).

The following IHC assays were performed using FDA-approved companion diagnostic or FDA-cleared tests consistent with the manufacturer's instructions: ALK (VENTANA ALK (D5F3) CDx Assay, Ventana), ER (CONFIRM anti-Estrogen Receptor (ER) (SP1), Ventana), FOLR1 (VENTANA FOLR1-2.1 RxDx, Ventana), CLDN18 (VENTANA, 43-14A RxDx Assay, Gastric/GEJ), PR (CONFIRM anti-Progesterone Receptor (PR) (1E2), Ventana), HER2/neu (PATHWAY anti-HER-2/neu (4B5), Ventana), Ki-67 (MIB-1 pharmDx, Dako), MAGE-A4 1F9 (pharmDx, Dako), PD-L1 22c3 (pharmDx, Dako), PD-L1 SP142 (VENTANA, non-small cell lung cancer), PD-L1 28-8 (pharmDx, Dako, Gastric/GEJ, non-small cell lung cancer), PD-L1 SP263 (Ventana, non-small cell lung cancer), and Mismatch Repair (MMR) proteins (MLH1, MSH2, MSH6, and PMS2; VENTANA MMR RxDx Panel, Ventana).

HER2 results and interpretation follow the ASCO/CAP scoring criteria. Bartley, A.N., J.A. Ajani, et al. (2016). "HER2 testing and clinical decision making in gastroesophageal adenocarcinoma: guideline from the College of American Pathologists, American Society for Clinical Pathology, and the American Society of Clinical Oncology". J Clin Oncol. 35(4):446-464.

Professional Component Performed: -- Location 18, Irvine, CA. Caleb Cheng, MD, MBA, Lab Director.

## References

| #  | Drug                                                                        | Biomarker | Reference                                                                                                                                                                                                                                                                                                                                                                                                          |
|----|-----------------------------------------------------------------------------|-----------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1  | alectinib                                                                   | ALK       | Camidge, D.R., A.T. Shaw, et al. (2019). "Updated Efficacy and Safety Data and Impact of the EML4-ALK Fusion Variant on the Efficacy of Alectinib in Untreated ALK-Positive Advanced Non-Small Cell Lung Cancer in the Global Phase III ALEX Study." J Thorac Oncol 14(7): 1233-1243. <a href="#">View Citation Online</a>                                                                                         |
| 2  | alectinib                                                                   | ALK       | Gadgeel, S., D.R. Camidge, et al. (2018). "Alectinib versus crizotinib in treatment-naïve anaplastic lymphoma kinase-positive (ALK+) non-small-cell lung cancer: CNS efficacy results from the ALEX study." Ann Oncol 29 (11): 2214-2222. <a href="#">View Citation Online</a>                                                                                                                                     |
| 3  | alectinib, brigatinib, ceritinib, crizotinib, lorlatinib                    | ALK       | Lindeman, N.I., Y. Yatabe, et al. (2018). "Updated Molecular Testing Guideline for the Selection of Lung Cancer Patients for Treatment With Targeted Tyrosine Kinase Inhibitors Guideline From the College of American Pathologists, the International Association for the Study of Lung Cancer, and the Association for Molecular Pathology." J Thorac Oncol 13(3): 323-358. <a href="#">View Citation Online</a> |
| 4  | alectinib, brigatinib, ceritinib, crizotinib, lorlatinib                    | ALK       | National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology. Non-Small Cell Lung Cancer Version 1.2020                                                                                                                                                                                                                                                                                    |
| 5  | brigatinib                                                                  | ALK       | Camidge, D.R., S. Popat, et al. (2018). "Brigatinib versus Crizotinib in ALK-Positive Non-Small-Cell Lung Cancer." N Engl J Med 379(21): 2027-2039. <a href="#">View Citation Online</a>                                                                                                                                                                                                                           |
| 6  | brigatinib                                                                  | ALK       | Lin, J.L., G.J. Riely, et al. (2018). "Brigatinib in Patients with Alectinib-refractory ALK-positive NSCLC." J Thorac Onc 13(10): 1530-1538. <a href="#">View Citation Online</a>                                                                                                                                                                                                                                  |
| 7  | brigatinib                                                                  | ALK       | Reckamp, K., J. Lee, et al. (2019). "Comparative efficacy of brigatinib versus ceritinib and alectinib in patients with crizotinib-refractory anaplastic lymphoma kinase-positive non-small cell lung cancer." Curr Med Res Opin 35(4):569-576. <a href="#">View Citation Online</a>                                                                                                                               |
| 8  | brigatinib, crizotinib                                                      | ALK       | Thorne-Nuzzo, T., P. Towne, et al. (2017). "A Sensitive ALK Immunohistochemistry Companion Diagnostic Test Identifies Patients Eligible for Treatment with Crizotinib." J Thorac Oncol 12(5): 804-813 <a href="#">View Citation Online</a>                                                                                                                                                                         |
| 9  | ceritinib, lorlatinib                                                       | ALK       | Shaw, A.T., E. Felip, et al. (2017). "Ceritinib versus chemotherapy in patients with ALK-rearranged non-small-cell lung cancer previously given chemotherapy and crizotinib (ASCEND-5): a randomised, controlled, open-label, phase 3 trial." Lancet Oncol 18 (7):874-886. <a href="#">View Citation Online</a>                                                                                                    |
| 10 | ceritinib, lorlatinib                                                       | ALK       | Solomon, B. J., A. T. Shaw, et al. (2018). "Lorlatinib in patients with ALK-positive non-small cell lung cancer: results from a global phase 2 study." Lancet Oncol 19:1654-1667. <a href="#">View Citation Online</a>                                                                                                                                                                                             |
| 11 | ceritinib, lorlatinib                                                       | ALK       | Soria, J.C., de Castro G., et al. (2017). "First-line ceritinib versus platinum-based chemotherapy in advanced ALK-rearranged non-small-cell lung cancer (ASCEND-4): a randomised, open-label, phase 3 study." Lancet 389: 917-929. <a href="#">View Citation Online</a>                                                                                                                                           |
| 12 | crizotinib                                                                  | ALK       | Solomon, B. J., T. S. Mok, et al. (2018). "Final overall survival analysis from a study comparing first-line crizotinib versus chemotherapy in ALK-mutation positive Non-small-cell-lung cancer." J Clin Oncol 36: 2251-2258. <a href="#">View Citation Online</a>                                                                                                                                                 |
| 13 | crizotinib                                                                  | ALK       | van der Wekken, A. J., H.J.M. Groen, et al. (2017). "Dichotomous ALK IHC is a better predictor for ALK inhibition outcome than traditional ALK FISH in advanced Non-small cell lung cancer." Clin Cancer Res 23(15): 4251-4258. <a href="#">View Citation Online</a>                                                                                                                                               |
| 14 | dabrafenib, dabrafenib + trametinib, encorafenib + binimetinib, vemurafenib | BRAF      | Hyman, D.H., J. Baselga, et al. (2015). "Vemurafenib in Multiple Nonmelanoma Cancers with BRAF V600 Mutations." NEJM 373(8):726-736. <a href="#">View Citation Online</a>                                                                                                                                                                                                                                          |
| 15 | dabrafenib, dabrafenib + trametinib, encorafenib + binimetinib, vemurafenib | BRAF      | Planchard, D., B.E. Johnson, et al. (2016). "An open-label phase II trial of dabrafenib (D) in combination with trametinib (T) in patients (pts) with previously treated BRAF V600E-mutant advanced non-small cell lung cancer (NSCLC; BR113928)." J Clin Oncol 34: 15_suppl, 107-107. <a href="#">View Citation Online</a>                                                                                        |
| 16 | dabrafenib, dabrafenib + trametinib, encorafenib + binimetinib, vemurafenib | BRAF      | Planchard, D., B.E. Johnson, et al. (2017). "Dabrafenib plus trametinib in patients with previously untreated BRAFV600E-mutant metastatic non-small-cell lung cancer: an open-label, phase 2 trial." Lancet Oncol 18(1):1307-1316. <a href="#">View Citation Online</a>                                                                                                                                            |

## References

| #  | Drug                                                                        | Biomarker | Reference                                                                                                                                                                                                                                                                                                                                                                           |
|----|-----------------------------------------------------------------------------|-----------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 17 | dabrafenib, dabrafenib + trametinib, encorafenib + binimetinib, vemurafenib | BRAF      | Riely, G. J., B. E. Johnson, et al. (2023). "Phase II, Open-Label Study of Encorafenib Plus Binimetinib in Patients With BRAFV600-Mutant Metastatic Non-Small-Cell Lung Cancer." J Clin Oncol. 41 (21): 3700-3711 <a href="#">View Citation Online</a>                                                                                                                              |
| 18 | dabrafenib, dabrafenib + trametinib, encorafenib + binimetinib, vemurafenib | BRAF      | Subbiah, V., D.M. Hyman, et al. (2019). "Efficacy of Vemurafenib in Patients With Non-Small-Cell Lung Cancer With BRAF V600 Mutation: An Open-Label, Single-Arm Cohort of the Histology-Independent VE-BASKET Study." JCO Precis Oncol 2019; 3, 1-9. <a href="#">View Citation Online</a>                                                                                           |
| 19 | afatinib, dacomitinib, osimertinib                                          | EGFR      | Sequist, L.V., M. Schuler, et al. (2013). "Phase III Study of Afatinib or Cisplatin Plus Pemetrexed in Patients with Metastatic Lung Adenocarcinoma With EGFR Mutations." J Clin Oncol ahead of print July 1, 2013, doi: 10.1200/JCO.2012.44.2806 <a href="#">View Citation Online</a>                                                                                              |
| 20 | afatinib, dacomitinib, osimertinib                                          | EGFR      | Wu, Y-L, T.S. Mok, et al (2017). "Dacomitinib versus gefitinib as first-line treatment for patients with EGFR-mutation-positive non-small-cell lung cancer (ARCHER 1050): a randomized, open-label, phase III trial." Lancet Oncol. 2017;18(11):1454-1466. <a href="#">View Citation Online</a>                                                                                     |
| 21 | afatinib, dacomitinib, osimertinib                                          | EGFR      | Yang, J.C., et al. (2013). "Symptom control and quality of life in LUX-Lung 3: a phase III study of afatinib or cisplatin/pemetrexed in patients with advanced lung adenocarcinoma with EGFR mutations." J Clin Oncol 31:3342-3350. <a href="#">View Citation Online</a>                                                                                                            |
| 22 | afatinib, dacomitinib, osimertinib                                          | EGFR      | Yang, J.C., et al. (2015). "Clinical activity of afatinib in patients with advanced non-small-cell lung cancer harbouring uncommon EGFR mutations: a combined post-hoc analysis of LUX-Lung 2, LUX-Lung 3, and LUX-Lung 6." Lancet Oncol; (7):830-838. <a href="#">View Citation Online</a>                                                                                         |
| 23 | erlotinib, gefitinib                                                        | EGFR      | Brugger, W., F. Cappuzzo, et. al. (2011). "Prospective molecular marker analyses of EGFR and KRAS from a randomized, placebo-controlled study of erlotinib maintenance therapy in advanced non-small-cell lung cancer." J. Clin. Oncol. 29:4113-4120. <a href="#">View Citation Online</a>                                                                                          |
| 24 | erlotinib, gefitinib                                                        | EGFR      | Fukuoka, M., T.S.K. Mok, et. al. (2011). "Biomarker analyses and final overall survival results from a phase III, randomized, open-label, first-line study of gefitinib versus carboplatin/paclitaxel in clinically selected patients with advanced non-small-cell lung cancer in Asia (IPASS). J. Clin. Oncol. DOI: 10.1200/JCO.2010.33.4235. <a href="#">View Citation Online</a> |
| 25 | erlotinib, gefitinib                                                        | EGFR      | Keedy, V.L., G. Gianconne, et. al. (2011). "American Society of Clinical Oncology Provisional Clinical Opinion: epidermal growth factor receptor (EGFR) mutation testing for patients with advanced non-small cell lung cancer considering first-line EGFR tyrosine kinase inhibitor therapy." J. Clin. Oncol. 29(15):2121-2127. <a href="#">View Citation Online</a>               |
| 26 | erlotinib, gefitinib                                                        | EGFR      | Maemondo, M., T. Nukiwa, et. al. (2010). "Gefitinib or chemotherapy for non-small cell lung cancer with mutated EGFR." N. Engl. J. Med. 362:2380-8. <a href="#">View Citation Online</a>                                                                                                                                                                                            |
| 27 | capmatinib, tepotinib                                                       | MET       | Paik, P. K., X. Le, et al. (2020). "Tepotinib in patients with MET exon 14 (METex14) skipping advanced NSCLC: Updated efficacy results from VISION Cohort A." IASLC 2020 World Conference on Lung Cancer (Abstract: 1361) <a href="#">View Citation Online</a>                                                                                                                      |
| 28 | capmatinib, tepotinib                                                       | MET       | Paik, P.K., X. Le, et al. (2020). "Tepotinib in Non-Small Cell Lung Cancer with MET Exon 14 Skipping Mutations." N Engl J Med 383(10): 931-943. <a href="#">View Citation Online</a>                                                                                                                                                                                                |
| 29 | capmatinib, tepotinib                                                       | MET       | Wolf, J., R.S. Heist, et al. (2019). "Capmatinib (INC280) in METΔex14-mutated advanced non-small cell lung cancer (NSCLC): efficacy data from the Phase 2 GEOMETRY mono-1 study." J Clin Oncol 37, (suppl; abstr 9004). <a href="#">View Citation Online</a>                                                                                                                        |
| 30 | crizotinib                                                                  | MET       | Paik, P.K., M. Ladanyi, et al. (2015). "Response to MET inhibitors in patients with stage IV lung adenocarcinomas harboring MET mutations causing exon 14 skipping". Cancer Discov. Published online May 13, 2015. <a href="#">View Citation Online</a>                                                                                                                             |
| 31 | crizotinib                                                                  | MET       | Wang, S.X.Y., J.W. Neal, et al. (2019). "Case series of MET exon 14 skipping mutation-positive non-small-cell lung cancers with response to crizotinib and cabozantinib." Anticancer Drugs 30(5):537-541. <a href="#">View Citation Online</a>                                                                                                                                      |

## References

| #  | Drug                       | Biomarker | Reference                                                                                                                                                                                                                                                                                                                      |
|----|----------------------------|-----------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 32 | telisotuzumab vedotin      | MET       | Camidge, D., S. Lu et al. "Telisotuzumab Vedotin Monotherapy in Patients With Previously Treated c-Met Protein-Overexpressing Advanced Nonsquamous EGFR-Wildtype Non-Small Cell Lung Cancer in the Phase II LUMINOSITY Trial." J Clin Oncol [published online ahead of print, 2024 Jun 6] <a href="#">View Citation Online</a> |
| 33 | pralsetinib                | RET       | Gainor J.F., V. Subbiah, et al. (2020). "Registrational dataset from the phase I/II ARROW trial of pralsetinib (BLU-667) in patients (pts) with advanced RET fusion+ non-small cell lung cancer (NSCLC)." J Clin Oncol. 38(suppl):9515. <a href="#">View Citation Online</a>                                                   |
| 34 | pralsetinib, selpercatinib | RET       | National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology. Non-Small Cell Lung Cancer Version 6.2020 <a href="#">View Citation Online</a>                                                                                                                                                           |
| 35 | selpercatinib              | RET       | Drilon, A., V. Subbiah, et al. (2020). "Efficacy of Selpercanib in RET Fusion-Positive Non-Small-Cell Lung Cancer." N Engl J Med. 383 (9): 813-824. <a href="#">View Citation Online</a>                                                                                                                                       |
| 36 | ceritinib, lorlatinib      | ROS1      | Lim SM, BC Cho, et al. (2017). "Open-Label, Multicenter, Phase II Study of Ceritinib in Patients With Non-Small-Cell Lung Cancer Harboring ROS1 Rearrangement". J Clin Oncol. May 18;JCO2016713701. <a href="#">View Citation Online</a>                                                                                       |
| 37 | ceritinib, lorlatinib      | ROS1      | Shaw, A.T., S.-H. I. Ou, et al. (2019). "Lorlatinib in advanced ROS-1 positive non-small-cell lung cancer: a multicentre, open-label, single-arm, phase 1-2 trial." Lancet Oncol pii: S1470-2045(19)30655-2. <a href="#">View Citation Online</a>                                                                              |
| 38 | crizotinib                 | ROS1      | Shaw, A.T., S.-H. I. Ou, et al. (2019). "Crizotinib in ROS1-rearranged advanced non-small-cell lung cancer (NSCLC): updated results, including overall survival, from PROFILE 1001." Ann Oncol 30(7): 1121-1126. <a href="#">View Citation Online</a>                                                                          |
| 39 | entrectinib, repotrectinib | ROS1      | Cho, B.C., A. Drilon, et al. (2023). "Repotrectinib in patients with ROS1 fusion-positive (ROS1+) NSCLC: Update from the pivotal phase 1/2 TRIDENT-1 trial." Presented at: 2023 World Conference on Lung Cancer. Abstract OA03.06. <a href="#">View Citation Online</a>                                                        |
| 40 | entrectinib, repotrectinib | ROS1      | Demetri, G.D., R.D., Doebele, et al. (2018). "Efficacy and safety of entrectinib in patients with NTRK fusion-positive tumors: pooled analysis of STARTRK-2, STARTRK-1 and ALKA-372-001. Presented at: 2018 ESMO Congress; October 19-23, 2018; Munich, Germany. Abstract LBA17. <a href="#">View Citation Online</a>          |
| 41 | entrectinib, repotrectinib | ROS1      | Desai AV, Brodeur GM, Foster J, et al. Phase I study of entrectinib (RXDX-101), a TRK, ROS1, and ALK inhibitor, in children, adolescents, and young adults with recurrent or refractory solid tumors. J Clin Oncol. 2018;36 (suppl;abstr 10536). doi: 10.1200/JCO.2018.36.15_suppl.10536. <a href="#">View Citation Online</a> |