
**Molecular Tumor Board Recommendation
May 23, 2025**

Dear Dr. Sekkath-Veedu:

The Markey Cancer Center Molecular Tumor Board met on May 20, 2025 and reviewed your patient, Amy Whitley (019092113; mucinous adenocarcinoma of the lower lobe of the lung). After reviewing the genomic report (Caris report dated May 11, 2025) and clinical characteristics of the case, the MTB recommends consideration of the following:

1. If progression, consider the following clinical trials due to the MET 3+, 35% result:
 - a. NCT04270591 (Ph 1b/2): Assessment of Anti-tumor and Safety in Glutemetinib in Patients With c-MET-positive Non-Small Cell Lung Cancer
 - b. NCT03175224 (Ph 1/2): APL-101 Study of Subjects With NSCLC With c-Met EXON 14 Skip Mutations and c-Met Dysregulation Advanced Solid Tumors (SPARTA)
 - c. NCT04693468 (Ph 1b): Talazoparib and Palbociclib, Axitinib, or Crizotinib for the Treatment of Advanced or Metastatic Solid Tumors, TalaCom Trial
 - d. NCT06083857 (Ph 1/2): Amivantamab and Tepotinib for the Treatment of Locally Advanced or Metastatic MET-Altered Non-Small Cell Lung Cancer
2. If progression, consider the following clinical trials targeting the KRAS mutation (G12D):
 - a. NCT06586515 (1a/b): MOONRAY-01, A Study of LY3962673 in Participants With KRAS G12D-Mutant Solid Tumors (MOONRAY-01)
 - b. NCT05382559 (Ph 1): A Study of ASP3082 in Adults With Advanced Solid Tumors
 - c. NCT06040541 (Ph 1/1b): Study of RMC-9805 in Participants With KRAS G12D-Mutant Solid Tumors
 - d. NCT05585320 (Ph 1/2a): A Phase 1/2a Study of IMM-1-104 in Participants With Advanced or Metastatic Solid Tumors
 - e. NCT06385925 (Ph 1/2): A Study of TSN1611 Treating Patients with Advanced Solid Tumors Harboring KRAS G12D Mutation
3. If progression beyond standard of care therapy, consider off-label telisotuzumab vedotin due to the MET 3+, 35% result. This is a category 2 evidence level recommendation. Please note, in a phase II trial of patients with nonsquamous EGFR-wildtype NSCLC, telisotuzumab vedotin treatment resulted in an ORR of 22.9%, median PFS of 6.0 months, and median OS of 14.2 months in patients with intermediate cMET overexpression after receiving prior systemic treatment (PMID: 38843488).
 - a. Reference: Camidge DR, Bar J, Horinouchi H, et al. Telisotuzumab Vedotin Monotherapy in Patients With Previously Treated c-Met Protein-Overexpressing Advanced Nonsquamous EGFR-Wildtype Non-Small Cell Lung Cancer in the Phase II LUMINOSITY Trial. J Clin Oncol. 2024;42(25):3000-3011. doi:10.1200/JCO.24.00720

Thank-you for submitting this case to the Markey Cancer Center MTB.

Signature: _____ Date: _____

Signature: _____ Date: _____

Signature: _____ Date: _____

Level of Evidence	Interpretation	Supporting Evidence
1	FDA-labeled indication or NCCN-recommended; substantial evidence supporting use (standard of care)	Meta-analysis, RCTs, etc. OncoKB level 1
2	Off-label use; substantial evidence supporting use	Phase II studies and above OncoKB level 2 and above
3	Off-label use; low to moderate evidence supporting use; MTB recommends, clinical trial preferred	Phase I, case studies and reports OncoKB level 3 and below
4	Off-label; no evidence supporting use; enrollment in clinical trial recommended; report experience to MTB if used outside clinical trial	Little to no evidence (pre-clinical, animal/cellular models), mechanistically plausible