



Home Tube Feeding & PEG Care Handbook — Chennai Edition

Why Safe Tube Feeding Matters

Feeding tubes such as **Ryle's (NG)** or **PEG/PEJ** help patients with swallowing difficulties, cancer, stroke, or neurological illness maintain proper nutrition at home. But poor technique can lead to **aspiration pneumonia, tube blockages, infections, and re-hospitalisation**. This handbook helps families prepare and manage **safe, sterile feeding** with confidence.

Supplies Checklist

- Prescribed formula or feeds (as advised by your doctor/dietician)
- Feeding sets / syringes (sterile)
- Gloves & PPE
- pH strips or feeding tube position confirmation tools
- Clean water / sterile flush solution
- Dressing materials for PEG site care
- Clamp or cap for tube closure

Tip: Create a dedicated, clean feeding station to avoid contamination.

Safe Feeding Routine

1. **Hand Hygiene First** — Wash hands or use alcohol rub before and after handling the tube.
2. **Position Patient Upright** — At least 30–45° during and for 30–60 min after feeding to reduce aspiration risk.
3. **Check Tube Patency** — Verify placement as advised (pH check, aspirate, visual markings).
4. **Flush Before & After Feed** — Use sterile water to prevent clogs.
5. **Use Correct Feed Volume & Speed** — Avoid rushing or overfilling.
6. **Watch for Intolerance** — Vomiting, coughing, abdominal bloating, diarrhoea.
7. **PEG Site Care** — Clean daily with sterile solution, keep dry, inspect for redness/pus.
8. **Store Formula Safely** — Refrigerate as directed; never use expired feeds.

Red Flags — Call for Help If

- Vomiting, coughing, or choking during feeds
- Sudden breathing difficulty or aspiration signs
- Tube pulled out, displaced or leaking
- Fever, foul-smelling discharge or redness at PEG site
- Persistent diarrhoea or dehydration

Call **Narpavi Homecare 9790016097** or your doctor immediately if any of these occur.

Infection Prevention Tips

- Always follow **WHO hand hygiene** before tube handling.
- Use **sterile syringes and water** for flushing.
- Keep **PEG site clean and dry**; change dressing as advised.
- Disinfect feeding equipment after each use.

Why Use Professional Support

Narpavi Homecare's tube-feeding-trained nurses:

- Safely manage **Ryle's/NG and PEG feeds, flushes, and aspiration prevention.**
- Provide **dietician coordination** for proper nutrition.
- Deliver **digital daily logs & WhatsApp updates** — ideal for **NRI families.**
- Offer **24×7 escalation & hospital backup** for emergencies.



Need Help With Tube Feeding or PEG Care at Home?

Call **9790016097** or **Request a Free Specialty Care Assessment.**

Narpavi Homecare — **Safe, sterile, and doctor-supervised tube feeding & PEG care at your doorstep in Chennai.**