

Using the Structured Coalescent for Phylogeography in BEAST 2

Lecturer: Tim Vaughan

Centre for Computational Evolution
Department of Computer Science, University of Auckland

February 2017



Phylogeography

Mugration models

Structured population
models

Phylogeographic
inference in BEAST 2

Summary

Tutorial

References

What is Phylogeography?

Phylogeography is a field of study concerned with the principles and processes governing the geographic distributions of genealogical lineages, especially those within and among closely related species.

[Avise, 2000]

Structured Coalescent

Phylogeography

Migration models

Structured population models

Phylogeographic inference in BEAST 2


Summary

Tutorial

References

What is Phylogeography?

Early human migrations:



[Wikipedia]

Phylogeography

Migration models

Structured population models

Phylogeographic inference in BEAST 2

Summary

Tutorial


References

[Phylogeography](#)[Migration models](#)[Structured population models](#)[Phylogeographic inference in BEAST 2](#)[Summary](#)[Tutorial](#)[References](#)

Sample	Sequence	Location	Age/Time
1	A A G C T T C A	Place A	0
2	A A G C T T T A	Place B	2
3	A A G T T T C A	Place B	3
4	A A G C T T C A	Place C	3

Phylogeographic inference questions

Common questions include:



Phylogeography

Migration models

Structured population models

Phylogeographic inference in BEAST 2


Summary

Tutorial

References

Phylogeographic inference questions

Common questions include:



Phylogeography

Migration models

Structured population models

Phylogeographic inference in BEAST 2


Summary

Tutorial

References

Phylogeographic inference questions

Common questions include:



Phylogeography

Migration models

Structured population models

Phylogeographic inference in BEAST 2


Summary

Tutorial

References

Phylogeographic inference questions

Common questions include:



Phylogeography

Migration models

Structured population models

Phylogeographic inference in BEAST 2


Summary

Tutorial

References

Phylogeographic inference questions

Common questions include:



Phylogeography

Migration models

Structured population models

Phylogeographic inference in BEAST 2

Summary

Tutorial

References

The usual phylogenetic posterior is:

$$P(T, \mu, \theta | A) = \frac{1}{P(A)} P(A | T, \mu) P(T | \theta) P(\mu) P(\theta)$$

where

$P(A | T, \mu)$ is the *tree likelihood*,

$P(T | \theta)$ is the *tree prior*, and

$P(\mu)$ and $P(\theta)$ are the *parameter priors*.

The usual phylogenetic posterior is:

$$P(T, \mu, \theta | A) = \frac{1}{P(A)} P(A | T, \mu) P(T | \theta) P(\mu) P(\theta)$$

where

$P(A | T, \mu)$ is the *tree likelihood*,

$P(T | \theta)$ is the *tree prior*, and

$P(\mu)$ and $P(\theta)$ are the *parameter priors*.

Where does geography fit in?

Currently there are two main classes of models:

► **Mugration models:**

- Given tree and root location, what is the probability of sample locations?
- Exist in continuous and discrete forms.
- Developed by Phillipe Lemey et al.
[Lemey et al., 2009, Lemey et al., 2010].

► **Structured population models:**

- Given sequences and locations, what is the probability of the location-coloured tree?
- Currently mostly discrete.
- Earliest examples by [Hudson, 1990] and [Notohara, 1990].

Discrete migration model

Structured Coalescent

Phylogeography

Migration models

Structured population
models

Phylogeographic
inference in BEAST 2

Summary

Tutorial

References

Discrete migration model

Structured Coalescent

Phylogeography

Migration models


Structured population
models

Phylogeographic
inference in BEAST 2

Summary

Tutorial

References



Discrete migration model

Structured Coalescent

Phylogeography

Migration models


Structured population
models

Phylogeographic
inference in BEAST 2


Summary

Tutorial

References



Discrete migration model



Structured Coalescent

Phylogeography

Migration models

Structured population models


Phylogeographic inference in BEAST 2

Summary

Tutorial

References

Discrete migration model



Structured Coalescent

Phylogeography

Migration models

Structured population models

Phylogeographic inference in BEAST 2

Summary

Tutorial

References

The usual phylogenetic posterior is:

$$P(T, \mu, \theta | A) = \frac{1}{P(A)} P(A | T, \mu) P(T | \theta) P(\mu) P(\theta)$$

where

A is a sequence alignment,

T is the tree.

Mugration Inference: Modified tree likelihood

Structured Coalescent

Phylogeography

Mugration models

Structured population models

Phylogeographic inference in BEAST 2

Summary

Tutorial

References

The standard phylogenetic posterior is modified:

$$P(T, \mu, \theta | A, L) = \frac{1}{P(A, L)} P(A | T, \mu) P(L | T, M) \\ \times P(T | \theta) P(\mu) P(M) P(\theta)$$

where

L are the sampled locations, and
 M is a matrix specifying the random walk.

Mugration Inference: Modified tree likelihood

Structured Coalescent

Phylogeography

Mugration models

Structured population models

Phylogeographic inference in BEAST 2

Summary

Tutorial

References

The standard phylogenetic posterior is modified:

$$P(T, \mu, \theta | A, L) = \frac{1}{P(A, L)} P(A | T, \mu) P(L | T, M) \\ \times P(T | \theta) P(\mu) P(M) P(\theta)$$

where

L are the sampled locations, and
 M is a matrix specifying the random walk.

Notice the similarity between the two likelihood terms.

Mugration Inference: Modified tree likelihood

Structured Coalescent

Phylogeography

Mugration models

Structured population models

Phylogeographic inference in BEAST 2

Summary

Tutorial

References

The standard phylogenetic posterior is modified:

$$P(T, \mu, \theta | A, L) = \frac{1}{P(A, L)} P(A | T, \mu) P(L | T, M) \\ \times P(T | \theta) P(\mu) P(M) P(\theta)$$

where

L are the sampled locations, and
 M is a matrix specifying the random walk.

Notice the similarity between the two likelihood terms.

Mugration models treat location as just another trait/character.

The following very important assumption made by the migration model posterior:

The following very important assumption made by the migration model posterior:

Samples are to be collected in a manner that is blind to their location.

The following very important assumption made by the migration model posterior:

Samples are to be collected in a manner that is blind to their location.

- Migration models use sample location as data.

The following very important assumption made by the migration model posterior:

Samples are to be collected in a manner that is blind to their location.

- ▶ Migration models use sample location as data.
- ▶ Just as for genetic data, non-random sampling procedures will **bias results**.

Equivalent population genetic model

Structured Coalescent

A helpful way to visualise the migration model is to imagine its effect on the population as a whole:

Phylogeography

Migration models

Structured population models

Phylogeographic inference in BEAST 2


Summary

Tutorial

References

Equivalent population genetic model

A helpful way to visualise the migration model is to imagine its effect on the population as a whole:



Structured Coalescent

Phylogeography

Migration models

Structured population models

Phylogeographic inference in BEAST 2


Summary

Tutorial

References


Equivalent population genetic model

A helpful way to visualise the migration model is to imagine its effect on the population as a whole:




Equivalent population genetic model

A helpful way to visualise the migration model is to imagine its effect on the population as a whole:




Equivalent population genetic model

A helpful way to visualise the migration model is to imagine its effect on the population as a whole:

[Phylogeography](#)[Migration models](#)[Structured population models](#)[Phylogeographic inference in BEAST 2](#)[Summary](#)[Tutorial](#)[References](#)

Equivalent population genetic model

A helpful way to visualise the migration model is to imagine its effect on the population as a whole:




► Migration \implies stochastically varying population sizes.

[Phylogeography](#)[Migration models](#)[Structured population models](#)[Phylogeographic inference in BEAST 2](#)[Summary](#)[Tutorial](#)[References](#)

Equivalent population genetic model

A helpful way to visualise the migration model is to imagine its effect on the population as a whole:



- Migration \implies stochastically varying population sizes.
- A “neutral” model.

Structured Wright-Fisher model

Imagine two sub-populations connected by weak migration:

Structured Coalescent

Phylogeography

Migration models

Structured population models

Phylogeographic inference in BEAST 2


Summary

Tutorial

References


Structured Wright-Fisher model

Imagine two sub-populations connected by weak migration:




Structured Wright-Fisher model

Imagine two sub-populations connected by weak migration:

[Phylogeography](#)[Migration models](#)[Structured population models](#)[Phylogeographic inference in BEAST 2](#)[Summary](#)[Tutorial](#)[References](#)


Structured Wright-Fisher model

Imagine two sub-populations connected by weak migration:

[Phylogeography](#)[Migration models](#)[Structured population models](#)[Phylogeographic inference in BEAST 2](#)[Summary](#)[Tutorial](#)[References](#)


Structured Wright-Fisher model

Imagine two sub-populations connected by weak migration:

[Phylogeography](#)[Migration models](#)[Structured population models](#)[Phylogeographic inference in BEAST 2](#)[Summary](#)[Tutorial](#)[References](#)


Structured Wright-Fisher model

Imagine two sub-populations connected by weak migration:

[Phylogeography](#)[Migration models](#)[Structured population models](#)[Phylogeographic inference in BEAST 2](#)[Summary](#)[Tutorial](#)[References](#)

Structured Wright-Fisher model

Imagine two sub-populations connected by weak migration:




► Model described by [Notohara, 1990].

[Phylogeography](#)[Migration models](#)[Structured population models](#)[Phylogeographic inference in BEAST 2](#)[Summary](#)[Tutorial](#)[References](#)

Structured Wright-Fisher model

Imagine two sub-populations connected by weak migration:



- Model described by [Notohara, 1990].
- Island populations are held constant by respective carrying capacities.

[Phylogeography](#)[Migration models](#)[Structured population models](#)[Phylogeographic inference in BEAST 2](#)[Summary](#)[Tutorial](#)[References](#)

Backwards-in-time process that generates both the tree and ancestral locations.

Phylogeography

Migration models

Structured population models

Phylogeographic inference in BEAST 2

Summary

Tutorial

References

Structured Coalescent

Backwards-in-time process that generates both the tree and ancestral locations.



Structured Coalescent

Backwards-in-time process that generates both the tree and ancestral locations.



Structured Coalescent


Backwards-in-time process that generates both the tree and ancestral locations.



- Structured Coalescent
- Phylogeography
- Migration models
- Structured population models
- Phylogeographic inference in BEAST 2
- Summary
- Tutorial
- References


Structured Coalescent

Backwards-in-time process that generates both the tree and ancestral locations.




Structured Coalescent

Backwards-in-time process that generates both the tree and ancestral locations.



Structured Coalescent

Backwards-in-time process that generates both the tree and ancestral locations.



Structured Coalescent

Phylogeography

Migration models

Structured population models

Phylogeographic inference in BEAST 2


Summary

Tutorial

References


Structured Coalescent

Backwards-in-time process that generates both the tree and ancestral locations.




Structured Coalescent

Backwards-in-time process that generates both the tree and ancestral locations.




Structured Coalescent

Backwards-in-time process that generates both the tree and ancestral locations.




Structured Coalescent

Backwards-in-time process that generates both the tree and ancestral locations.



Structured Coalescent

Backwards-in-time process that generates both the tree and ancestral locations.



Again, the standard phylogenetic posterior is modified:

$$\begin{aligned} P(T, \mathbf{C}, \mu, \theta, \bar{\mathbf{M}}, \vec{\mathbf{N}} | \mathbf{A}, L) &= \frac{1}{P(\mathbf{A} | L)} P(\mathbf{A} | T, \mu) \\ &\times P(T, \mathbf{C} | \vec{\mathbf{N}}, \bar{\mathbf{M}}, L) \\ &\times P(\mu) P(\theta) P(\bar{\mathbf{M}}) P(\vec{\mathbf{N}}) \end{aligned}$$

where

- L are the sampled locations,
- $\bar{\mathbf{M}}$ is the migration rate matrix, and
- \mathbf{C} are the ancestral locations on the tree.

The sample locations and SC model affect the **tree prior**.

The *shape* of the tree is affected by structure.

[Phylogeography](#)[Migration models](#)[Structured population models](#)[Phylogeographic inference in BEAST 2](#)[Summary](#)[Tutorial](#)[References](#)

- ▶ The coalescent tree prior is explicitly conditioned on the sample times.
- ▶ Similarly, the structured coalescent tree prior is conditioned on sample locations.

The structured coalescent makes no assumption about the manner in which samples are collected with respect to location.

- ▶ Sample distribution not used as data.
- ▶ Uneven sampling can reduce inference power, but will *not* bias results!

Phylogeography

Migration models

Structured population models

Phylogeographic inference in BEAST 2

Summary

Tutorial

References


- ▶ Introduced by [\[Kühnert et al., 2016\]](#).
- ▶ A birth-death model of population dynamics in which individuals are permitted to change location due to discrete migration events.
- ▶ Sampling process is explicitly modelled.
- ▶ Birth and death rates may be location-dependent: not "neutral"! (Tree shape affected by structure.)
- ▶ Inference is performed using modified tree prior.

Required packages:

▶ BEAST_CLASSIC

- ▶ Very well supported, BEAUti analysis setup.
- ▶ Tutorial on beast2.org/tutorials.
- ▶ Very fast, allows inference of which migrations are necessary to describe data.
- ▶ Prone to sampling biases.

Discrete Phylogeography (Mugration)



Phylogeography

Mugration models

Structured population models

Phylogeographic inference in BEAST 2

Summary

Tutorial

References

Required packages:

▶ GEO_SPHERE

- ▶ Also well supported, BEAUti analysis setup.
- ▶ Tutorial on beast2.org/tutorials.
- ▶ Output can be summarized using [Spread](#) and visualized using Google Earth.
- ▶ Prone to sampling biases.

Continuous Phylogeography (Mugration)

Structured Coalescent

Phylogeography

Mugration models


Structured population models

Phylogeographic inference in BEAST 2

Summary

Tutorial

References




Required packages:

▶ MultiTypeTree

- ▶ Newer analysis option, BEAUti setup.
- ▶ Tutorial at beast2.org/tutorials.
- ▶ No built-in assumptions regarding sampling procedure.
- ▶ More computationally demanding than mugration models, only smaller numbers of demes are feasible.

Structured Coalescent (full model)



Phylogeography

Migration models

Structured population models

Phylogeographic inference in BEAST 2

Summary

Tutorial

References


[Vaughan et al., 2014]

Required packages:

► BASTA

- Very new analysis option, **no BEAUti setup**.
- Temporary tutorial at github.com/tgvaughan/MultiTypeTree/wiki
- Approximation cuts down on the computational demands of the SC model, allowing many more locations to be considered.
- Produces similar results to MultiTypeTree.

Structured Coalescent (approximation)



[De Maio et al., 2015]

Required packages:

- ▶ bdmm
- ▶ MultiTypeTree
- ▶ SA
- ▶ MASTER

- ▶ Very new analysis option, BEAUti setup possible.
- ▶ Information can be found on the GitHub repository at github.com/denisekuehnert/bdmm.

Birth-death Migration Model

Structured Coalescent

Phylogeography

Migration models


Structured population models

Phylogeographic inference in BEAST 2

Summary


Tutorial

References



[Kühnert et al., 2016]

Birth-death Migration Model



Phylogeography

Migration models

Structured population models

Phylogeographic inference in BEAST 2

Summary

Tutorial

References

[Kühnert et al., 2016]

- ▶ Bayesian phylogeographic methods provide a systematic way of combining geographic and genetic data.
- ▶ BEAST 2 provides two main routes:
 - ▶ Migration models
 - ▶ Structured population models
- ▶ Migration models tend to allow computationally efficient inference, but have questionable foundation and are subject to sampling biases.
- ▶ Structured population models may be more closely tied to the biology and don't necessarily depend on the sampling process.

1. Open MultiTypeTree tutorial at taming-the-beast.github.io/tutorials/Structured-coalescent
2. Start the tutorial!

- Avise, J. C. (2000). *Phylogeography: the history and formation of species*. Harvard university press.
- De Maio, N., Wu, C.-H., O'Reilly, K. M., and Wilson, D. (2015). New routes to phylogeography: A bayesian structured coalescent approximation. *PLoS Genet*, 11(8):e1005421.
- Hudson, R. R. (1990). Gene genealogies and the coalescent process. *Oxford Surveys in Evolutionary Biology*, 7:1.
- Kühnert, D., Stadler, T., Vaughan, T. G., and Drummond, A. J. (2016). Phylodynamics with migration: A computational framework to quantify population structure from genomic data. *Mol Biol Evol*, 33:2102–2116.
- Lemey, P., Rambaut, A., Drummond, A. J., and Suchard, M. A. (2009). Bayesian phylogeography finds its roots. *PLoS Comput Biol*, 5(9):e1000520.
- Lemey, P., Rambaut, A., Welch, J. J., and Suchard, M. A. (2010). Phylogeography takes a relaxed random walk in continuous space and time. *Mol Biol Evol*, 27:1877–1885.
- Notohara, M. (1990). The coalescent and the genealogical process in geographically structured population. *J Math Biol*, 29(1):59–75.
- Vaughan, T. G., Kühnert, D., Poppinga, A., Welch, D., and Drummond, A. J. (2014). Efficient bayesian inference under the structured coalescent. *Bioinformatics*, 30(16):2272–2279.

[Phylogeography](#)[Migration models](#)[Structured population models](#)[Phylogeographic inference in BEAST 2](#)[Summary](#)[Tutorial](#)[References](#)