IV Antibiotic Renal Dose Adjustment (Adults)

Dose by CrCl (Cockcroft-Gault, mL/min). Adults only. NOT for dialysis patients (no RRT on site) - refer. Lab eGFR is a rough screen only; calculate Cockcroft-Gault CrCl for dosing. Status: **green=no adjustment**, **amber=adjust by CrCl**, **red=special caution**.

#	Drug (brand)	Strength	Status	CrCl >50	CrCl 30-50	CrCl 10-29	CrCl <10	Comments
1	**Amikacin** Amikin	500 mg/2 mL	Special	No adj (CrCl >=60). 50-59: 5-7.5 mg/kg q12h.	40-50: 5-7.5 mg/kg q12h. 30-39: q24h.	20-29: q24h. 10-19: single dose, then individualise.	Single dose, then individualise.	Dose on actual BW (adj if TBW>1.2xIBW). No TDM: monitor renal fn & hearing. Native >=60/40-60/20-40/<20.
2	**Amoxicillin/clavulanate** Augmentin	1.2 g	Adjust	No adjustment.	No adj (CrCl >=30).	Initial 1 g, then 500 mg q12h.	Initial 1 g, then 500 mg q12-24h.	Doses as amoxicillin component (IV Canadian product).
3	**Amphotericin B (conv. deoxycholate)** Fungizone	50 mg	Special	No dose adj by CrCl.	No dose adj by CrCl.	No dose adj by CrCl.	No dose adj by CrCl.	Nephrotoxic: lipid form preferred if impaired. Na load (1 L NS), longer infusion, avoid nephrotoxins; monitor kidney fn.
4	**Ampicillin** Ampicin	1 g	Adjust	1-2 g q6h (no change).	1-2 g q8h.	15-29: 1-2 g q12h. <15: 1-2 g q24h.	1-2 g q24h.	Expert opinion. If usual 2 g q4h: q4h/q6h/q8h/q12h.
5	**Cefazolin** Ancef, Kefzol	1 g	Adjust	1-2 g q8h (no change).	1-2 g q8-12h.	500 mg-1 g q12h (2 g q12h if severe).	500 mg-1 g q24h (<=10).	Consider an initial full dose for severe infection before reducing.
6	**Cefotaxime** Claforan	1 g	Adjust	1-2 g q8h (no change).	1-2 g q12h.	1-2 g q12h.	1-2 g q24h (<=10).	US limited availability. If usual q6h: q8h/q12h; if 2 g q4h: q6-8h/q12h.
7	**Ceftazidime** Fortum, Fortaz	1 g	Adjust	1 g q8h (no change).	1 g q12h (31-50).	16-30: 1 g q24h. <=15: 500 mg q24h.	500 mg q24h (<=15).	If usual 2 g q8h: 2 g q12h / 2 g q24h / 1 g q24h.
8	**Ceftriaxone** Rocephin	1 g	No adj	No adjustment.	No adjustment.	No adjustment.	No adj (>2 g/day: monitor).	Biliary elimination; not significantly dialysed.
9	**Ciprofloxacin** Ciprobay, Cipro	400 mg/vial	Adjust	400 mg q8-12h.	400 mg q8-12h.	200-400 mg q12-24h (load 400 if 200 q24h).	200-400 mg q12-24h.	Severe infections: higher end of interval.
10	**Clindamycin** Dalacin C, Cleocin	600 mg/4 mL	No adj	No adjustment.	No adjustment.	No adjustment.	No adjustment.	Hepatically eliminated; not dialysed.
11	**Cloxacillin** Cloxapen	1 g	No adj	No adjustment.	No adjustment.	No adj (CrCl >=10).	No specific adj; monitor; consider <=8 g/day.	US not-available product. Nephrotoxicity at >=8 g/day.
12	**Colistimethate sodium** Coly-Mycin	150 mg/vial	Special	300 mg CBA/day (>=50).	183-250 mg CBA/day.	150-183 mg CBA/day.	117 mg CBA/day (<10).	Load 300 mg CBA. CBA 1 mg = 30,000 IU; 150 mg vial is CBA. Adjusted BW. Full ladder in notes.
13	**Ertapenem** Invanz	1 g	Adjust	1 g daily (no change).	1 g daily (>30).	500 mg daily (<=30).	500 mg daily.	No BSA adjustment of CrCl needed.
14	**Gentamicin** Garamycin	80 mg/2 mL	Special	No adj (CrCl >=60). 50-59: usual dose q12h.	40-50: usual dose q12h. 30-39: q24h.	20-29: q24h. 10-19: single dose, then individualise.	Single dose, then individualise.	Dose on actual BW. No TDM: monitor renal fn & hearing. Native >=60/40-60/20-40/<20.
15	**Meropenem** Meronem, Merrem	1 g/vial	Adjust	1 g q8h (no change).	1 g q12h (>25-50).	10-25: 500 mg q12h. 26-29: 1 g q12h.	500 mg q24h.	If usual 2 g q8h: 2 g q12h / 1 g q12h / 1 g q24h or 500 mg q12h. Extended 3-h infusion improves attainment.
16	**Metronidazole** Flagyl	0.5% 100 mL (500 mg)	No adj	No adjustment.	No adjustment.	No adjustment.	No adj (500 mg q12h may suffice).	Single-dose regimens need no adj. Monitor in CrCl <30.
17	**Penicillin G sodium** Benzylpenicillin	5 MU (3 g)	Adjust	No adjustment (100%).	50-75% of usual dose.	25-50% of usual dose.	20-25% of usual dose.	% of usual indication-specific dose; continuous or divided q4-6h. Monitor for neurotoxicity.
18	**Piperacillin/tazobactam** Tazocin, Zosyn	4.5 g/vial	Adjust	4.5 g q6h (40-100; >100 extended infusion).	40-50: 4.5 g q6h. 30-39: 4.5 g q8h or 3.375 g q6h.	20-29: 4.5 g q8h or 3.375 g q6h. <20: 4.5 g q12h or 2.25 g q6h.	4.5 g q12h or 2.25 g q6h (<20).	Combined pip+tazo. Extended infusion (3-4 h) preferred for high CrCl/severe.
19	**Remdesivir** Veklury	100 mg/vial	No adj	No adjustment.	No adjustment.	No adjustment.	No adjustment.	Earlier eGFR <30 restriction no longer applied per current UpToDate.

#	Drug (brand)	Strength	Status	CrCl >50	CrCl 30-50	CrCl 10-29	CrCl <10	Comments
20	**Vancomycin** Vancocin	500 mg/vial	Special	50-90: 15-20 mg/kg q12h. >90: q8-12h.	10-15 mg/kg q24h (15 to <50).	15-29: 10-15 mg/kg q24h. <15: q48-72h.	10-15 mg/kg q48-72h (<15).	Initial dose on actual BW. Normally titrated to levels; no TDM: monitor renal fn & response. Max loading 3 g.

Colistin full maintenance ladder (total daily CBA, 2 divided doses): CrCl >=90: 360; 80-90: 340; 70-80: 300; 60-70: 275; 50-60: 245; 40-50: 220; 30-40: 195; 20-30: 175; 10-20: 160; 5-10: 145; <5: 130 mg/day.
No on-site TDM: aminoglycosides (amikacin, gentamicin) and vancomycin are normally titrated to serum levels; monitor renal function, hearing, and response. Clinical reference only - verify against current UpToDate. Sole reference: UpToDate / Lexicomp, accessed 2026-06-23.