



Proximity Effects *in* *Altermagnetic Systems*

~ o ~

a bachelor thesis.

Written by Arto Steffan,
under the supervision of prof. dr. Jacob Wüsthoff Linder,
and corrected by prof. dr. Wolfgang Belzig.

Universität
Konstanz



NTNU – Trondheim
Norwegian University of
Science and Technology

Acknowledgment

...

_____ ~ _____

Contents

| | | |
|----------|---|-----------|
| 1 | Theoretical Background | 1 |
| 1.1 | Bosons and fermions | 1 |
| 1.2 | The second quantisation | 3 |
| 1.2.1 | Second quantisation for fermions | 3 |
| 1.2.2 | Second quantisation for bosons | 5 |
| 1.3 | Basis transformation | 5 |
| 1.3.1 | Field operators | 6 |
| 1.4 | Interactive electron gas | 7 |
| 2 | Superconductivity | 9 |
| 2.1 | Theoretical framework and BCS theory | 9 |
| 2.1.1 | The non-interacting electron gas | 10 |
| 2.1.2 | Fermi-Liquid - the interacting case | 11 |
| 2.1.3 | Instability due to attractive Interactions | 12 |
| 2.1.4 | Phonon-mediated attractive interactions reformulate | 12 |
| 2.1.5 | Contraction of the effective Hamiltonian | 14 |
| 2.1.6 | On our way to the BCS-theory | 15 |
| 2.1.7 | Transition temperature and energy gap | 18 |
| 2.1.8 | Generalized gap equation, s-wave and d-wave superconductivity | 18 |
| 3 | Bogoliubov-de Gennes Formalism | 21 |
| 3.1 | Tigh Binding Model | 22 |
| 3.1.1 | Non-interacting electrons | 22 |
| 3.1.2 | Superconductivity | 23 |
| 3.2 | A more symmertic Hamiltonian | 24 |
| 3.3 | Eigenvalues | 25 |
| 3.4 | Diagonalization | 25 |
| 3.4.1 | Superconducting Gap | 26 |
| 4 | Results | 27 |
| 5 | Literature | 30 |

1 Theoretical Background

In order to describe the superconductors we are going to introduce the second quantisation formalism. It allows us to describe the wavefunction of a system using creation and annihilation operators over energy states of the system and simplify a lot the notation. The mathematical foundation of this formalism lays in the Hilbert space, its dual space and furthermore we are going to introduce the Fock space.

It's also relevant for our study that we are going to work on fermions. The formalism stays the same for bosons but the results are fundamentally different. One can mention the Pauli principle as an example which only applies on fermions and can be derived with the help of the second quantisation.

1.1 Bosons and fermions

We consider without loss of generality the following hamiltonian.

$$\hat{H} = \hat{H}_0 + \hat{H}_I \quad (1)$$

with the single particle operator \hat{H}_0 and the interaction operator \hat{H}_I :

$$\hat{H}_0 = \sum_{i \in \llbracket N \rrbracket} \hat{h}_i(x_i), \quad \hat{h}_i(x_i) = -\frac{\hbar^2}{2m} \nabla_i^2 + \hat{U}(x_i)$$

Where we introduce the notation $\llbracket N \rrbracket = \{n \in \mathbb{N} : n \leq N\}$. We call it single particle operator because the operator only applies on a particle. It may depend from the particle's position \mathbf{r} or spin s : $x_i := (\mathbf{r}, s) \in \mathcal{X} \subseteq \mathbb{R}^3 \times \mathbb{S}$. For example we have for an electron $\mathbb{S} = -\frac{1}{2}, \frac{1}{2}$. A single particle operator is in this case the sum of the kinetic- and potential energy operators.

Further we describe a quantum state that a particle can occupy with a wavefunction $\phi_\nu(x)$, which own a certain energy $\epsilon_\nu \in \mathbb{R}$. This energy depends on the wavevector and the spin of the particle: $\nu = (\mathbf{k}, \sigma)$. The fundamental equation of quantum mechanics relates the wave function with the hamiltonian using the energy of the state:

$$\hat{h}\phi_\nu(x) = \epsilon_\nu\phi_\nu(x)$$

The wave function lay in the Hilbert space [more details?]. Therefor $\phi_\nu(x)$ are eigenfunction or -states of the Hamiltonian with eigenvalues ϵ_ν . Further the wavefunction should build an orthonormal basis:

$$\int_{\mathcal{X}} \phi_{\nu'}(x)\phi_\nu(x)dx = \delta_{\nu'\nu}.$$

ν and ν' are two different states. We introduced here the korenker delta $\delta_{\nu'\nu}$ which is one when the two indicies are equal and zero otherwise. Because the spin s is not continuous one can understand the integral in the following way:

$$\int_{\mathcal{X}} dx = \sum_{s \in \mathbb{S}} \int_{\mathbb{R}^3} d^3r$$

where $\int_{\mathbb{R}^3} d^3r = \int_{\mathbb{R}} \int_{\mathbb{R}} \int_{\mathbb{R}} dr_1 dr_2 dr_3$. We integrate over all possible states.

Now that we can describe one particle we want to describe a system containing many instance of that particle. The wavefunction sums up all possible combination of wavefunction in the system and should stay normalized. A combination is discribed as the product of the wavefunction of the particle in a certain state. These particle can be swaped and therefore we need to consider all combinations. We restrict ourselves to fermions and bosons. We admit having $N \in \mathbb{N}$ paricles in the system.

Bosons The many-particle wavefunction of the bosons is a symetric (exponent S) under swap of two particles.

$$\Phi^{(S)}(x_1, \dots, x_N) = \left(N! \prod_N (n_\nu)! \right)^{-\frac{1}{2}} \sum_{P \in S_n} P \phi_{\nu_1}(x_1) \cdot \dots \cdot \phi_{\nu_N}(x_N)$$

This represents the an eigenfunction of the non interacting bosonic-Hamiltonian. We used n_ν , which represents the number of particle in the state ν . Therefor we usaly call it the occupation number of the state ν . For fermion this integer has no constrain in general. The permutation set S_n contains all the possbile combinations of x_i in the state ν_j for $i, j \in \llbracket N \rrbracket$.

Fermions Fermions are a bit different, their many-particle fermion wavefunction is antisymmetric under swap of two particles. We denote it as

$$\Phi^{(A)}(x_1, \dots, x_N) = (N!)^{-\frac{1}{2}} \sum_{P \in S_n} \text{sgn}(P) \cdot P \phi_{\nu_1}(x_1) \cdot \dots \cdot \phi_{\nu_N}(x_N).$$

which is an eigenfunction of the non interacting fermionic-Hamiltonian. sgn represents the signum function. Applied on a permutation P it is one if P is even and minus one if P is even. We already know that the Pauli principle implies that it can be up to one particle in each energy state. We therefore have $n_\nu \in \{0, 1\}$. The normalisization factor is the same but the product over the ones vanishes.

At this point one might have recognised the formula of the determinant

$$\Phi^{(A)}(x_1, \dots, x_N) = (N!)^{-\frac{1}{2}} \det \begin{pmatrix} \phi_{\nu_1}(x_1) & \dots & \phi_{\nu_1}(x_N) \\ \vdots & & \vdots \\ \phi_{\nu_N}(x_1) & \dots & \phi_{\nu_N}(x_N) \end{pmatrix},$$

which vanishes if two rows or columns are identical. We usually describe this expression as the Slater determinant. This means that the probability of finding two fermions in the same state is zero. This is the Pauli principle. Only one or no particle may occupy each state.

Further we encounter a major problem. The many-particle wave function of fermions is defined up to a sign. For instance if we consider two particles “having” x_1 and x_2 , we have two possible states ν_1 and ν_2 . Two possible solutions are

$$\begin{aligned}\Phi^{(A_1)} &= \frac{1}{\sqrt{2}}(\varphi_{\nu_1}(x_1)\varphi_{\nu_2}(x_2) - \varphi_{\nu_1}(x_2)\varphi_{\nu_2}(x_1)) \\ \text{or } \Phi^{(A_2)} &= \frac{1}{\sqrt{2}}(\varphi_{\nu_1}(x_2)\varphi_{\nu_2}(x_1) - \varphi_{\nu_1}(x_1)\varphi_{\nu_2}(x_2)) \\ &= -\Phi^{(A_1)}.\end{aligned}$$

This sign difference may lead to computation errors. We aim to give a labeling to our states when we count them and keep it when it comes to build the Slater determinant.

These bosonic and fermionic wavefunctions are eigenstates of the Hamiltonian \hat{H}_0 and the corresponding eigenvalue E_ν is given by summing the energy of each state times its occupation number: $E_\nu = \sum_\nu \epsilon_\nu n_\nu$. For this reason it's important that they build an orthonormal basis:

$$\int_{\mathcal{X}^N} \Phi_a^*(x_1, \dots, x_N) \Phi_b(x_1, \dots, x_N) d^N x = \delta_{ab}.$$

Therefore we can expand any many-particle wavefunction Ψ as the linear combination of these:

$$\Psi = \sum_a f_a \Phi_a(x_1, \dots, x_N)$$

where f_a is a coefficient and a a labeling.

What we just discussed is the so called first quantisation- or wave function formalism. Now we intend to introduce a better way of describing our system.

1.2 The second quantisation

For a better description of the many-particle system we introduce a simpler notation. The second quantisation lays on three important objects. States described as “ket”. We put any relevant information between the ket e.g. $|\mathbf{k}, \sigma, \dots\rangle$. Then we need operators that act on these states to allow interactions in the system. We need an operator that creates a state and another that annihilates a state.

States In this section we describe a state as the number of particles that occupies each single-particle state. Therefore we order the states $1 < 2 < \dots < N$. We then can describe the wave function as follows $|n_1, \dots, n_N\rangle$.

Further the state where no particles are present is called the vacuum state and we denote it as $|0_{\nu_1}, \dots, 0_{\nu_N}\rangle = |0\rangle$.

1.2.1 Second quantisation for fermions

Creation operator c_ν^\dagger The creation operator adds a particle in the state that is concerned and rephases the state:

$$c_\nu^\dagger |n_1, \dots, n_\nu, \dots\rangle = (-1)^{\sum_{\mu < \nu} n_\mu} (1 - n_\nu) |n_1, \dots, n_\nu + 1, \dots\rangle$$

We notice the $(1 - n_\nu)$ term which avoids to create a particle at the state, if one already exists. This is the expression of the Pauli-principle. and we can then construct a state by applying this operator after another on the vacuum state. To avoid the minus one to add a negative sign, we start from the end and add the particles backwards in the order of the state:

$$|n_1, \dots, n_N\rangle = (c_1^\dagger)^{n_1} \cdot \dots \cdot (c_N^\dagger)^{n_N} |0\rangle$$

Annihilation operator c_ν Likewise the annihilation operator destroys a particle in the corresponding state. The operator reads

$$c_\nu^\dagger |n_1, \dots, n_\nu, \dots\rangle = (-1)^{\sum_{\mu < \nu} n_\mu} (n_\nu) |n_1, \dots, n_\nu - 1, \dots\rangle.$$

We can easily recognise that due to the n_ν -term, destroying a particle that doesn't exist gives zero, so it's only possible to destroy particle that exist. Further we intend to introduce some few computation rules that are going to help us later.

The anticommutator of two operator reads $[A, B]_+$ or $\{A, B\} := AB + BA$ and is an operator as well. We're going to stick with $[A, B]_+$ since it's more consistent with the commutator notation $[A, B]_-$ (or simply $[A, B]$).

The following results are obtained by separating the $\nu = \mu$ from the $\nu \neq \mu$. We must also say that the dagger \dagger should be understood as the complex transpose of the operator and $(AB)^\dagger = B^\dagger A^\dagger$.

$$[c_\nu, c_\mu]_+ = 0 \quad (\text{fer}_1)$$

$$[c_\nu^\dagger, c_\mu^\dagger]_+ = 0 \quad (\text{fer}_2)$$

$$[c_\nu^\dagger, c_\mu]_+ = \delta_{\mu, \nu} \quad (\text{fer}_3)$$

We can then combine the creation and annihilation operator to count the number of particles in a state:

$$c_\nu^\dagger c_\nu |n_1, \dots, n_\nu, \dots\rangle = n_\nu |n_1, \dots, n_\nu, \dots\rangle.$$

From this we can define the number operator $\hat{n}_\nu := c_\nu^\dagger c_\nu$ which we can combine in the total number operator

$$\hat{N} = \sum_\nu \hat{n}_\nu, \quad \text{where logically } N = \sum_\nu n_\nu$$

if we apply the operator on a state.

Second quantisation description of the single- and two- particle operators We first need to make an important observation between the Slater determinant and the single particle state to understand the following. First we introduce two basis element $|\Phi_\alpha\rangle$ and $|\Phi_\beta\rangle$, which can be many-particles eigenstate of the system. We can also call them Slater determinant. Further we introduce the probability of the configuration $|\Phi_\alpha\rangle$ to scatter into the $|\Phi_\beta\rangle$ due to the action of an operator A (momentum, potential, interactions,...). This is described by the matrix element $\langle\Phi_\alpha|A|\Phi_\beta\rangle$ which involves the single particle states $|\alpha_1\rangle, \dots, |\alpha_N\rangle$ of $|\Phi_\alpha\rangle$ and $|\beta_1\rangle, \dots, |\beta_N\rangle$ of $|\Phi_\beta\rangle$.

$$\langle\Phi_\alpha|A|\Phi_\beta\rangle = \sum_{i,j} C_{ij} \langle\alpha_i|A|\beta_j\rangle$$

involving some constants C_{ij} . This describes the overlap of the two configurations, after that we modified $|\Phi_\beta\rangle$ with A . On the right hand side (r.h.s) we introduced the bracket scalar product notation. The bra $\langle\alpha|$ lives in the dual space of the Hilbert space. One reads it as the complex transpose of $|\alpha\rangle$.

We recall the single particle Hamiltonian we introduced earlier. Its second quantisation representation reads

$$\hat{H}_0 = \sum_{i \in \mathbb{N}} \hat{h}(x_i) \rightsquigarrow \sum_{\alpha, \beta} \langle\alpha|\hat{h}|\beta\rangle c_\alpha^\dagger c_\beta \quad (2)$$

where α and β are single-particle states of the system. $c_\alpha^\dagger c_\beta$ tries to transfer a fermion from the state $|\beta\rangle$ to the state $|\alpha\rangle$. We have

$$\langle\alpha|\hat{h}|\beta\rangle = \int \varphi_\alpha^*(x) \hat{h}(x) \varphi_\beta(x) dx \quad (3)$$

where x still represents the position and the spin of the particle.

\hat{h} is a single particle operator, it means it acts on one particle at a time. Two states are going to be changed. $|\alpha\rangle$ loses a particle and $|\beta\rangle$ gains one. We say for instance, that the configuration before the scattering is $|\Phi\rangle$ and after the scattering is $|\Phi'\rangle$. This means, if our two Slater

determinant $|\Phi\rangle$ and $|\Phi'\rangle$ differs in more than two state, there are some scattering that we can't describe, so the overlap must be zero. We allow only two states to be modified. Otherwise the single-particle states differ and due to their orthonal properties, we get a zero.

Similarly for the two-particle operator we have

$$\hat{H}_I = \frac{1}{2} \sum_{i \neq j \in \llbracket N \rrbracket} \hat{v}(x_i, x_j) \rightsquigarrow \frac{1}{2} \sum_{\alpha, \beta, \gamma, \delta} \langle \alpha \beta | \hat{v} | \gamma \delta \rangle c_\alpha^\dagger c_\beta^\dagger c_\gamma c_\delta \quad (4)$$

involving a more nested overlap of the four states:

$$\langle \alpha \beta | \hat{v} | \gamma \delta \rangle = \int \int \varphi_\alpha^*(x) \varphi_\beta^*(x') \hat{v}(x, x') \varphi_\gamma(x) \varphi_\delta(x') dx dx'. \quad (5)$$

which modifies four states so the overlap of two slatter determinant vanishes if the determinant differ in at least four states.

The l.h.s of the equation is the matrix element $\langle \Phi_\alpha | \hat{v} | \Phi_\beta \rangle$ of the the operator \hat{v} , which involves two basis state $|\Phi_\alpha\rangle$ and $|\Phi_\beta\rangle$. On the r.h.s we have a description with wavefunctions, which own to the first quantisation. What we have here is the bridge between the first and the second quantisation. One could compute each side separately and notice that both formalism lead to the same result.

1.2.2 Second quantisation for bosons

1.3 Basis transformation

Until now considered the wavefunction in a restricted basis $\{\varphi_\alpha(x)\}$. A wave function is defined as the overlap between the basis and the state:

$$\phi(x) = \langle x | \alpha \rangle.$$

Let us now define a more general new operator that create a states $|\alpha\rangle = a_\alpha |0\rangle$ for fermion and bosons. We asume that we have another basis $\{|\tilde{\alpha}\rangle\}$. We now want to show, that the new operator a_α can be expressed as a linear combination of the other operator $a_{\tilde{\alpha}}$. This will be a usefull too to jump from a basis to another. We first notice the following identity relations:

$$\mathbb{I} = \sum_\alpha |\alpha\rangle \langle \alpha| = \sum_{\tilde{\alpha}} |\tilde{\alpha}\rangle \langle \tilde{\alpha}|$$

this allow us to write

$$a_\alpha^\dagger |0\rangle = |\alpha\rangle = \sum_{\tilde{\alpha}} |\tilde{\alpha}\rangle \underbrace{\langle \tilde{\alpha} | \alpha \rangle}_{\text{scalar}} = \sum_{\tilde{\alpha}} \langle \tilde{\alpha} | \alpha \rangle |\tilde{\alpha}\rangle$$

inverting the indicies leads to the same result, which leads to the transformation rules

$$\begin{aligned} a_\alpha^\dagger &= \sum_{\tilde{\alpha}} \langle \tilde{\alpha} | \alpha \rangle a_{\tilde{\alpha}}^\dagger \\ a_\alpha &= \sum_{\tilde{\alpha}} \langle \alpha | \tilde{\alpha} \rangle a_{\tilde{\alpha}}. \end{aligned}$$

Further we can use these relations to express a wavefunction in the basis of another wavefunction. We have

$$\phi_\alpha(x) = \langle x | \alpha \rangle = \langle x | \left(\sum_{\tilde{\alpha}} \langle \tilde{\alpha} | \alpha \rangle |\tilde{\alpha}\rangle \right) = \sum_{\tilde{\alpha}} \langle \tilde{\alpha} | \alpha \rangle \langle x | \tilde{\alpha} \rangle = \sum_{\tilde{\alpha}} \langle \tilde{\alpha} | \alpha \rangle \varphi_{\tilde{\alpha}}(x).$$

Inverting α and $\tilde{\alpha}$ leads as well $\varphi_{\tilde{\alpha}}(x) = \sum_\alpha \langle \alpha | \tilde{\alpha} \rangle \phi_\alpha(x)$.

Morover we can show that the basis transformation is unitary. This is an important feature because we can simplify the calculation by changing the basis but the result we seek for would remain the same after the unitary transformation. Such transformations plays a major role

later in the thesis. We can save the $\langle \tilde{\alpha} | \alpha \rangle$ in a matrix D and prove that this matrix is unitary. We have

$$\begin{aligned} \langle \tilde{\alpha} | \tilde{\beta} \rangle &= \sum_{\gamma} \langle \tilde{\alpha} | \gamma \rangle \langle \gamma | \beta \rangle = \sum_{\gamma} \langle \tilde{\alpha} | \gamma \rangle \langle \beta | \gamma \rangle^* \\ &= \sum_{\gamma} D_{\alpha\gamma} D_{\beta\gamma}^* = \sum_{\gamma} D_{\alpha\gamma} D_{\gamma\beta}^\dagger = (DD^\dagger)_{\alpha\beta}. \end{aligned}$$

meanwhile $\langle \tilde{\alpha} | \tilde{\beta} \rangle = \delta_{\tilde{\alpha}\tilde{\beta}}$ so that $DD^\dagger = \mathbb{I}$. The matrix D is therefore unitary. [but do we also have the orthonormality of if we take two different basis?](#).

The last important step is to show that the basis transformation keeps the anti-, commutation relations. Let us for the seek of readability use the notation $[A, B]_\xi = AB - \xi BA$ where $\xi = -$ for bosons and $\xi = +$ for fermions. We have for exemple using $[a_\alpha, a_{\alpha'}^\dagger]_\xi = \delta_{\alpha\alpha'}$

$$[a_{\tilde{\alpha}}, a_{\tilde{\alpha}'}^\dagger]_\xi = \sum_{\alpha\alpha'} \langle \tilde{\alpha} | \alpha \rangle \langle \alpha' | \tilde{\alpha}' \rangle [a_\alpha, a_{\alpha'}^\dagger]_\xi = \langle \tilde{\alpha} | \tilde{\alpha}' \rangle = \delta_{\tilde{\alpha}\tilde{\alpha}'}.$$

The first step follows after splitting the commutator in two parts, insert the transformation and recombine the new commutator. The last step involves the orthonormality of the basis. On the same way one can prove $[a_{\tilde{\alpha}}, a_{\tilde{\alpha}'}] = [a_{\tilde{\alpha}}^\dagger, a_{\tilde{\alpha}'}^\dagger] = 0$.

1.3.1 Field operators

Later in this thesis we are going to use field operators to describe an order parameter of a superconductive system. These are creation and annihilation operators that are defined in the $|x\rangle$ -space-spin basis regarding another basis $\{|\alpha\rangle\}$. We here give the state as an argument and not as an index anymore. Despite the notation the following operators musn't be confused with a wavefunction.

$$\hat{\psi}^\dagger(x) = \sum_{\alpha} \langle \alpha | x \rangle a_{\alpha}^\dagger = \sum_{\alpha} \varphi_{\alpha}^*(x) a_{\alpha}^\dagger \quad (6)$$

$$\hat{\psi}(x) = \sum_{\alpha} \langle \alpha | x \rangle a_{\alpha} = \sum_{\alpha} \varphi_{\alpha}(x) a_{\alpha} \quad (7)$$

involving fermionic or bosonic operators a . We can then annihilate and create a particle at a spin-space location x . Because these operators are involved, the commutation property respects which particle we are describing. Using the result we had for $[a_{\alpha}, a_{\alpha'}^\dagger]_\xi$, $[a_{\alpha}, a_{\alpha'}^\dagger]_\xi$ and $[a_{\alpha}^\dagger, a_{\alpha'}^\dagger]_\xi$ where $\xi = -$ for the bosons and $+$ for the fermions. We find the following commutation relations:

$$\begin{aligned} \left[\hat{\psi}(x), \hat{\psi}(x') \right]_\xi &= \left[\hat{\psi}(x)^\dagger, \hat{\psi}^\dagger(x') \right]_\xi = 0 \\ \left[\hat{\psi}(x), \hat{\psi}^\dagger(x') \right]_\xi &= \delta(x - x') \end{aligned}$$

In the last expression we obtain a $\langle x | x' \rangle$ which can't normalize the $\{|x\rangle\}$ -basis so instead of a Kronecker delta $\delta_{xx'}$ we get a delta distribution $\delta(x - x')$. The goal is now to describe the Hamiltonian using these field operators. Therefore we rebuild the field operator in the Hamiltonian using a $\{|x\rangle\}$ basis

$$\begin{aligned} \hat{H}_0 &= \int \hat{\psi}^\dagger(x) \hat{h}(x) \hat{\psi}(x) dx \\ \hat{H}_I &= \frac{1}{2} \int \int \hat{\psi}^\dagger(x) \hat{\psi}^\dagger(x') \hat{v}(x, x') \hat{\psi}(x') \hat{\psi}(x) dx' dx \end{aligned}$$

We already know that the first and second quantisation are equivalent. We could now insert the definition of the field operators 6 and 7 in the Hamiltonian and obtain a wave function description. Then we can just use the first to second quantisation translation 3 and 5 to a the second quantised representation. The result is a generalisation of the expression with the operators a .

1.4 Interactive electron gas

As we are later going to study, the electron are allowed to interact with each other. During such processes a photon is usaly carrying the momentum transfert of the scattering. We're going to use the formalism we introduced to find a second quantisation representation of the interacting Hamiltonian.

The system we're stying is a cube of side length L with wolume $\Omega = L^3$ with N electrons. Further we consider a periodic boundary condition for an arbitrary position vector $\mathbf{r} = (x, y, z)$:

$$(L, y, z) = (0, y, z) , (x, L, z) = (x, 0, z) , (x, y, L) = (x, y, 0).$$

We use the general form of the Hamiltonian 1 introduced in the beggining. We have a Coulomb potential and a kinetic energy term.

$$H_0 = T + U = - \sum_{i \in \llbracket N \rrbracket} \frac{\hbar^2}{2m} \nabla_i^2 + \sum_{i \in \llbracket N \rrbracket} u(x_i) \quad (8)$$

$$H_I = V = \sum_{i \neq j \in \llbracket N \rrbracket} v(x_i, x_j) \quad (9)$$

Where the pairwise potential is just a Coulomb potential $v(x_i, x_j) = \frac{e^2}{4\pi\epsilon_0|x_i - x_j|} = v(|x_i - x_j|)$. [drop of the hat?](#) We aim to describe the Hamiltonian in the momentum space, which is more convenient for the second quantisation. Therefore we first need to find an expression for the wavevector \mathbf{k} . We describe a state $\alpha = (\mathbf{k}, \sigma)$ at a spin-space coordinate $x = (\mathbf{r}, s)$. First we take a plane wave solution of the Schrödinger equation:

$$\psi_{\mathbf{k}}(\mathbf{r}) = \frac{1}{\sqrt{\Omega}} e^{i\mathbf{k} \cdot \mathbf{r}} \chi_{\sigma}(s)$$

The periodicity of the system allows us to write $\psi(x=0) = \psi(x=L)$ using the boundary condition and obtain $1 = e^{ik_x L}$ as well as for the two other coordinates. The results reflects in the wavevector \mathbf{k} :

$$\mathbf{k} = \frac{2\pi}{L} (n_x, n_y, n_z) , \quad n_i \in \mathbb{Z}.$$

As we saw earlier the eigenfunctions build a complete basis:

$$\begin{aligned} \int \psi_{\alpha'}^*(x) \psi_{\alpha}(x) dx &= \int \sum_s \psi_{\alpha'}^*(x) \psi_{\alpha}(x) dx \\ &= \frac{1}{\Omega} \int e^{i\mathbf{r} \cdot (\mathbf{k} - \mathbf{k}')} \underbrace{\sum_s \delta_{s\sigma} \delta_{s\sigma'}}_{\delta_{\sigma\sigma'}} dx \\ &= \delta_{\mathbf{k}\mathbf{k}'} \delta_{\sigma\sigma'} = \delta_{\alpha\alpha'}. \end{aligned}$$

where the integral over the exponential function diverges if $\mathbf{k} \neq \mathbf{k}'$ so we use the case $\mathbf{k} = \mathbf{k}'$ and the integral is zero otherwise.

The kinetic energy operator is a single particle operator so we have according to 2 we have:

$$T = \sum_{\alpha, \alpha'} \langle \alpha | \frac{\hat{\mathbf{p}}^2}{2m} | \alpha' \rangle c_{\alpha}^{\dagger} c_{\alpha'}.$$

Now we can juste use $\hat{\mathbf{p}} = i\hbar \nabla$ and to compute this expression we use its first quantised form (see 3) indvolving the $\delta_{\sigma\sigma'}$ trick we introduced in the derivation of the complete basis. We also use $\mathbf{k}' = \mathbf{k}$ and hide the $\frac{1}{\Omega}$ is the $\delta_{\mathbf{k}\mathbf{k}'}$. The result is

$$T = \sum_{\alpha, \alpha'} \delta_{\alpha\alpha'} \frac{\hbar \mathbf{k}^2}{2m} c_{\alpha'}^{\dagger} c_{\alpha} = \sum_{\alpha} \underbrace{\frac{\hbar \mathbf{k}^2}{2m}}_{=: \epsilon_{\mathbf{k}}} c_{\alpha}^{\dagger} c_{\alpha}$$

we recognise the occupation number operator $\hat{n}_{\alpha} = c_{\alpha}^{\dagger} c_{\alpha}$ and the energy of the state $\epsilon_{\mathbf{k}}$. This is a quite meaningful result, the non-interacting energy part of the system is the product of

the energy of a state with the number of particle within that state, summed over all states.

We don't consider the single particle external potential for now.

For the interaction potential, we have a two-particle operator. This is described by equation 4 and requires 5 to be solved.

$$V = \frac{1}{2} \sum_{\alpha, \beta, \gamma, \delta} \langle \alpha \beta | v | \gamma \delta \rangle c_{\alpha}^{\dagger} c_{\beta}^{\dagger} c_{\gamma} c_{\delta}$$

Introducing these in V using $v(\mathbf{r}, \mathbf{r}') = v(\mathbf{r} - \mathbf{r}')$ leads to:

$$\langle \alpha \beta | v | \gamma \delta \rangle = \frac{1}{\Omega^2} \delta_{\sigma_{\alpha} \sigma_{\gamma}} \delta_{\sigma_{\beta} \sigma_{\delta}} \int \int e^{i\mathbf{r}(\mathbf{k}_{\gamma} - \mathbf{k}_{\alpha})} e^{i\mathbf{r}'(\mathbf{k}_{\delta} - \mathbf{k}_{\beta})} v(\mathbf{r} - \mathbf{r}') d\mathbf{r}' d\mathbf{r}$$

we can make a substitution $\mathbf{R} = \mathbf{r} - \mathbf{r}'$, add and subtract a $(\mathbf{k}_{\gamma} - \mathbf{k}_{\alpha})\mathbf{r}'$ and obtain

$$\langle \alpha \beta | v | \gamma \delta \rangle = \frac{1}{\Omega^2} \delta_{\sigma_{\alpha} \sigma_{\gamma}} \delta_{\sigma_{\beta} \sigma_{\delta}} \underbrace{\int e^{-i\mathbf{R}(\mathbf{k}_{\alpha} - \mathbf{k}_{\delta})} v(\mathbf{R}) d\mathbf{R}}_{v_{\mathbf{k}_{\delta} - \mathbf{k}_{\alpha}}} \underbrace{\int e^{i\mathbf{r}'(\mathbf{k}_{\gamma} - \mathbf{k}_{\beta} + \mathbf{k}_{\delta} - \mathbf{k}_{\alpha})} d\mathbf{r}'}_{=\delta_{\mathbf{k}_{\gamma} - \mathbf{k}_{\beta} + \mathbf{k}_{\delta}, \mathbf{k}_{\alpha}}}.$$

Almost finished, we derive a combuersom equation

$$V = \frac{1}{2\Omega} \sum_{\substack{\mathbf{k}_{\alpha} \mathbf{k}_{\beta} \mathbf{k}_{\gamma} \mathbf{k}_{\delta} \\ \sigma_{\alpha} \sigma_{\beta} \sigma_{\gamma} \sigma_{\delta}}} \delta_{\sigma_{\alpha} \sigma_{\gamma}} \delta_{\sigma_{\beta} \sigma_{\delta}} \delta_{\sigma_{\alpha} \sigma_{\gamma}} \delta_{\sigma_{\beta} \sigma_{\delta}} \delta_{\mathbf{k}_{\alpha}, \mathbf{k}_{\gamma} - \mathbf{k}_{\beta} + \mathbf{k}_{\delta}} v_{\mathbf{k}_{\alpha} - \mathbf{k}_{\delta}} c_{\mathbf{k}_{\alpha} \sigma_{\alpha}}^{\dagger} c_{\mathbf{k}_{\beta} \sigma_{\beta}}^{\dagger} c_{\mathbf{k}_{\gamma} \sigma_{\gamma}} c_{\mathbf{k}_{\delta} \sigma_{\delta}}.$$

we can sum over the \mathbf{k}_{α} , rename $\sigma_{\alpha} \rightarrow \sigma$ and $\sigma_{\beta} \rightarrow \sigma'$ and sum up over σ_{γ} and σ_{δ} to simplify the Kroneker deltas.

$$V = \frac{1}{2\Omega} \sum \sigma \sigma' \sum_{\mathbf{k}_{\beta} \mathbf{k}_{\gamma} \mathbf{k}_{\delta}} v_{\mathbf{k}_{\gamma} - \mathbf{k}_{\beta}} c_{\mathbf{k}_{\delta} + \mathbf{k}_{\gamma} - \mathbf{k}_{\beta}, \sigma}^{\dagger} c_{\mathbf{k}_{\beta} \sigma'}^{\dagger} c_{\mathbf{k}_{\gamma} \sigma'} c_{\mathbf{k}_{\delta} \sigma'}$$

where using $\mathbf{k}_{\alpha} = \mathbf{k}_{\gamma} - \mathbf{k}_{\beta} + \mathbf{k}_{\delta}$ from the Kroneker delta we get $v_{\mathbf{k}_{\alpha} - \mathbf{k}_{\delta}} = v_{\mathbf{k}_{\gamma} - \mathbf{k}_{\beta} + \mathbf{k}_{\delta} - \mathbf{k}_{\delta}}$. Then we can introduce the following variable transformations:

$$\mathbf{k}_{\delta} \rightarrow \mathbf{k}, \quad \mathbf{k}_{\gamma} \rightarrow \mathbf{k}', \quad \mathbf{k}_{\beta} \rightarrow \mathbf{k}' - \mathbf{q}$$

which yields

$$\begin{aligned} \mathbf{k}_{\delta} + \mathbf{k}_{\gamma} - \mathbf{k}_{\beta} &= \mathbf{k} + \mathbf{q} \\ \mathbf{k}_{\gamma} - \mathbf{k}_{\beta} &= -\mathbf{q} \end{aligned}$$

and we finally get our second quantised interaction operator

$$V = \frac{1}{2\Omega} \sum_{\mathbf{q}} v_{\mathbf{q}} \sum_{\substack{\mathbf{k} \sigma \\ \mathbf{k}' \sigma'}} c_{\mathbf{k} + \mathbf{q}, \sigma}^{\dagger} c_{\mathbf{k}' - \mathbf{q}, \sigma'}^{\dagger} c_{\mathbf{k}' \sigma'} c_{\mathbf{k} \sigma}.$$

This describes an electron transferring a momentum \mathbf{q} to another electron. The formula tells that we kill the electron we had before the interaction and create two in states where one loosed some momentum, that has been transfered the other electron. As we saw earlier this depends on the distance $\mathbf{r} - \mathbf{r}'$ between the electrons, but not \mathbf{r} and \mathbf{r}' respectively, so this process is translation invariant. The following diagramm is a good illustration of this effect.




Figure 2: The interaction of two electrons modulated by a photon of momentum \mathbf{q} .

This closes the introduction theorie on the second quantisation. We saw how we can express integral over wavefunctions as bra-kets scalarproducts. This allowed us to introduce some operators that creates and annihilates particles in a certain state. Using this formalism we were able to compute the Hamiltonian of the system in the momentum sapce. We found that the non interacting-part relies on the energy of each state times there occupation number and that an interaction between two electrons can be described as a momentum transfert between them wich is modulated by the fourier transform of the Coulomb potential.

Now we can move on to the descrition of a superconductive state.

2 Superconductivity

Superconductivity can be illustrated as a phase transition of a meterial under a crital temperature. In the superconductive state the material become a perfect diamaget and its resistivity vanishes. We then observe some shielding currents that arise on it's surface and we can let flew a current for a very long time without loosing energy. The superconductive state is also described as Meissner state.

Suppose that we heat the material to the critical temperature T_c , some fluctuation effects arise and break the superconductive state. The shielding effects reacts different on the material. We usely distinguish type I and type II superconductors. The type I superconductor loose abruptly their magnetisation over T_c . Type II have a mixed state where the magnetisation slowly decreases until we can't measure it anymore.

The break of the superconductive state can be described as letting more and more filed flew inside of the material. Asuming that some particle are responsible for the superconductivity, the field achieve to penetrate where wo observe a lower density of these particles. The penetration is described as some magnetic field vortecies reaching a certain depth in the material.

The Meissner state is a thermodynamical state. We can show that the free energy of the superconductive state is higher than the normal state. This results in a lower entropy compered to the normal state.

For readability reason we set the redacted Planck constant $\hbar = 1$ in the following.

2.1 Theoretical framework and BCS theory

The Hamiltonian of the system is discribed by the solid state physics. The consider the energy of the electrons and the ions in a lattice.

$$H = H_{e^-e^-} + H_{e^-ion} + H_{ion ion}$$

Each term consist of a kinetik and potential energy term. For a more mathematical approch we consider a system of N electrons and L ions.

$$\begin{aligned} H_{e^-e^-} &= \sum_{i \in \llbracket N \rrbracket} \frac{p_i^2}{2m} + \sum_{i,j \in \llbracket N \rrbracket} V_{\text{Coulomb}}^{e^-e^-}(\mathbf{r}_i - \mathbf{r}_j) \\ H_{ion ion} &= \sum_{i \in \llbracket M \rrbracket} \frac{p_i^2}{2M} + \sum_{i,j \in \llbracket L \rrbracket} V_{\text{Coulomb}}^{ion-ion}(\mathbf{R}_i - \mathbf{R}_j) \\ H_{e^-ion} &= \sum_{i \in \llbracket N \rrbracket, j \in \llbracket L \rrbracket} V_{\text{Coulomb}}^{e^-ion}(\mathbf{r}_i - \mathbf{R}_j) \end{aligned}$$

We have m and M as the mass of the electron and the ion. \mathbf{r} and \mathbf{R} are the position of the electron and the ion. The ion-ion potential freezes the ions into the lattice. We first going to introduce some concepts by describing a non-interacting electron and then improve it to include the interactions.

2.1.1 The non-interacting electron gas

In this case of study the Hamiltonian only include a kinetic term

$$H = \sum_{\mathbf{k}, \sigma} \epsilon_{\mathbf{k}} c_{\mathbf{k}, \sigma}^{\dagger} c_{\mathbf{k}, \sigma}.$$

We assume that it exist a the ground state $|0\rangle$, where the system is filled up with a certain amount of electron until the Fermi-energy ϵ_F is reached. Associated with this energy we find a wave vector \mathbf{k}_F , the Fermi-momentum. The set of energy up to ϵ_F is called the Fermi-sea, as an analogy to the level zero of the topographic maps.

$$\hat{n}_{\mathbf{k}, \sigma} |0\rangle = \Theta(\epsilon_F - \epsilon_{\mathbf{k}}) |0\rangle.$$

We introduced here a very useful tool called the Heavyside-step function wich is defined as:

$$\Theta(x) = \begin{cases} 1, & x > 0 \\ 0, & x < 0 \end{cases}, \quad \Theta(-x) = 1 - \Theta(x). \quad (10)$$

This means that if we count the particle that have an energy heigher than the Fermi-energy ($\mathbf{k} > \mathbf{k}_F$) than we get zero.

We now want to study how the electron can scatter in different states. The function that we're using is called the propagator and gives the probailty to find the particle at $|\mathbf{k}', \sigma\rangle$ at t' know it at $|\mathbf{k}, \sigma\rangle$, t . An important fact is that without interaction, the particle shouldn't scatter in another state due to energy conservation. Therefore

$$G_0(\mathbf{k}, \mathbf{k}', t' - t) = G_0(\mathbf{k}, t' - t) \delta_{\mathbf{k}, \mathbf{k}}.$$

which is zero if the wave-vectors between the two timepoint differs. We observe that only the past time $t' - t$ is relevant. This is due to the time independent property of the Hamiltonian. We are going to use the representation in the frequency space, using a Fourier-transformation.

$$G_0(\mathbf{k}, \omega) = \int_{\mathbb{R}} e^{i\omega t} G_0(\mathbf{k}, t) dt = \frac{1}{\omega - \epsilon_{\mathbf{k}} + i\delta_{\mathbf{k}}} \quad (11)$$

where $\delta_{\mathbf{k}} = \delta \cdot \text{sgn}(\epsilon_{\mathbf{k}} - \epsilon_F)$ involving a very small non zero number δ . We observe that this analytical function has a pole given by

$$\begin{aligned} \omega - \epsilon_{\mathbf{k}} + i\delta_{\mathbf{k}} &= 0 \\ \iff \omega &= \epsilon_{\mathbf{k}} - i\delta_{\mathbf{k}} \end{aligned}$$

where we denote i as the imaginary unit to avoid confusion with the index i . The frequency ω gives the so called spectrum of the exitation from the unique-particle system. The imaginary part serves as a damping term and is inversly proportional to the lifetime of the particle. δ is a small number due to the infinitely long lifetime. This is a direct result of the absence of scattering.

Further the propagator yields important informations on the system when considering the integration over its different arguments. First we take the imaginary part of the propagator called the single particle spectral weight.

$$\begin{aligned} A(\mathbf{k}, \omega) &= -\frac{1}{\pi} \text{Im} [G_0(\mathbf{k}, \omega)] = \frac{1}{\pi} \frac{\delta_{\mathbf{k}}}{(\omega - \epsilon_{\mathbf{k}})^2 + \delta_{\mathbf{k}}^2} \\ &= \delta(\omega - \epsilon_{\mathbf{k}}) \end{aligned} \quad (12)$$

which informs us about the occupation of a state $|\mathbf{k}\rangle$ with energy ω . We can find a form for the momentum distribution $n(\mathbf{k})$

$$n(\mathbf{k}) = \int A(\mathbf{k}, \omega) d\omega$$

and for the density of state

$$D(\omega) = \int A(\mathbf{k}, \omega) d^3k, \quad \text{or for discontinuous state } \sum_{\mathbf{k}} A(\mathbf{k}, \omega). \quad (13)$$

2.1.2 Fermi-Liquid - the interacting case

Now that we've described the non interacting system, let us complexify the model by introducing the interactions. In an earlier section we saw how

$$H = \sum_{\mathbf{k}, \sigma} \epsilon_{\mathbf{k}} c_{\mathbf{k}, \sigma}^{\dagger} c_{\mathbf{k}, \sigma} + \sum_{\mathbf{k}, \sigma, \mathbf{k}', \sigma'} V_{\mathbf{k}\mathbf{k}', \mathbf{q}} c_{\mathbf{k}-\mathbf{q}, \sigma}^{\dagger} c_{\mathbf{k}+\mathbf{q}, \sigma'}^{\dagger} c_{\mathbf{k}, \sigma} c_{\mathbf{k}', \sigma'} \quad (14)$$

represent the pairwise interaction of multiple electrons and their respective energy.

To extend the model we now want to introduce two new quantities, the propagator G and the one-particle irreducible self-energy Σ . The propagator maps in the complex space and gives the probability amplitude of finding a the particle in the state $|\mathbf{k}, \sigma\rangle$ at a time t . On the other hand $\Sigma = \Sigma_R + i\Sigma_I$ contains the lifetime of the particle in this state and shift of energy of the particle due to the interaction with the surroundings. The framework defines the non-interacting energy of the particle as $\epsilon_{\mathbf{k}}$. When put in an interacting system the spectrum shifts and becomes $\tilde{\epsilon}_{\mathbf{k}} = \epsilon_{\mathbf{k}} + \Sigma_R$. Due to the interactions, the particle then has a much smaller lifetime. Σ_I is antiproportional to the particle's lifetime $\tau_{\mathbf{k}}$. We therefore expect Σ_I to be really small in the non interacting case. These two quantities are linked through the Dyson equation, which reads

$$(G(\mathbf{k}, \omega))^{-1} = (G_0(\mathbf{k}, \omega))^{-1} - \Sigma(\mathbf{k}, \omega).$$

sigmaR?

One can use a Fourier-transformation to switch from the time representation to the frequency representation ω . Reordering the equation and using the result from 11 we obtain

$$G(\mathbf{k}, \omega) = \frac{1}{\omega - \epsilon_{\mathbf{k}} - \Sigma}.$$

This function has a pole at $\omega = \epsilon_{\mathbf{k}} + \Sigma_R + i\Sigma_I$, where in the none interacting case $\omega = \epsilon_{\mathbf{k}} + i\Sigma_I$. It makes sense, the particle spectrum is now shifted due to the finite lifetime of the particle as a result of the interaction.

In complex analysis the order of a pole is given as n if $f(z)$ is meromorphic and has a pole at z_0 where

$$(z - z_0)^n f(z)$$

is also meromorphic in the neighbourhood of z_0 . In our case we're interested in the 0.th order of the pole *why?*. We therefore ignore the imaginary part of the pole and we get

$$\omega = \epsilon_{\mathbf{k}} + \Sigma_R(\mathbf{k}, \epsilon_{\mathbf{k}}) = \tilde{\epsilon}_{\mathbf{k}}.$$

Taking in account a small imaginary part requires a Taylor expansion of Σ_R *why?* in the neighbourhood of $\omega = \tilde{\epsilon}_{\mathbf{k}}$.

Follow with the book and check

We define the residue of the propagator as

$$z_{\mathbf{k}} = \frac{1}{1 - \left. \frac{\partial \Sigma}{\partial \omega} \right|_{\omega = \tilde{\epsilon}_{\mathbf{k}}}} \quad (15)$$

This residue is a decreasing function of the energy, which means that its influence is more important for low energies. An interpretation could be that the slow moving electrons have less time to interact with their homologues. *Interacting electrons are degraded version of the non-interacting case. There remains a $z_{\mathbf{k}} < 1$.*

Quasi-particles The main question we have now is how does the residue look like on the fermi surface? We set us in the context of a low energy electron, close to the Fermi-surface and once we had a interaction. It turns out that if there is a $z_{\mathbf{k}_F} > 0$ we find a precise low energy single particle excitation. This excitation is very close to the exact eigenfunction of a non-interacting Hamiltonian. We call this state akin to the free electron a quasi-particle.

This is it, a Fermi-liquid is a system of interacting electrons and quasi-particles. These quasi-particles are not eigenstates of the interacting Hamiltonian anymore. We can't consider them as an electron like excitation in the interaction context. The interactions allows to scatter some states in and out of the Bloch-State. [Introduce what a Bloch state is.](#)

The above expression for the momentum distribution $n(\mathbf{k})$ can be plotted and we can recognise a gap of $z_{\mathbf{k}}$ or k_F ? wich is an important feature of the Fermi-liquid.

Repulsive interactions In condensed matter physics the dominant effect is the repulsive interactions between the electrons due to the coulomb potential. We already described it in the Hamiltonian H_{e-e} using some pairwise interactions in 14. Therefore the goal is now to find an expression for the potential $V_{\mathbf{k}\mathbf{k}',\mathbf{q}}$.

As we can see in eq. 14 the system is describe in the momentum-space. For this reason we need to consider the Fourier-transform of the real-sapce potential. We start with the coulomb potential which is predominant in the solid state physics. However the integral is going to diverge for $r \rightarrow 0$. To solve this problem We introduce the Yakuma potential which exponentially modulates the coulomb potential:

$$V_\lambda(r) = \frac{1}{4\pi\epsilon_0} \frac{e^2}{r} e^{-r\lambda} \quad (16)$$

this is a solid approximation since we can the fourier transform and let $\lambda \rightarrow 0$ in the result. [Result in appendix?](#). The result we get is

$$V_{\mathbf{k},\mathbf{k}',\mathbf{q}} = V_{\text{el}}(\mathbf{q}) := \frac{1}{4\pi\epsilon_0} \frac{2\pi e^2}{q^2} \quad (17)$$

with \mathbf{q} the momentum transfert during the interaction. The Fermi-liquid remains stable to the repulsive processes.

However as we're going to see in the next section, attractive interactiosn also takes place due to an exchange of phonons between two electrons. This will be the ground stone to our description of the Meissner state.

2.1.3 Instability due to attractive Interactions

[Hier we are going to show how the attractive interactions destabilise the Fermi-liquid:new ground states are open.](#) Now that we showed the influence of attractive interactions, we seek some candidate process that are attractive.

2.1.4 Phonon-mediated attractive interactions [reformulate](#)

As known from condensed matter physics, the lattice can have some intern oscillations called phonons resulting from the spring coupling between the ions. Now we can imagine that due to the Coulomb interactions, an electron can shift an ions producing a phonon. If this phonon travels and influences another electron on it way, we result in an effective electron-electron interaction thanks to the phonon. A similar case would be the exchange of a photon between two electrons. We are going to show how this exchange can lead to an attractive interaction.

Tn a dense lattice the ions moves much slowly around their equilibrium positions than the lights electrons who pass by. The electron moves the charges of the ions resulting in a small dipol moment. A second electron that also passes in the surounding is going to feel the dipol moment and will be attracted. Then the ion shifts back in its new equilibrium position and the dipol moment vanishes long after the first electron passed.

Morover due to the Coulomb interaction, the electrons aim to put as most distance as they can between them in a minimal amount of time. Therefore we can say that the \mathbf{k} -quantum number should be opposite between the two electron. If we target to put these concepts in a mathematical form, we use our previous Hamiltonian and add an electron-phonon interaction term.

$$H = \sum_{\mathbf{k},\sigma} \epsilon_{\mathbf{k}} c_{\mathbf{k},\sigma}^\dagger c_{\mathbf{k},\sigma} + \sum_{\mathbf{k},\sigma,\mathbf{k}',\sigma'} V_{\mathbf{k}\mathbf{k}',\mathbf{q}} c_{\mathbf{k}-\mathbf{q},\sigma}^\dagger c_{\mathbf{k}+\mathbf{q},\sigma'} c_{\mathbf{k},\sigma} c_{\mathbf{k}',\sigma'} + V_{e\text{-phonon}}$$

where $V_{e\text{-phonon}}$ usely depends on the sum of the phonons-modes λ . These modes are similar to the oscillations modes we have in a CO_2 -molecule. The expression form the phonon-depend potential in momentum space reads

$$V_{e\text{-phonon}} = \sum_{\mathbf{k}, \mathbf{q}, \sigma} M_{\mathbf{q}} (a_{-\mathbf{q}\dagger} + a_{\mathbf{q}}) c_{\mathbf{k}+\mathbf{q}, \sigma}^\dagger c_{\mathbf{k}, \sigma}. \quad (18)$$

$M_{\mathbf{q}} (a_{-\mathbf{q}\dagger} + a_{\mathbf{q}})$ is a matrix element of the coupling between the electron and the phonon. The $a_{\mathbf{q}}$ and $a_{\mathbf{q}\dagger}$ are annihilation and creation operators of the phonon with wavevector \mathbf{q} . Further researches have shown that [source](#)

$$[a_{\mathbf{q}}, a_{\mathbf{q}\dagger}] = \delta_{\mathbf{q}, \mathbf{q}'}$$

and therefor phonons act like bosons. Their number is however not conserved in a solid. The matrix element M is a function of the eigenfrequency [or energy?](#) of the phonon $\omega_{\mathbf{q}, \lambda}$ and the fourier transform \tilde{V} of the electrostatic potential $V_{\lambda}(\mathbf{q})$ between the electron and the phonon of mode λ , if included. [more details?](#)

$$M_{\mathbf{q}, \lambda} = i(\mathbf{q} \cdot \boldsymbol{\xi}_{\lambda}) \sqrt{\frac{\hbar}{2M\omega_{\mathbf{q}, \lambda}}} \tilde{V}_{\lambda}(\mathbf{q}).$$

M can't be realy computed due to its complexity, we hold it as a parameter here. This pairing is much waker than the electron-photon interaction. An other important fact ist than for $\mathbf{q} \rightarrow 0$ the matrix element M vanishes. M is proportional to \mathbf{q} which illustrate the the electron-phonon interaction happens between a point charge and a dipol.

The goal is now the implicitly express the phonon exchange with an effective electron-electron process. If we consider two diagrams, one aiming to describe the absorption of a phonon and one the emission of a phonon, we can combine them to get a new effective interaction like the photon exchange case.




Figure 3: The emission (left) and absorption (right) digramm of a phonon of wavevector \mathbf{q} by an electron.

If we represent this nderaction by linking both \mathbf{q} -edges. The energy of phonon is simply defined as

$$H_{\text{phonon}} = \sum_{\mathbf{q}} \omega_{\mathbf{q}} a_{\mathbf{q}}^\dagger a_{\mathbf{q}}$$

and we can write a propagator which describes the dashed line in a connected context similar to to photon case described in the figure ??.

$$D_0(\mathbf{q}, \omega) = \frac{1}{\omega^2 - \omega_{\mathbf{q}}^2 + i\eta}$$

involving a very small quantity η . From this [how?](#) we obtain an expression for the phonon-mediated interaction of two electrons:

$$V_{\text{eff}}^{(ph)}(\mathbf{q}, \omega) = \frac{2|M_{\mathbf{q}}|^2 \omega_{\mathbf{q}}}{\omega^2 - \omega_{\mathbf{q}}^2} \quad (19)$$

with \mathbf{q} the momentum transfert. We can now use a more complete potential in the Hamiltonian involving both the electrostatic 17 and effective phonon-mediated interactions 19.

$$V_{\text{eff}}(\mathbf{k}, \mathbf{k}', \mathbf{q}) = V_{\text{el}}(\mathbf{q}) + V_{\text{eff}}^{(ph)}(\mathbf{q}, \omega) = \frac{1}{4\pi\epsilon_0} \frac{2\pi e^2}{q^2} + \frac{2|M_{\mathbf{q}}|^2 \omega_{\mathbf{q}}}{\omega^2 - \omega_{\mathbf{q}}^2} \quad (20)$$

Here we have reached a very important point. This new potential can be in some case negative, which means we result in an attractive interaction between the electrons. With other word, in some cases the phonon exchange can be attractive and even overcome the strong repulsive coulomb potential. [Graph of V and the simplified version of it, which is also a very good predication.](#)

2.1.5 Contraction of the effective Hamiltonian

We want to restrict ourselves in the case where the effective Hamiltonian is attractive. This happens in a small shell around the Fermi-surface. If we want to maximise the phase space for the scattering, the state before and after the scattering have to be in this shell. A good idea is to consider that the two electrons have opposite wavevectors. The folowing figure illustrates this process.




Figure 4: The scattering process of two electrons with opposite wavevectors \mathbf{k} and \mathbf{k}' . We have a momentum transfert \mathbf{q} between them. The thick line illustrates the Fermi-surface and the dotted one the thin shell. As we see if the electrons have opposite wavevectors, and the \mathbf{k} electrons scatters into the shell, then the \mathbf{k}' electron scatters in the shell as well. This is not allways the case if the wavevectors don't agree as we can see on the left figure. The right figure points out the maximisation process. With opposite wavevectors, we get the largest possibility spectrum for the scattering.

Further the attractivity is a short range effect, so if we want to consider it, we must think that the electrons are very close to eachother. This requires the electrons to have opposite spins due to the Pauli principle. This is a same as a lattice site. This is the only possibility for them to cohabit the same neighbourhood. [check](#) The approximation turns out to be a good model.

Now we allow us to rename some variables:

$$\mathbf{k} + \mathbf{q} \longrightarrow \mathbf{k}, \quad \mathbf{k} \longrightarrow \mathbf{k}'$$

The Hamiltonian that follows from these transformations is called the BCS-reduced Hamiltonian

$$H = \sum_{\mathbf{k}, \sigma} \epsilon_{\mathbf{k}} c_{\mathbf{k}, \sigma}^\dagger c_{\mathbf{k}, \sigma} + \sum_{\mathbf{k}, \sigma, \mathbf{k}'} V_{\mathbf{k}\mathbf{k}'} c_{\mathbf{k}, \sigma}^\dagger c_{-\mathbf{k}, -\sigma}^\dagger c_{-\mathbf{k}', -\sigma} c_{\mathbf{k}', \sigma} \quad (21)$$

$V_{\mathbf{k}\mathbf{k}'}$ is now a matrix element that acts if the wavevectors are close to the Fermi-surface. The electrons have to move in opposite directions with opposite spins. Due to the retardation processes we introduced earlier, there remains a distortion in the lattice long after the electrons passed. Due to the inducing dipol moment, the other electron is attracted towards the distortion with $M_{\mathbf{q}}$?. As we also saw, the coulomb repulsion causes a colinear displacement, close to the distortion of the homologue. This phenomenon is called the Cooper-pairing and is a

coupling that happens in momentum space.

An interacting fact is that these interactions are the source of the superconductivity but they are also to main origin of resistivity in clean materials.

2.1.6 On our way to the BCS-theory

After this introduction on the phonon coupling between the electrons in the momentum space, or Cooper-pairing, we aim to describe the energy of the superconductor in a mean-field approach. The goal of it is to reduce the description with the neighbours to the description of a site, in the mean field of the other sites. Therefore we are going to describe a one-body problem which is easier to compute. As known mean-fields approaches requier self-consistent equations that will as well follow.

The first step is to introduce the following expectation values:

$$b_{\mathbf{k}} = \langle c_{-\mathbf{k}\downarrow} c_{\mathbf{k}\uparrow} \rangle \quad (22)$$

$$b_{\mathbf{k}}^\dagger = \langle c_{\mathbf{k}\uparrow}^\dagger c_{-\mathbf{k}\downarrow}^\dagger \rangle \quad (23)$$

which lead to a new expression for the c operators:

$$c_{-\mathbf{k}\downarrow} c_{\mathbf{k}\uparrow} = b_{\mathbf{k}} + \underbrace{c_{-\mathbf{k}\downarrow} c_{\mathbf{k}\uparrow} - b_{\mathbf{k}}}_{\delta_{b_{\mathbf{k}}}} \quad (24)$$

where we can see the $\delta_{b_{\mathbf{k}}}$ as a deviation, or fluctuation term. If we introduce it back into the Hamiltonian, we can write

$$H = \sum_{\mathbf{k}\sigma} \epsilon_{\mathbf{k}} c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{k}\sigma} + \sum_{\mathbf{k}, \mathbf{k}'} V_{\mathbf{k}\mathbf{k}'} (b_{\mathbf{k}}^\dagger + \delta_{b_{\mathbf{k}}}^\dagger) (b_{\mathbf{k}} + \delta_{b_{\mathbf{k}}}).$$

We can compute the product of the two terms in parenthesis and forget the $\mathcal{O}(\delta_{b_{\mathbf{k}}}^2)$ because the fluctuation are small. We then obtain the following expression

$$H = \sum_{\mathbf{k}\sigma} \epsilon_{\mathbf{k}} c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{k}\sigma} + \sum_{\mathbf{k}, \mathbf{k}'} V_{\mathbf{k}\mathbf{k}'} \left(b_{\mathbf{k}}^\dagger c_{-\mathbf{k}'\downarrow} c_{\mathbf{k}'\uparrow} + b_{\mathbf{k}'}^\dagger c_{\mathbf{k}\uparrow}^\dagger c_{-\mathbf{k}\downarrow}^\dagger - b_{\mathbf{k}}^\dagger b_{\mathbf{k}'} \right).$$

The next step is to define the superconduction gap parameter Δ :

$$\Delta_{\mathbf{k}} := - \sum_{\mathbf{k}'} V_{\mathbf{k}\mathbf{k}'} b_{\mathbf{k}}^\dagger \quad (25)$$

$$\Delta_{\mathbf{k}}^\dagger := - \sum_{\mathbf{k}'} V_{\mathbf{k}\mathbf{k}'} b_{\mathbf{k}'} \quad (26)$$

which brings our Hamiltonian in another form:

$$H = \sum_{\mathbf{k}\sigma} \epsilon_{\mathbf{k}} c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{k}\sigma} - \sum_{\mathbf{k}} \left(\Delta_{\mathbf{k}}^\dagger c_{-\mathbf{k}\downarrow} c_{\mathbf{k}\uparrow} + \Delta_{\mathbf{k}} c_{\mathbf{k}\uparrow}^\dagger c_{-\mathbf{k}\downarrow}^\dagger - b_{\mathbf{k}}^\dagger \Delta_{\mathbf{k}}^\dagger \right). \quad (27)$$

We took the liberty to split the sum, rename the \mathbf{k}' to \mathbf{k} and recombine the sum. We notice that this form involves a lot of creation and annihilation terms that are not common for an effective non-interacting electron gas. We remember that we aim at a one-particle description in a mean field of its neighbours. This complexity will lead to some difficulties to express the quasi-particle spectrum. A good solution is to rotate the basis of the c operators to land in a basis that diagonalises the Hamiltonian and therefore minimises the number of operators.

The transformation involves two new fermionic operators η and γ that therefore respect \mathfrak{fer}_1 to \mathfrak{fer}_3 and reads

$$\begin{pmatrix} c_{\mathbf{k}\uparrow} \\ c_{-\mathbf{k}\downarrow}^\dagger \end{pmatrix} = \begin{pmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{pmatrix} \begin{pmatrix} \eta_{\mathbf{k}} \\ \gamma_{\mathbf{k}} \end{pmatrix} \quad (28)$$

along with the conjugate transpose of each component of the l.h.s vector rebuild into a matrix-vector equation

$$\begin{pmatrix} c_{\mathbf{k}\uparrow}^\dagger \\ c_{-\mathbf{k}\downarrow} \end{pmatrix} = \begin{pmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{pmatrix} \begin{pmatrix} \eta_{\mathbf{k}}^\dagger \\ \gamma_{\mathbf{k}}^\dagger \end{pmatrix}. \quad (29)$$

We can reintroduce these into the Hamiltonian 27. Some of the multiplication involves $\cos(\theta)^2 - \sin(\theta)^2 = \cos(2\theta)$ and $\cos(\theta)^2 + \sin(\theta)^2 = 1$. Further due to the anticommutations we have $\eta\gamma^\dagger = -\gamma^\dagger\eta$ and so on for $\gamma\eta^\dagger = -\eta^\dagger\gamma$. We obtain the following expression

$$\begin{aligned}
H = & \sum_{\mathbf{k}} \epsilon_{\mathbf{k}} \cdot 1 + \Delta_{\mathbf{k}} b_{\mathbf{k}}^\dagger \\
& + \sum_{\mathbf{k}} \left[\epsilon_{\mathbf{k}} \cos(2\theta) - \cos(\theta) \sin(\theta) \left(\Delta_{\mathbf{k}}^\dagger + \Delta_{\mathbf{k}} \right) \right] \eta_{\mathbf{k}}^\dagger \eta_{\mathbf{k}} \\
& - \sum_{\mathbf{k}} \left[\epsilon_{\mathbf{k}} \cos(2\theta) - \sin(\theta) \cos(\theta) \left(\Delta_{\mathbf{k}}^\dagger + \Delta_{\mathbf{k}} \right) \right] \gamma_{\mathbf{k}}^\dagger \eta_{\mathbf{k}} \\
& - \sum_{\mathbf{k}} \left[\Delta_{\mathbf{k}} \cos(\theta)^2 - \Delta_{\mathbf{k}}^\dagger \sin(\theta)^2 + 2\epsilon_{\mathbf{k}} \cos(\theta) \sin(\theta) \right] \eta_{\mathbf{k}}^\dagger \gamma_{\mathbf{k}} \\
& - \sum_{\mathbf{k}} \left[\Delta_{\mathbf{k}} \cos(\theta)^2 - \Delta_{\mathbf{k}}^\dagger \sin(\theta)^2 + 2\epsilon_{\mathbf{k}} \cos(\theta) \sin(\theta) \right] \gamma_{\mathbf{k}}^\dagger \eta_{\mathbf{k}}.
\end{aligned} \tag{30}$$

The goal is to diagonalise the Hamiltonian in the (γ, η) basis. Therefore have to rotate with θ such that the terms with $\gamma^\dagger\eta$ and $\eta^\dagger\gamma$ vanish. A difficulty that we may encounter along with this idea is that Δ is an order parameter and own a complex phase fluctuation. With other words one can write $\Delta = |\Delta|e^{i\tau}$ where τ has some fluctuations. We are going to ignore them.

We can set

$$\Delta_{\mathbf{k}} = \Delta_{\mathbf{k}}^\dagger \quad \text{and} \quad \tan(2\theta) = -\frac{\Delta_{\mathbf{k}}}{\epsilon_{\mathbf{k}}}$$

and introduce two new variables called the coherence factors

$$v_{\mathbf{k}}^2 := \sin(\theta)^2 = \frac{1}{2} \left(1 - \frac{\epsilon_{\mathbf{k}}}{E_{\mathbf{k}}} \right), \tag{31}$$

$$u_{\mathbf{k}}^2 := \cos(\theta)^2 = \frac{1}{2} \left(1 + \frac{\epsilon_{\mathbf{k}}}{E_{\mathbf{k}}} \right) \tag{32}$$

along with a new energy $E_{\mathbf{k}} = \sqrt{\epsilon_{\mathbf{k}}^2 + |\Delta_{\mathbf{k}}|^2}$. [Historical context?](#) These factors play an important role in NMR as well as in the ultra sound propagation in superconductors. Cooper and Schrieffer made some correct predictions in an advanced many-body system. This is labeled as one of the greatest achievements in condensed matter physics in the 20th century.

[source](#)

We obtain a Hamiltonian that looks very similar to a free fermion quasiparticle gas [source?](#)

$$\begin{aligned}
H = & \underbrace{\sum_{\mathbf{k}} \left(\epsilon_{\mathbf{k}} + \Delta_{\mathbf{k}} b_{\mathbf{k}}^\dagger \right)}_{=: H_0 \text{ constant mean-field term}} + \underbrace{\sum_{\mathbf{k}} E_{\mathbf{k}} \left(\eta_{\mathbf{k}}^\dagger \eta_{\mathbf{k}} - \gamma_{\mathbf{k}}^\dagger \gamma_{\mathbf{k}} \right)}_{\text{Spinless fermion system with two types of fermions of energies } E_{\mathbf{k}} \text{ and } -E_{\mathbf{k}}}.
\end{aligned} \tag{33}$$

where we don't describe any interaction, but two types of particles in a mean field. As [1] described it in chapter 3, p.83-84, one can note that we don't have any spins involved. This is due to the fact that we describe the quasiparticles as a linear combination of electrons and holes with opposite spins. [source](#) Therefore talking about degrees of freedom the two electron-hole singlets replace the \uparrow, \downarrow degrees of freedom and they are preserved.

The energy $E_{\mathbf{k}}$ shows up a gap in the energy of the quasiparticle when looking at the fermi-surface $\epsilon_{\mathbf{k}} = 0$ [We count relative to it? But the way it is defined in a non interacting Ham. is not relative to it right?](#). This is due to the fact that the expectation values we introduced in 22 and 23 are not zero. The gap is a result of the Cooper-pairing. These expectation values are the order parameters of the Meissner state and should be confused with the superconducting gap Δ . They are however zero at the same time.

This context of a free case is a good opportunity to introduce the grand canonical ensemble of the Hamiltonian. Even if we could give the direct result, deriving this ensemble involves a lot of important steps, so for the sake of completeness we are going to derive it.

We define the particle number operator $N = \sum_{\mathbf{k}} \mu (n_{\eta\mathbf{k}} + n_{\gamma\mathbf{k}})$ where the index \mathbf{k} is proper to this equation. Further the possible states \mathbf{k} are in a “continuous” set $\mathfrak{K} = \{\mathbf{k}_1, \mathbf{k}_2, \dots\}$. $\{n_{\mathbf{k}}\}$ is the set of the different occupation numbers of all the states \mathbf{k} . Further we consider fermions so the occupation numbers of the particle type η or γ in state \mathbf{k} are labeled $n_{\eta\mathbf{k}}$ and $n_{\gamma\mathbf{k}}$ and equals either 0 or 1.

$$\begin{aligned}
Z_G &= \text{Tr} \left(e^{-\beta(H+\mu N)} \right) \\
&= \sum_{\{n_{\mathbf{k}}\}} e^{-\beta H_0} \langle \{n_{\mathbf{k}}\} | \exp \left(\sum_{\mathbf{k}'} -\beta E_{\mathbf{k}'} \eta_{\mathbf{k}'}^\dagger \eta_{\mathbf{k}'} + \beta E_{\mathbf{k}'} \gamma_{\mathbf{k}'}^\dagger \gamma_{\mathbf{k}'} \right) \exp(-\beta \mu N) | \{n_{\mathbf{k}}\} \rangle \\
&= \sum_{\{n_{\eta\mathbf{k}}\}} \sum_{\{n_{\gamma\mathbf{k}}\}} e^{-\beta H_0} \exp \left(\sum_{\mathbf{k}} -\beta \left[E_{\mathbf{k}} \eta_{\mathbf{k}}^\dagger \eta_{\mathbf{k}} + \mu \right] + \beta \left[E_{\mathbf{k}} \gamma_{\mathbf{k}}^\dagger \gamma_{\mathbf{k}} - \mu \right] \right) \\
&= e^{-\beta H_0} \sum_{\substack{n_{\eta\mathbf{k}_1}, \\ n_{\eta\mathbf{k}_2}, \dots}} \sum_{\substack{n_{\gamma\mathbf{k}_1}, \\ n_{\gamma\mathbf{k}_2}, \dots}} \exp \left(\sum_{\mathbf{k}} -\beta \left[E_{\mathbf{k}} n_{\eta\mathbf{k}} + \mu \right] + \beta \left[E_{\mathbf{k}} n_{\gamma\mathbf{k}} - \mu \right] \right) \\
&= e^{-\beta H_0} \sum_{\substack{n_{\eta\mathbf{k}_1}, \\ n_{\eta\mathbf{k}_2}, \dots}} \prod_{\mathbf{k}} \exp(-\beta [E_{\mathbf{k}} n_{\eta\mathbf{k}} + \mu]) \sum_{\substack{n_{\gamma\mathbf{k}_1}, \\ n_{\gamma\mathbf{k}_2}, \dots}} \prod_{\mathbf{k}} \exp(\beta [E_{\mathbf{k}} n_{\gamma\mathbf{k}} - \mu]) \\
&= e^{-\beta H_0} \sum_{n_{\eta\mathbf{k}_1}} \exp(-\beta [E_{\mathbf{k}_1} n_{\eta\mathbf{k}_1} + \mu]) \sum_{n_{\eta\mathbf{k}_2}} \exp(-\beta [E_{\mathbf{k}_2} n_{\eta\mathbf{k}_2} + \mu]) \dots \\
&\quad \sum_{n_{\gamma\mathbf{k}_1}} \exp(\beta [E_{\mathbf{k}_1} n_{\gamma\mathbf{k}_1} - \mu]) \sum_{n_{\gamma\mathbf{k}_2}} \exp(\beta [E_{\mathbf{k}_2} n_{\gamma\mathbf{k}_2} - \mu]) \dots \\
&= e^{-\beta H_0} \prod_{\mathbf{k}} \left(1 + e^{-\beta(E_{\mathbf{k}} + \mu)} \right) \left(1 + e^{\beta(E_{\mathbf{k}} - \mu)} \right)
\end{aligned}$$

This partition function isn't very useful like this in our case. But we can derive the free energy which will help us to solve the self consistency equation for the gap. According to [2] p.99 the free energy can be derived from the partition function as ([Appendix for more details?](#))

$$F = -\frac{1}{\beta} \ln(Z_G) = H_0 - \frac{1}{\beta} \sum_{\mathbf{k}} + \ln(1 + e^{-\beta E_{\mathbf{k}}}) \ln(1 + e^{\beta E_{\mathbf{k}}})$$

According to [1] we obtain in the limit of low temperatures ($\beta \rightarrow \infty$)

$$\begin{aligned}
F &= H_0 + \sum_{\mathbf{k}} E_{\mathbf{k}} \theta(-E_{\mathbf{k}}) + E_{\mathbf{k}} \theta(E_{\mathbf{k}}) \\
&= H_0 + \sum_{\mathbf{k}} E_{\mathbf{k}} - E_{\mathbf{k}} \theta(E_{\mathbf{k}}) + E_{\mathbf{k}} \theta(E_{\mathbf{k}}) \\
&= \sum_{\mathbf{k}} \epsilon_{\mathbf{k}} + \Delta_{\mathbf{k}} b_{\mathbf{k}}^\dagger E_{\mathbf{k}}.
\end{aligned}$$

where we used the properties from the Heavyside step-function described in 10. The minimization of the free energy F should self-consistently determine the gap $\Delta_{\mathbf{k}}$. This is a statement that depends neither on the momentum space structure nor on the attractiveness of the potential [1]. $E_{\mathbf{k}}$ has an implicit $\Delta_{\mathbf{k}}$ -dependence. We compute the derivation $\frac{\partial F}{\partial \Delta_{\mathbf{k}}}$ and search the argument of the zero-postion. Derivating one \mathbf{k} from the sum is enough. We demand the following to be satisfied:

$$\frac{\partial F}{\partial \Delta_{\mathbf{k}}} = 0, \quad \frac{\partial F}{\partial \Delta_{\mathbf{k}}^\dagger} = 0 \quad (34)$$

This leads to [Appendix?](#)

$$b_{\mathbf{k}}^\dagger = \Delta_{\mathbf{k}} \underbrace{\frac{\tanh(\beta E_{\mathbf{k}}/2)}{E_{\mathbf{k}}}}_{\chi(\mathbf{k})}$$

$\chi(\mathbf{k})$ is the pair- susceptibility and gives how capable the system is to create Cooper-pairs [1]. If we allow us to relabel $\mathbf{k} \rightarrow \mathbf{k}'$ then multiply on both sides with $-\sum_{\mathbf{k}} V_{\mathbf{k}\mathbf{k}'}$ and introduce 26

we obtain the self-consistency equation for the gap. This is usually denominated as the BCS gap equation.

$$\Delta_{\mathbf{k}} = - \sum_{\mathbf{k}'} V_{\mathbf{k}\mathbf{k}'} \Delta_{\mathbf{k}'} \frac{\tanh(\beta E_{\mathbf{k}'}/2)}{E_{\mathbf{k}'}}. \quad (35)$$

Fossheim and Sudbø [1] emphasize that this equation is not limited to phonon mediated interaction, as long we don't specify the potential. Further Cooper-pairs can condensate because in the pair the electron involved have opposite spins and this adds up to zero. The pair is not ruled anymore by the Pauli principle and can take part into the superconducting condensate. Even if the formation temperature of Cooper-pairs is higher than the critical temperature T_C , this mean field approach makes this temperatures agree.

As we introduced earlier the gap has a complex phase fluctuation. This phase is hard to vary in good metals because the number of carrier electrons is high. On the other side in poor metals the fluctuations that break the pair are more easily reached, i.e. at a lower temperature than in good metal. Superconductivity is therefore more stable in metals of good value.

In this section we saw how to reduce the many body Hamiltonian to a single-body problem in the mean field of its neighbours. The system that follows from this is a non interacting gas of a type electrons and holes. This involves the introduction of the superconducting gap parameter that we can thanks to the statistical mechanics formalism express in a self-consistent way.

2.1.7 Transition temperature and energy gap

The goal is to derive a universal ratio between Δ and the critical temperature.

2.1.8 Generalized gap equation, s-wave and d-wave superconductivity


As we introduced in an earlier section, the formalism follows from the attractive phonon exchange but can be generalized. The superconductivity is the result of the pairing of the electrons into Cooper-pairs. These pairs shift into a condensate and take part of a coherent matter-wave. Making the analogy with the coherent light wave, superconductivity can be viewed as the *analog* of lasers like Fossheim and Sudbø highlight it in [1].

To achieve such generalisation we let the formalism be open to other kind of interactions and don't restrict it to the thin shell around the Fermi-surface. The potential matrix-element can therefore take a complex form than introduced in 20.

We recall that the electrons move into a lattice. We can assume that this crystal owns some symmetries that are reflected in its crystallographic (complete) basis $\{g_\eta(\mathbf{k})\}$. Similar to the Bloch state, we could imagine that the symmetries emphasize some physical quantities. Assuming this we propose a form for the potential:

$$V_{\mathbf{k}\mathbf{k}'} = \sum_{\eta} \lambda_{\eta} g_{\eta}(\mathbf{k}) g_{\eta}(\mathbf{k}')$$

η is also often labeled as the channel [1]. Assuming we have a square lattice,



with interaction strength $U/2$ on site, for nearest neighbours $2V$ and $4W$ for second-nearest neighbours. [need of more details from Jacob](#). Taking the Fourier transform of this leads:

$$\begin{aligned} f(k, k') &= \cos(k) \cos(k') + \sin(k) \sin(k') \\ V_{\mathbf{k}\mathbf{k}'} &= U + 2V (f(k_x, k'_x) + f(k_y, k'_y)) \\ &\quad + 4W (f(k_x, k'_x) \cdot f(k_y, k'_y)) \end{aligned}$$

We can then introduce the basis-functions $\{g_\eta(\mathbf{k})\}$ we talked about a few paragraphs ago.

$$\begin{aligned}
g_1(\mathbf{k}) &= \frac{1}{2\pi} \\
g_2(\mathbf{k}) &= \frac{1}{2\pi} (\cos(k_x) + \cos(k_y)) & \text{(s-wave)} \\
g_3(\mathbf{k}) &= \frac{1}{2\pi} \cos(k_x) \cos(k_y) \\
g_4(\mathbf{k}) &= \frac{1}{2\pi} (\cos(k_x) - \cos(k_y)) & \text{(d-wave)} \\
g_5(\mathbf{k}) &= \frac{1}{2\pi} \sin(k_x) \sin(k_y)
\end{aligned}$$

from which involves the following lambdas $\lambda_1 = 2U\pi^2$, $\lambda_2 = \lambda_4 = 4V\pi^2$, $\lambda_3 = \lambda_5 = 4W\pi^2$ and for greater η ge set $\lambda_{\eta \geq 6} = 0$.

For $\eta \in \{1, 2\}$, the corresponding g is the identity under all symmetries of C_{v4} and for $\eta \in \{3, 4, 5\}$ we observe a change of sign under $\pi/2$ rotations. The same behaviour is found in the spherical harmonics for resp. the quantum numbers $l = 0$ and $l = 2$. From this we can deduct the use of the terms s-wave and d-wave. [Does this need a source?](#)

Furhter we can reintroduce the BCS gap equation 35 and obtain a lattice-dependent form for the gap:

$$\begin{aligned}
\Delta_{\mathbf{k}} &= - \sum_{\mathbf{k}'} V_{\mathbf{k}\mathbf{k}'} \Delta_{\mathbf{k}'} \chi(E_{\mathbf{k}}) \\
&= - \sum_{\eta \in \llbracket 5 \rrbracket} \lambda_\eta g_\eta(\mathbf{k}) \underbrace{\sum_{\mathbf{k}'} g_\eta(\mathbf{k}') \Delta_{\mathbf{k}'} \chi(E_{\mathbf{k}'})}_{=: \Delta_\eta / \lambda_\eta} \\
&= \sum_{\eta \in \llbracket 5 \rrbracket} \Delta_\eta g_\eta(\mathbf{k})
\end{aligned} \tag{36}$$

This is a nice form to have as Fossheim and Sudbø [1] point out p.92. The gap is a linear combination of the physical quantites $g_\eta(\mathbf{k})$ which gives a physical quantities as well. The newly introduced Δ_η are independent from the wavevector \mathbf{k} but are function if the temperature. We can express them in the basis of antoher η' :

$$\begin{aligned}
\Delta_\eta &= \sum_{\eta' \in \llbracket 5 \rrbracket} \Delta_{\eta'} \mathcal{M}_{\eta, \eta'} \\
\mathcal{M}_{\eta, \eta'} &= - \lambda_\eta \sum_{\mathbf{k}} g_\eta(\mathbf{k}) g_{\eta'}(\mathbf{k}) \chi(E_{\mathbf{k}}).
\end{aligned}$$

These nummericaly combuersom to compiute but taking $U > 0$, $V < 0$ and $W = 0$ in the square lattice we obtain no attraction $V_{\mathbf{k}\mathbf{k}'} \eta=1 > 0$? in the λ_1 -channel but attraction in the s- and d-wave channels. This tells us a lot like [1] analised it. First the gap is highly associated to the Fourier transform of the wavefunction of the Cooper-pairss [Which variable describes their wavefunction?](#) Further the gap accomodate itself to be zero in the channel where the wavefuction is non-zero [“for zero separation between the electrons”](#). The s- and d-wave channels are prefered which as consequence avoid the on site coulomb repulsion (i.e the Coulomb froce between the two members of the pair). This can be put in comparasion with the special case of the phonon pairing. In the latest the electrons used retardation processes to cancel out the repulsion. They avoided themselves in time. Here the avoidance takes place in space. ([How to recognise the angular momentum coupeling here?](#))

The manifestation of the retardation effects are acctually a direct consequence of restricting ourselves to a thin shell ouround the Fermi-surface. [But members of a Cooper pairs must have oposite spin and momenta. The opposite mometum is a result of the shell so is this statement valid in general?](#). This means without restrictions, the whole Brillouin zone is taken into account to compute the gap.

Further we can express the Fourier transform of the gap as $\Delta(\mathbf{r}) = \sum_{\mathbf{k}} \Delta_{\mathbf{k}} e^{i\mathbf{k}\mathbf{r}}$ which leads $\Delta(0) = \sum_{\mathbf{k}} \Delta_{\mathbf{k}}$. However a repulsive interaction is obtained for $\Delta(0) \neq 0$ so we need to find $\Delta_{\mathbf{k}}$ s whose sum satisfy this. This can be done using the s-wave and d-wave channels, i.e $\Delta_{\mathbf{k}} = \Delta_0(T)g_2(\mathbf{k})$ and $\Delta_{\mathbf{k}} = \Delta_0(T)g_4(\mathbf{k})$. We already motivated superconductivity and its difficulty to maintain due to the phase fluctuations above the (freezing) critical temperature T_c . However superconductivity involving d-wave channels is believed [1] p.92 to be found in materials with high T_c . $\Delta_{\mathbf{k}} = \Delta_0(T)g_4(\mathbf{k})$ is therefore a good case of study.

Density of Meissner states

We imagine some fluctuation $g(\mathbf{k})$ that only depend on $E_{\mathbf{k}} = \sqrt{\epsilon_{\mathbf{k}}^2 + \Delta_{\mathbf{k}}^2}$. Further we assume Δ being \mathbf{k} -independent for now. We can introduce the normal density of states.

$$D_n(\epsilon) = \frac{1}{N} \sum_{\mathbf{k}} \delta(\epsilon - \epsilon_{\mathbf{k}})$$

with N the number of Fourier modes. This is the same as the number of lattice sites for us [1] p.93.

$$\sum_{\mathbf{k}} g(\mathbf{k}) = \int D_n(\epsilon) g(\sqrt{\epsilon^2 - \Delta^2}) d\epsilon$$

as well as a variable transformation $E = \sqrt{\epsilon^2 + \Delta^2}$ which leads to

$$dE = \frac{\epsilon}{\sqrt{\epsilon^2 + \Delta^2}} d\epsilon \iff d\epsilon = \frac{E}{\epsilon} dE,$$

$$\sum_{\mathbf{k}} g(\mathbf{k}) = \int D_n(\epsilon) g(E) \frac{E}{\epsilon} dE.$$

Further we have $\frac{\partial \epsilon}{\partial E} = \frac{E}{\sqrt{E^2 - \Delta^2}} = \frac{E}{\epsilon}$ so that

$$\sum_{\mathbf{k}} g(\mathbf{k}) = \int g(E) \underbrace{D_n(\epsilon) \frac{\partial \epsilon}{\partial E}}_{D_s(E)} dE. \quad (\text{take total or partial derivative?})$$

Then we use a methode we already used (sure?) to simplify the expression. We assume that $D_n(\epsilon)$ is slowly varying so that $D_n(\epsilon) \approx D_n(0)$ where ϵ is measured relative to the Fermi-surface. Hopefully we don't have any sharpness around the Fermi-surface [1] p.93. We can therefore exclude the Van Hove singularities close to the Fermi-surface.

$$\frac{D_s(E)}{D_n(0)} \frac{E}{\sqrt{E^2 - \Delta^2}} = \frac{E}{\sqrt{E^2 - \Delta^2}} \underbrace{\frac{D_n(\epsilon)}{D_n(0)}}_{=\Theta(E-|\Delta|)}.$$

For the \mathbf{k} dependence of the gap we can use the spectral weight introduced in 12. In analogy to the propagator or Green function introduced in ?? for the electrons, we can define one for the superconducting condensate that involves time-dependent fermionic operators.

$$G(\mathbf{k}, t; \sigma) = -i \langle 0 | c_{\mathbf{k}\sigma}(t) c_{\mathbf{k}\sigma}^\dagger(0) | 0 \rangle$$

involving the BCS superconducting ground state $|0\rangle$ which is an eigenstate of 33. As before we can consider the Fourier transform of the propagator $G(\mathbf{k}, \omega; \sigma)$

$$G(\mathbf{k}, \omega; \sigma) = \frac{1}{2\pi} \int e^{i\omega t} G(\mathbf{k}, \omega; \sigma) dt$$

We recall that Cooper pairs are two made of two electrons with opposite spin. We can first take a look at the density of state of the pair-member with $\sigma = \uparrow$. However are only considering spin singlet pairing so proving that the spectral weight A is spin independent should show that computing it for $\sigma = \uparrow$ is enough as Fossheim and Sudbø argued in [1] p.94.

So without loss of generality we set $\sigma = \uparrow$ and reintroduce the rotation of the basis of the c operators we performed in 28 and 29.

$$G(\mathbf{k}, t; \uparrow) = -i \langle 0 | \left(\cos(\theta) \eta_{\mathbf{k}}^\dagger(t) - \sin(\theta) \gamma_{\mathbf{k}}^\dagger(t) \right) \cdot \left(\cos(\theta) \eta_{\mathbf{k}}(0) - \sin(\theta) \gamma_{\mathbf{k}}(0) \right) | 0 \rangle$$

The goal of these rotated operators was to diagonalise the hamiltonian. It follows

$$\langle 0 | \eta_{\mathbf{k}}^\dagger \gamma_{\mathbf{k}}^\dagger | 0 \rangle = 0 = \langle 0 | \eta_{\mathbf{k}} \gamma_{\mathbf{k}} | 0 \rangle$$

and therefore

$$G(\mathbf{k}, t; \uparrow) = -i \cos(\theta)^2 \langle 0 | \eta_{\mathbf{k}}^\dagger(t) \eta_{\mathbf{k}}(0) | 0 \rangle - i \sin(\theta)^2 \langle 0 | \eta_{\mathbf{k}}^\dagger(t) \eta_{\mathbf{k}}(0) | 0 \rangle. \quad (37)$$

We can nicely split the propagator in the sum of free η - and γ -particles in the superconducting state. Employing to coherence factors 32 and 31 in the Fourier transform of we obtain

$$G(\mathbf{k}, \omega; \uparrow) = \frac{u_{\mathbf{k}}^2}{\omega - E_{\mathbf{k}} + i\delta_{\mathbf{k}}} + \frac{v_{\mathbf{k}}^2}{\omega + E_{\mathbf{k}} - i\delta_{\mathbf{k}}}.$$

which gives the spectral weight

$$A(\mathbf{k}, \omega; \uparrow) = u_{\mathbf{k}}^2 \delta(\omega - E_{\mathbf{k}}) + v_{\mathbf{k}}^2 \delta(\omega + E_{\mathbf{k}}).$$

This value is spin independent and using the arguments we've just given in such a case, we find

$$\begin{aligned} D_s(\omega) &= \frac{1}{N} \sum_{\mathbf{k}} A(\mathbf{k}, \omega) \\ &= \frac{1}{N} \sum_{\mathbf{k}} (u_{\mathbf{k}}^2 \delta(\omega - E_{\mathbf{k}}) + v_{\mathbf{k}}^2 \delta(\omega + E_{\mathbf{k}})). \end{aligned}$$

The density of state of the superconductive condensate D_s is also spin independent since we're considering spin-singlet pairings.

3 Bogoliubov-de Gennes Formalism

The Bogoliubov-de Gennes transformation allows us to express the hamiltonian in a diagonal way and express our quantities by looking at the eigenvectors of the hamiltonian. The resulting matrix is expressed in a huge space and is very sparse.

To give a taste of it, it will allow us to rewrite our hamiltonian as following

$$H = E_0 - \frac{1}{2} \tilde{c}^\dagger \tilde{H} \tilde{c}, \quad (38)$$

involving $\tilde{c} = (\tilde{c}_1, \dots, \tilde{c}_N)$, where each \tilde{c}_i is a vector containing the creation and annihilation operators of a lattice site i : $\tilde{c}_i = (c_{i,\uparrow}, c_{i,\downarrow}, c_{i,\uparrow}^\dagger, c_{i,\downarrow}^\dagger)$.

As we see we just describe each site with the four possible c -operators. This means for each lattice site, we have a 4×4 -submatrix that reflects the possible combinations of creation and annihilation operators of both spins. For the readability we are going to drop the comma between the site and spin indices.

For exemple if one has (without loss of generality) a chemical potential at the site i , then the hamiltonian is described in the following way:

$$H_{\text{chem},i} = \sum_{\sigma} \mu_i c_{i,\sigma}^\dagger c_{i,\sigma}$$

If we want to describe it in terms of \tilde{c}_i we have:

$$H_{\text{chem},i} = \tilde{c}_i^\dagger \cdot \mu_i \mathbb{I}_4 \cdot \tilde{c}_i = \begin{pmatrix} c_{i,\uparrow}^\dagger \\ c_{i,\downarrow}^\dagger \\ c_{i,\uparrow} \\ c_{i,\downarrow} \end{pmatrix} \cdot \mu_i \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} c_{i,\uparrow} \\ c_{i,\downarrow} \\ c_{i,\uparrow}^\dagger \\ c_{i,\downarrow}^\dagger \end{pmatrix}$$

Depending on the interaction we wish to describe, we can figure out what combination of operators we want and design the 4×4 matrix accordingly. To achieve a full description of the system we can consider the interaction between site i, j as a 4×4 matrix involving the \hat{c}_i^\dagger and c_j operators. Then we can build a huge matrix \tilde{H} based on 4×4 matrices at $\tilde{H}_{i,j}$ and the vector we multiply it to is just the \hat{c}_i^\dagger and c_j operators stacked above one another forming the above-introduced \tilde{c} vector. As a result, one gets the first formula introduced in this section 38. We can then compute the eigenvalues and -vectors express the quantities we're interested in. This is what we call the Bogoliubov-de Gennes transformation.

Now that the motivation is clear, we need to bring our Hamiltonian in a form that involves the fermionic operators $c_{i\sigma}$ and $c_{i\sigma}^\dagger$.

3.1 Tigh Binding Model

Our goal is now to fix our particle on lattice sites and describe their interactions. We are therefore going to translate our wavefunction formalism in an on site plus nearest neighbour description.

For the generalities, assume we have the Hamiltonian in the second quantisation formalism:

$$H = \sum_{\sigma\sigma'} \int \phi_\sigma^\dagger(\mathbf{r}) H_{\sigma\sigma'}(\mathbf{r}) \psi_{\sigma'}(\mathbf{r}) d^3r \\ + \sum_{\sigma\sigma'} \int \int \phi_\sigma^\dagger(\mathbf{r}) \phi_{\sigma'}^\dagger(\mathbf{r}') V_{\sigma\sigma'}(\mathbf{r}, \mathbf{r}') \phi_{\sigma'}(\mathbf{r}') \phi_\sigma(\mathbf{r}) d^3r' d^3r$$

We introduce a basis of so called Wannier orbitals $w(\mathbf{r} - \mathbf{R}_i)$ with \mathbf{R}_i an atom location. The should be large in the neighbourhood of \mathbf{R}_i and vanishes when the distance tends to infinity. They are therefore called "localised". The basis is complete, the orbitals verify the orthonormality condition:

$$\int w^*(\mathbf{r} - \mathbf{R}_i) w(\mathbf{r} - \mathbf{R}_j) d^3r = \delta_{ij}.$$

therefore we can define some field operator in this basis, based on creation and annihilation operators that acts on a lattice site i :

$$\phi_\sigma(\mathbf{r}) := \sum_i w(\mathbf{r} - \mathbf{R}_i) c_{i\sigma} \quad \phi_\sigma^\dagger(\mathbf{r}) := \sum_i w^*(\mathbf{r} - \mathbf{R}_i) c_{i\sigma}^\dagger \quad (39)$$

which is not a continuous description anymore. Inserting these operator back into our above Hamiltonian and using the orthonormality allows us to have an on site/nearest neighbour Hamiltonian. Taking for instance the first part of the Hamiltonian:

$$H = \sum_{\sigma\sigma'} \int \psi_\sigma^\dagger(\mathbf{r}) H_{\sigma\sigma'}(\mathbf{r}) \psi_{\sigma'}(\mathbf{r}) d^3r \\ = \sum_{ij\sigma\sigma'} c_{i\sigma}^\dagger c_{j\sigma'} \int w^*(\mathbf{r} - \mathbf{R}_i) H_{\sigma\sigma'}(\mathbf{r}) w(\mathbf{r} - \mathbf{R}_j) d^3r \\ := \sum_{i\sigma\sigma'} \epsilon_i^{\sigma\sigma'} c_{i\sigma}^\dagger c_{i\sigma} - \sum_{\langle ij \rangle \sigma\sigma'} t_{ij}^{\sigma\sigma'} c_{i\sigma}^\dagger c_{j\sigma'} + \dots$$

In the last line we include a local energy term ϵ and the so called hopping term t_{ij} , which is the interaction with the nearest neighbour sites j of i . For a more precise description one could consider more neighbour. The spin dependent term can be used to describe spin orbit coupling or spin-flip processes.

We now aim to define the useful process for this thesis using this formalism.

3.1.1 Non-interacting electrons

The two main components of the non-interacting system Hamiltonian H_N are the chemical potential μ_i which is specific to each site and the hopping term t_{ij} . The chemical potential is modulated by the number of particles on the site i and the hopping term gives the amplitudes

of moving a electron from site i to j . We assume it as spin-independent here.

$$H_N = - \sum_{i\sigma} \mu_i c_{i\sigma}^\dagger c_{i\sigma} - \sum_{\langle ij \rangle \sigma} t_{ij} c_{i\sigma}^\dagger c_{j\sigma} \quad (40)$$

where $\langle ij \rangle$ is a commonly-used notation to sum over i and its nearest neighbours j , skipping $i = j$. We label it the normal Hamiltonian.

The hopping amplitude can be computed from the overlap of the orbitals under a kinetic operator $-\nabla^2/(2m)$, which explains the meaning ‘‘hopping’’:

$$\begin{aligned} t_{ij} &= - \int w^*(\mathbf{r} - \mathbf{R}_i) \frac{\nabla^2}{2m} w(\mathbf{r} - \mathbf{R}_j) d^3r \\ &= + \frac{1}{2m} \int (\nabla w(\mathbf{r} - \mathbf{R}_i))^* (\nabla w(\mathbf{r} - \mathbf{R}_j)) d^3r. \end{aligned}$$

We used a partial integration considering the boundary conditions of the Wannier orbitals $w(\pm\infty) = 0$. Therefore one part of the partial integration vanishes and we integrate/differentiate the integrands in the other integral, leading to two ∇ s. Further we see that $t_{ij} = t_{ji}^*$ by swapping the two integrands.

3.1.2 Superconductivity

Previous study of ours on the superconductivity have led us to the following Hamiltonian:

$$H_S = - \int U(\mathbf{r}) \psi_\downarrow^\dagger(\mathbf{r}) \psi_\uparrow^\dagger(\mathbf{r}) \psi_\uparrow(\mathbf{r}) \psi_\downarrow(\mathbf{r}) d^3r$$

on which we can apply a mean field approximation $\Delta(\mathbf{r}) = U(\mathbf{r}) \langle \psi_\uparrow(\mathbf{r}) \psi_\downarrow(\mathbf{r}) \rangle$. This yields to a common BCS-Hamiltonian for regular superconductors.

$$H_S = - \int \left(\Delta(\mathbf{r}) \psi_\downarrow^\dagger(\mathbf{r}) \psi_\uparrow^\dagger(\mathbf{r}) + \Delta(\mathbf{r})^* \psi_\uparrow(\mathbf{r}) \psi_\downarrow(\mathbf{r}) \right) d^3r.$$

we see that the second integrand is just the complex conjugate of the first one. To spare some place, we are going to focus ourselves on the first one and denote its homologue with *h.c.* ‘‘hermitian conjugate’’.

We insert 39 and obtain:

$$\begin{aligned} H_S &= - \sum_{ij} c_{i\downarrow}^\dagger c_{i\uparrow}^\dagger \int \Delta(\mathbf{r}) w^*(\mathbf{r} - \mathbf{R}_i) w(\mathbf{r} - \mathbf{R}_j) d^3r + \text{h.c.} \\ &:= - \sum_{ij} \Delta_{ij} c_{i\downarrow}^\dagger c_{i\uparrow}^\dagger + \text{h.c.} \end{aligned}$$

$\Delta(\mathbf{r})$ is an order parameter and doesn’t vary too much in the coherence length, which is much bigger than the atomic length. Therefore we can say that the orbitals vary faster than the gap. Moreover these orbitals are peaked in the neighbourhood of the atomic location \mathbf{R}_i and \mathbf{R}_j . Achieving the integral we get $\Delta_{ij} = \Delta_i \delta_{ij}$. We can from then reintroduce the *h.c.* and we get

$$H_S = - \sum_i \Delta_i c_{i\downarrow}^\dagger c_{i\uparrow}^\dagger + \Delta_i^* c_{i\uparrow} c_{i\downarrow}. \quad (41)$$

We however we’re missing the mean field term E_0 :

$$E_0 = \int U \langle \psi_\downarrow^\dagger \psi_\uparrow^\dagger \rangle \langle \psi_\uparrow \psi_\downarrow \rangle d^3r = \int U \frac{\Delta^*}{U} \frac{\Delta}{U} d^3r = \int \frac{|\Delta|^2}{U} d^3r.$$

and after applying the tight binding formalism we get:

$$E_0 = \sum_i \frac{|\Delta_i|^2}{U},$$

which is a term we can add to the Hamiltonian 41. From these equations we have the final Hamiltonian for the superconducting system:

$$H = E_0 + H_N + H_S.$$

3.2 A more symmertic Hamiltonian

As we introduced it while motivating the Bogoliubov-de Gennes formalism, we aspire to describe each state as a vector-matrix-vector product of

$$\hat{c}_i = \left(c_{i\uparrow}, c_{i\downarrow}, c_{i\uparrow}^\dagger, c_{i\downarrow}^\dagger \right).$$

However using the form we have in the superconducting 41 and normal 40 Hamiltonian will later not act as a fermionic operator upon the transformation we're about to do. We need to rewrite the Hamiltonian in a more symmertic way to later respect the anticommutation relations.

The chemical potential term can be expressed using the anticommutation relations of the fermionic operators $[c_{i\sigma}^\dagger, c_{i\sigma}]_+ = 1$:

$$\sum_{i\sigma} \mu_i c_{i\sigma}^\dagger c_{i\sigma} = \frac{1}{2} \sum_{i\sigma} \mu_i \left(c_{i\sigma}^\dagger c_{i\sigma} - c_{i\sigma} c_{i\sigma}^\dagger + 1 \right) \quad (42)$$

The trick we used is quite straight forward but not obvious:

$$c^\dagger c = \frac{1}{2} c^\dagger c + \frac{1}{2} c^\dagger c = \frac{1}{2} c^\dagger c + \underbrace{\frac{1}{2} c^\dagger c + \frac{1}{2} c c^\dagger}_{\frac{1}{2} [c^\dagger, c]_+ = \frac{1}{2}} - \frac{1}{2} c c^\dagger \quad (\mathfrak{I}r1)$$

In the same way the hopping term can be expressed as:

$$\sum_{\langle ij \rangle \sigma} t_{ij} c_{i\sigma}^\dagger c_{j\sigma} = \frac{1}{2} \sum_{\langle ij \rangle \sigma} t_{ij} \left(c_{i\sigma}^\dagger c_{j\sigma} - c_{j\sigma} c_{i\sigma}^\dagger \right).$$

we can take the liberty to reorder the indicies in a term of a sum and use the fact that $t_{ij} = t_{ji}^*$:

$$\sum_{\langle ij \rangle \sigma} t_{ij} c_{i\sigma}^\dagger c_{j\sigma} = \frac{1}{2} \sum_{\langle ij \rangle \sigma} t_{ij} c_{i\sigma}^\dagger c_{j\sigma} - t_{ji} c_{i\sigma} c_{j\sigma}^\dagger = \frac{1}{2} \sum_{\langle ij \rangle \sigma} t_{ij} c_{i\sigma}^\dagger c_{j\sigma} - t_{ji}^* c_{i\sigma} c_{j\sigma}^\dagger. \quad (43)$$

We then finish this section by using 42 and 43 in the Hamiltonian and obtain the following form:

$$H = E_0 - \frac{1}{2} \sum_{i\sigma} \left(c_{i\sigma}^\dagger c_{i\sigma} - c_{i\sigma} c_{i\sigma}^\dagger \right) - \frac{1}{2} \sum_{\langle ij \rangle \sigma} t_{ij} c_{i\sigma}^\dagger c_{j\sigma} - t_{ji}^* c_{i\sigma} c_{j\sigma}^\dagger.$$

The constant term $\frac{1}{2} \sum_{i\sigma} \mu_i$ of the normal Hamiltonian just vanished in the E_0 . right? We can now rewrite the Hamiltonian in a more compact way:

$$H = E_0 - \frac{1}{2} \sum_{i,j} \hat{c}_i^\dagger \hat{H}_{ij} \hat{c}_j \quad (44)$$

where the on site matrix reads

$$\hat{H}_{ij} = \begin{pmatrix} \mu_i \mathbb{I}_2 \delta_{ij} + t_{ij} & -i\sigma_2 \Delta_i \delta_{ij} \\ i\sigma_2 \Delta_i^* \delta_{ij} & -\mu_i \mathbb{I}_2 \delta_{ij} - t_{ij}^* \end{pmatrix} = \begin{pmatrix} H_{ij} & \Delta_{ij} \\ \Delta_{ij}^\dagger & -H_{ij}^* \end{pmatrix} \quad (45)$$

where we use \mathbb{I}_n as an n -dimensional identity matrix. We havn't explicitly removed the t_{ij} if we're not considering nearest neighbours. At this point it's intersting to note that if we wish to build some periodic boundary conditions, it might be the case that a site on side is neighbour with a site on the other side.

We can further compress our \hat{c}_i operator by introducing

$$\check{c} = (\hat{c}_1, \dots, \hat{c}_N)$$

along with the system Hamiltonian-matrix $\check{H}_{ij} := \hat{H}_{ij}$ wich allows us to rewrite the Hamiltonian 44 as:

$$H = E_0 - \frac{1}{2} \check{c}^\dagger \check{H} \check{c}. \quad (46)$$

3.3 Eigenvalues

We now have a look at the following eigenvalue problem, which later helps from the diagonalization of the Hamiltonian:

$$\tilde{H}\tilde{\chi}_n = E_n\tilde{\chi}_n$$

n runs over the number of the eigenvalue and $\tilde{\chi}_n$ is the corresponding eigenvector. We can decompose the $\tilde{\chi}_n$ to reflect each lattice site: $\tilde{\chi}_n = (\tilde{\chi}_{n1}, \dots, \tilde{\chi}_{nN})$. This means $\chi_{n,i}$ refers to a 4×4 block, i.e. the one on the submatrix we had earlier talked about. Therefore this $\chi_{n,i}$ contains four values, grouped in two vectors of length two, one for each spin: $\chi_{n,i} = (u_{ni}, v_{ni})$. Further $u_{ni} = (u_{ni\uparrow}, u_{ni\downarrow})$ couples to the two first components $(c_{i\uparrow}, c_{i\downarrow})$ we had in \hat{c} and similarly $v_{ni} = (v_{ni\uparrow}, v_{ni\downarrow})$ to the two last components $(c_{i\uparrow}^\dagger, c_{i\downarrow}^\dagger)$ of the four operator \hat{c} .

We can simplify the eigenvalue problem by taking a look only at a site i . We then only sum up over i -th row of \tilde{H}_{ij} with the components of $\tilde{\chi}_n$:

$$\sum_{j \in \llbracket N \rrbracket} \hat{H}_{ij} \tilde{\chi}_{nj} = E_n \tilde{\chi}_{ni}.$$

We remember that \tilde{H}_{ij} represent a complex scalar and \hat{H}_{ij} is a 4×4 matrix with complex entries. So it follows by reintroducing the following set of equations:

$$\begin{cases} \sum_{j \in \llbracket N \rrbracket} H_{ij} u_{nj} + \Delta v_{nj} = & E_n u_{nj} \\ \sum_{j \in \llbracket N \rrbracket} \Delta^\dagger u_{nj} - H_{ij}^* v_{nj} = & E_n v_{nj}. \end{cases} \xrightarrow{(1)} \begin{cases} \sum_j H_{ij} u_{nj} + \Delta v_{nj} = & E_n u_{nj} \\ \sum_j H_{ij} v_{nj}^* + \Delta^\dagger u_{nj}^* = & -E_n v_{nj}^*. \end{cases} \quad (47)$$

Where in (1) we took the conjugate of the second equation and used $\Delta^\dagger = -\Delta^*$. This is an important result, because it shows that if $\tilde{\chi}_n = (u_{n1}, v_{n1}, u_{n2}, v_{n2}, \dots)$ is an eigenvector with eigenvalue E_n , then so should be $(v_{n1}^*, u_{n1}^*, v_{n2}^*, u_{n2}^*, \dots)$ with the eigenvalue $-E_n$.

This leads to a symmetry in the energy spectrum of $H = E_0 \pm \frac{1}{2} \tilde{c}^\dagger \tilde{H} \tilde{c}$. This flexibility allows us to choose the version of H with the positive sign, which is more commonly used.

3.4 Diagonalization

Our goal is now to express the Hamiltonian relative to its energy eigenvalues, which is more practical to work with. As we have seen in the last section, eigenvectors χ_n allows us to compute the energies. Therefore we are going to diagonalize the Hamiltonian by using the eigenvectors χ_n to express the Hamiltonian according to its eigenvalues.

First we define a row-vector of our eigenstate $\tilde{X} = [\tilde{\chi}_{\pm 1}, \dots, \tilde{\chi}_{\pm 2N}]$ and introduce a diagonal matrix $\tilde{D} = \text{diag}(E_{\pm 1}, \dots, E_{\pm 2N})$ with the eigenvalues. Then we can write the Hamiltonian as:

$$\tilde{H} = \tilde{X} \tilde{D} \tilde{X}^{-1} = \tilde{X} \tilde{D} \tilde{X}^\dagger$$

we can then transform the Hamiltonian with $\tilde{c} := \tilde{X} \tilde{\gamma}$

$$\begin{aligned} H &= E_0 - \frac{1}{2} \tilde{c}^\dagger \tilde{H} \tilde{c} = E_0 - \frac{1}{2} \tilde{\gamma}^\dagger \tilde{X}^\dagger \tilde{H} \tilde{X} \tilde{\gamma} \\ &= E_0 - \frac{1}{2} \tilde{\gamma}^\dagger \underbrace{\tilde{X}^\dagger \tilde{X}}_{=\mathbb{I}} \tilde{D} \underbrace{\tilde{X}^{-1} \tilde{X}}_{=\mathbb{I}} \tilde{\gamma} \\ &= E_0 - \frac{1}{2} \tilde{\gamma}^\dagger \tilde{D} \tilde{\gamma} \\ &= E_0 - \frac{1}{2} \sum_{n \in \mathcal{N}} \end{aligned}$$

where $\mathcal{N} = \{\pm n : n \in \llbracket N \rrbracket\}$. Rearranging the transformation of \tilde{c} we get $\gamma = \tilde{X}^\dagger \tilde{c}$. Now that we've made the structure of the involved variables clear in the last section, we find the

expression of the γ wich is $2N$ -dimensional:

$$\begin{aligned}\gamma_n &= \sum_i \left(u_{ni\uparrow}^* c_{i\uparrow} + v_{ni\uparrow}^* c_{i\uparrow}^\dagger + u_{ni\downarrow}^* c_{i\downarrow} + v_{ni\downarrow}^* c_{i\downarrow}^\dagger \right) \\ &= \sum_{i\sigma} \left(u_{ni\sigma}^* c_{i\sigma} + v_{ni\sigma}^* c_{i\sigma}^\dagger \right)\end{aligned}$$

and due to the symmerty we saw erlier,

$$\gamma_{-n} = \sum_{i\sigma} \left(v_{ni\sigma} c_{i\sigma} + u_{ni\sigma} c_{i\sigma}^\dagger \right)$$

for $n \in \llbracket N \rrbracket$. We now take a look at the conjugate transpose of γ_{-n} . Because scalar are dimension 1×1 we have $(uc^\dagger)^\dagger = (c^\dagger)^\dagger u^\dagger = c^\dagger u^* = u^* c$ and it follows:

$$\gamma_{-n}^\dagger = \sum_{i\sigma} \left(v_{ni\sigma}^* c_{i\sigma}^\dagger + u_{ni\sigma}^* c_{i\sigma} \right) = \gamma_n.$$

Using this we can link each γ_i to the corresponding eigenvalue E_i : γ_n to the corresponding eigenvalue E_n and γ_{-n} to the corresponding eigenvalue $E_{-n} = -E_n$. We recall that we had $2N$ degrees of freedom $c_{i\sigma}$ due to the spins and after the transformation we get $4N$ degrees into \hat{c}_i . But because our energies E_n and E_{-n} are realted to eachother, we can keep the positive $2N$ eigenvalues and this maintain the total number of degree of freedom.

We can split the sum over the $n \in \mathcal{N}$ in two parts: $\mathcal{N}_+ = \{n \in \mathcal{N} : n > 0\}$, $\mathcal{N}_- = \{n \in \mathcal{N} : n < 0\}$

$$\begin{aligned}H &= E_0 + \frac{1}{2} \sum_{n \in \mathcal{N}_+} E_n \gamma_n^\dagger \gamma_n + \frac{1}{2} \sum_{n \in \mathcal{N}_-} E_n \gamma_n^\dagger \gamma_n \\ &= E_0 + \frac{1}{2} \sum_{n \in \mathcal{N}_+} E_n \gamma_n^\dagger \gamma_n + \frac{1}{2} \sum_{n \in \mathcal{N}_+} E_{-n} \gamma_{-n}^\dagger \gamma_{-n} \\ &= E_0 + \frac{1}{2} \sum_{n \in \mathcal{N}_+} E_n \gamma_n^\dagger \gamma_n - \frac{1}{2} \sum_{n \in \mathcal{N}_+} E_n \gamma_{-n}^\dagger \gamma_{-n} \\ &= E_0 + \frac{1}{2} \sum_{n \in \mathcal{N}_+} E_n \gamma_n^\dagger \gamma_n - \frac{1}{2} \sum_{n \in \mathcal{N}_+} E_n \gamma_n \gamma_n^\dagger \\ &= E_0 + \frac{1}{2} \sum_{n \in \mathcal{N}_+} E_n (\gamma_n^\dagger \gamma_n - \gamma_n \gamma_n^\dagger)\end{aligned}$$

where with used the energy symmetry and $\gamma_{-n}^\dagger = \gamma_n, \gamma_{-n} = \gamma_n^\dagger$.

Using this knowledge, we can express a final formula for the Hamiltonian by using the anti-commutation properties of the fermionic γ -operators: $[\gamma_n^\dagger, \gamma_n]_+ = 1$, so using the trick $\mathfrak{T}\mathfrak{r}1$ and bringing the $\frac{1}{2}$ prefactor in the sum:

$$H = E_0 - \sum_{n \in \llbracket N \rrbracket} E_n \left(\gamma_n^\dagger \gamma_n - \frac{1}{2} \right). \quad (48)$$

This is the final form of the Hamiltonian in the Bogoliubov-de Gennes formalism. As a user one should build the Hamiltonian and computes its eigenvalues,-vector and transform them into the γ operators.

3.4.1 Superconducting Gap

We already covered how the superconducting gap Δ is a relevant property of the Meissner state. We now aim to use the mean field theorie in order to find the gap. This requires a self consistency equation, which we can be derived from the Hamiltonian.

The gap was defined as $\Delta(\mathbf{r}) := U(\mathbf{r})\langle\psi_\uparrow(\mathbf{r})\psi_\downarrow(\mathbf{r})\rangle$. Back to the tight binding formalism, the gap now depends on the lattice site i and reads $\Delta_i = \langle c_{i\uparrow}c_{i\downarrow} \rangle$ and we can express $c_{i\sigma}$ in terms of the γ -operators:

$$\begin{aligned} c_{i\sigma} &= \sum_{n \in \mathcal{N}} u_{ni\sigma} \gamma_n \\ &= \sum_{n \in \mathcal{N}_+} u_{ni\sigma} \gamma_n + u_{-n,i\sigma} \gamma_{-n} \\ &= \sum_{n \in \mathcal{N}_+} u_{ni\sigma} \gamma_n + v_{ni\sigma}^* \gamma_n^\dagger \end{aligned}$$

where we used the symmetry of the eigenvectors. We can now compute expectation value involved in the gap:

$$\begin{aligned} \langle c_{i\uparrow}c_{i\downarrow} \rangle &= \sum_{n,m \in \mathcal{N}_+} \langle (u_{ni\uparrow} \gamma_n + v_{ni\uparrow}^* \gamma_n^\dagger) (u_{mi\downarrow} \gamma_m + v_{mi\downarrow}^* \gamma_m^\dagger) \rangle \\ &= \sum_{n,m \in \mathcal{N}_+} \langle (u_{ni\uparrow} u_{mi\downarrow} \gamma_n \gamma_m + u_{ni\uparrow} v_{mi\downarrow}^* \gamma_n \gamma_m^\dagger + v_{ni\uparrow}^* u_{mi\downarrow} \gamma_n^\dagger \gamma_m + v_{ni\uparrow}^* v_{mi\downarrow}^* \gamma_n^\dagger \gamma_m^\dagger) \rangle \\ &\stackrel{(*)}{=} \sum_{n \in \mathcal{N}_+} \langle u_{ni\uparrow} v_{ni\downarrow}^* \gamma_n \gamma_n^\dagger \rangle + \langle v_{ni\uparrow}^* u_{ni\downarrow} \gamma_n^\dagger \gamma_n \rangle \\ &= \sum_{n \in \mathcal{N}_+} u_{ni\uparrow} v_{ni\downarrow}^* \langle \gamma_n \gamma_n^\dagger \rangle + v_{ni\uparrow}^* u_{ni\downarrow} \langle \gamma_n^\dagger \gamma_n \rangle \\ &= \sum_{n \in \mathcal{N}_+} u_{ni\uparrow} v_{ni\downarrow}^* (1 - f(E_n)) + v_{ni\uparrow}^* u_{ni\downarrow} f(E_n) \end{aligned}$$


where f is the Fermi-Dirac distribution. In $(*)$ we notice no $\gamma\gamma$ or $\gamma^\dagger\gamma^\dagger$ terms in the Hamiltonian, so their expectation value is zero ¹.

The expectation value $\langle a\hat{A} \rangle_\Phi$ of a scalar times an operator reads $\langle \Phi | a\hat{A} | \Phi \rangle_\Phi = a \langle \Phi | \hat{A} | \Phi \rangle_\Phi = a \langle \hat{A} \rangle_\Phi$. To convince ourselves, we just take a look at the first quantisation expression of this bracket. This result leads to the self consistency equation:

$$\Delta_i = U_i \sum_{n \in \mathcal{N}_+} u_{ni\uparrow} v_{ni\downarrow}^* (1 - f(E_n)) + u_{ni\downarrow} v_{ni\uparrow}^* f(E_n)$$


We plan to solve this equation numerically, inserting some guess in the Hamiltonian, diagonalize it, update Δ_i and reinsert it into H and repeat until we reach a fixpoint.

4 Results




□ ————— □

¹This is like the expectation value of killing twice a fermion in a state. It is not possible, because we can't annihilate a state that has a possession number of zero. And in the same way due to the Pauli-principle we can't have more than one particle in the same state, so $\langle \gamma^\dagger \gamma^\dagger \rangle = 0$. Here we forget the indices.. [why?](#) The Hamiltonian is diagonal in $\gamma\gamma^\dagger$ and $\gamma^\dagger\gamma$ [right?](#)




(a) Surrounded with vacuum.




(b) Vacuum on the right and left with vertical periodic boundary conditions.


(c) Both vertical and horizontal periodic boundary conditions.



(a) Surrounded with vacuum.



(b) Vacuum on the right and left with vertical periodic boundary conditions.



(c) Both vertical and horizontal periodic boundary conditions.

5 Literature

Books

- [1] Kristian Fossheim and Asle Sudbø. John Wiley & Sons, Ltd, 2004. ISBN: 9780470020784. DOI: <https://doi.org/10.1002/0470020784>. eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/0470020784>. URL: <https://onlinelibrary.wiley.com/doi/abs/10.1002/0470020784>.

Online

- [2] Prof. Dr. Matthias Fuchs and Philipp Baumgärtel. Last public available version 2010. 2023. URL: http://theorie.physik.uni-konstanz.de/lmfuchs/lectures/statmech0910/Stat.Mech0910_Skript.pdf.