

# HOMework 2B: What is a Green's Function

COURSE: Physics 231, *Methods of Theoretical Physics* (2018)

INSTRUCTOR: Professor Flip Tanedo ([flip.tanedo@ucr.edu](mailto:flip.tanedo@ucr.edu))

DUE BY: Mon, October 29

## 1 A Green's function by completeness

Three formulaic ways of solving for Green's functions are:

1. Fourier transforming to turn the differential operator into an algebraic one, and then doing a contour integral to go back to position space.
2. Solving the homogeneous equation and patching together two solutions over the  $\delta$ -function.
3. Using the completeness of eigenfunctions.

We will spend most of this course highlighting method 1. Method 2 will be familiar from electrodynamics. Here we use the third method, which is based on the eigenfunction completeness relation,

$$\sum_n e_n^*(y) e_n(x) = \delta(x - y) . \quad (1.1)$$

If you know the eigenvalues of the linear differential operator,  $L_x$ , then you can use this relation to hack together the Green's function  $G$  because  $L_x G(x, y) = \delta(x - y)$ . The subscript  $x$  in  $L_x$  means that the operator is a function of  $x$  and derivative operators with respect to  $x$ .

REFERENCES: This method is discussed in Matthews & Walker chapter 9-4, Stone & Goldbart chapter 5.4, Cahill chapter 6.38, and Byron & Fuller chapter 7.2.

In this problem, we consider a second order differential equation acting on a state  $\psi(x)$  with a source  $s(x)$ .

$$\left( -\frac{d^2}{dx^2} + k^2 \right) \psi(x) = s(x) .$$

We consider the domain  $x \in [0, 1]$  with boundary conditions  $\psi(0) = \psi(1) = 0$ .

### 1.1 Eigenfunctions

What are the eigenfunctions  $e_n(x)$  and the corresponding eigenvalues  $\lambda_n$ ? Recall that an eigenfunction of a linear operator  $L_x$  satisfies  $L_x e_n(x) = \lambda_n e_n(x)$ .

HINT: The eigenfunctions of  $-d^2/dx^2$  are  $\psi_n(x) = \sqrt{2} \sin(n\pi x)$  with eigenvalues  $n^2\pi^2$ . What changes when the operator  $L_x$  includes an additive constant?

## 1.2 Completeness

Write down the Green's function  $G(x, y)$  using the completeness trick with respect to eigenfunctions  $e_n$  with eigenvalues  $\lambda_n$ :


$$G(x, y) = \sum_n \frac{e_n^*(y) e_n(x)}{\lambda_n} . \quad (1.2)$$

HINT: Once you did the previous part, this step takes about twenty seconds depending on how quickly you write.

## 1.3 Extra Credit: Hands on (recommended)

You have a free license for several types of scientific software through UCR through MySoftware<sup>1</sup>. This problem requires some plotting. I suggest using *Mathematica*<sup>2</sup>.

Plot the solution to the Green's function  $G(x, y)$  for  $y = 0.5$  and  $k = 0.2$ , summing from  $n = 1$  to  $n = 10$ . Here's the general way to format it in *Mathematica*:



The highlighted piece and the example plot are *completely wrong*! Fill it in with the correct  $G(x, y)$  and plot it. It may help to use the ‘Basic Math Input’ palette if you’re unfamiliar with this. If this task is very painful, please ask a friend. If this is still very painful, then you may use any other plotting program that you wish.

Does this shape make sense? What do you expect will happen as  $n$  becomes larger? Try it for  $n = 100$ . Explore what happens as you change  $k$  and  $y$ . (Make sure  $y$  is in the domain of the function space!) Take a moment to have a nice warm cup of tea, enjoy the agreeable weather outside, and meditate upon why this shape of the Green's function makes sense and the kinds of physical problem where Green's function may be relevant. You do not have to write up these meditations, but I do strongly suggest the tea and outdoors.

---

<sup>1</sup><http://cnc.ucr.edu/mysoftware/>

<sup>2</sup>It sucks that this is not open source software. The closest open source alternative is SciPy with a Jupyter notebook.

## 2 A Green's function by patching

*This looks like a long problem, but it's mostly reading and understanding. Each step is modest.*

In this problem we solve for a Green's function by patching together solutions of the homogeneous equation. We examine the same system that you did in Homework 4, Problem #2: consider a second order differential equation acting on a state  $\psi(x)$  with a source  $s(x)$ .

$$\mathcal{O}\psi(x) = \left(-\frac{d^2}{dx^2} + k^2\right)\psi(x) = s(x) .$$

We consider the domain  $x \in [0, 1]$  with boundary conditions  $\psi(0) = \psi(1) = 0$ . **For simplicity**, we'll take  $k^2 = 0$  and leave the  $k^2 \neq 0$  case for extra credit.

We want to solve the Green's function equation,

$$\mathcal{O}G(x, y) = \delta(x - y) .$$

In this method, we solve the *homogeneous* equation  $\mathcal{O}G(x, y) = 0$  for  $x \neq y$  in the two regions  $x < y$  and  $x > y$ . These two solutions will each have independent coefficients that we must patch together at  $x = y$ .

### 2.1 Solve the homogeneous equation

Solve the two *homogeneous* equations in the regions away from the  $\delta$ -function spike:

$$-\left(\frac{d}{dx}\right)^2 G_{<}(x, y) = 0 \quad \text{in} \quad 0 < x < y \leq 1 \quad (2.1)$$

$$-\left(\frac{d}{dx}\right)^2 G_{>}(x, y) = 0 \quad \text{in} \quad 0 < y < x \leq 1 \quad (2.2)$$

Don't forget to impose the boundary conditions  $G(0, y) = G(1, y) = 0$ . Understand why it makes sense that  $\psi(0) = \psi(1) = 0$  imposes the same condition on the Green's functions. If the Green's function did not satisfy these conditions, then you could construct sources that violate the boundary conditions because  $\psi(1) = \int G(1, y)s(y) dy \neq 0$ .

The Green's function  $G(x, y)$  is now piecewise defined

$$G(x, y) = \begin{cases} G_{<}(x, y) & \text{if } x < y \\ G_{>}(x, y) & \text{if } x > y \end{cases} . \quad (2.3)$$

Check to make sure that you have the correct number of undetermined coefficients.

HINT: if you're not sure what the solution to the homogeneous differential equation is, then you're over thinking it. It's not a trigonometric function. If you're still confused, refer back to your plot the previous problem.

## 2.2 Patching, Part 1

For fixed  $y$ , integrate the Green's function equation over an infinitesimal sliver,  $x \in (y - \epsilon, y + \epsilon)$ , with  $\epsilon \rightarrow 0$ :

$$\int_{y-\epsilon}^{y+\epsilon} dx \left( \frac{d}{dx} \right)^2 G(x, y) = \int_{y-\epsilon}^{y+\epsilon} dx \delta(x - y) . \quad (2.4)$$

HINT: It may be useful to remember that

$$\int_{y-\epsilon}^{y+\epsilon} dx \frac{df(x)}{dx} = f(y + \epsilon) - f(y - \epsilon) . \quad (2.5)$$

Write out the resulting equation in terms of  $dG_{<}/dx$  and  $dG_{>}/dx$  at  $x = y$ . You will find that the slope of the Green's function,  $dG/dx$ , is *discontinuous* at  $x = y$ .

## 2.3 Patching, Part 2

Integrate (2.4) again over  $x \in (y - \epsilon, y + \epsilon)$  to relate  $G_{<}$  and  $G_{>}$  at  $x = y$ .

ANSWER:  $dG/dx$  has a finite discontinuity at  $x = y$ , therefore  $G$  is *continuous* at  $x = y$ . The discontinuity in the slope gives a kink in  $G$ ,

$$G_{<}(x, y)|_{x=y} = G_{>}(x, y)|_{x=y} . \quad (2.6)$$

We've found that the second derivative is a  $\delta$ -function (singular), the first derivative is simply discontinuous, and the zeroth derivative (the Green's function itself) is continuous.

## 2.4 Patching, Part 3

Take all of the above results and write down the piece-wise definition of  $G(x, y)$  in (2.3) as an explicit function of  $x$  and  $y$  with all coefficients determined. Sketch  $G(x, y)$  for  $y = 0.5$  and compare it to your plot from Problem #2.

## 2.5 Extra Credit: $k^2 \neq 0$

*This problem is not graded and is purely for your "enjoyment."* Perform the same steps for the case of  $k^2 \neq 0$ . The homogeneous solution is now a sine, which makes things a bit more complicated. The procedure is completely the same and the  $k^2 \lesssim 1$  case is well approximated by  $k^2 = 0$ , as you will have noticed from the plot in the previous problem. SOLUTION: Matthews & Walker, Chapter 9–4.

### 3 Practice with complex integration

#### 3.1 This is not a closed loop

The goal of this problem is to review line integrals in a 2D space. Perform the integral

$$\int_i^{1+i} dz \quad (3.1)$$

with respect to the path that connects  $z = i$  and  $z = i + 1$  by a straight line segment. This is *not* a closed loop, there's no magic formula for this. Just parameterize the path by writing  $z$  as a function of a real parameter. Then write out  $dz$  in terms of the parameter. Do the integral.

#### 3.2 Contour integrals

Calculate the following integrals using the residue theorem:

(a)  $\int_0^\infty \frac{x^2}{x^4 + 16} dx$

(b)  $\int_{-\infty}^\infty \frac{\sin x}{x^2 + 4x + 5} dx$

(c)  $\int_0^\infty \frac{1}{x^6 + 1} dx$  using a contour including the real line and a large semicircle in the complex plane

(d)  $\int_0^\infty \frac{1}{x^6 + 1} dx$  using a contour that encloses the 'pizza slice' wedge between  $\theta = 0$  and  $\theta = \pi/3$ .

HINT: for (d), the line integral along the diagonal is proportional to the integral you want.

### 4 Preparing for a personal statement

The National Science Foundation Graduate Research Fellowship is a prestigious graduate fellowship that supports students for three years with a stipend. This frees you from teaching obligations, provides research flexibility, and is an excellent item on a CV. While not everyone is eligible for the fellowship, the components are fairly typical of the many kinds of funding opportunities you will see in your research career (including grants, telescope time requests, etc.). Find information about the GRFP here: <https://www.nsfgrfp.org>

- (a) What are the eligibility requirements for an NSF graduate research fellowship with respect to citizenship?
- (b) What are the main components of the application? How many pieces of writing, what are the goals of each essay?
- (c) How many letters of recommendation are required? If you were to apply this year, who would those letters be from?
- (d) What are the main criteria for evaluating the application?

- (e) What resources are available at UCR to help you with writing?
- (f) When is the application due, when is the last time one is eligible to apply?
- (g) Write a list of five ideas that you would include in your research statement if you were to apply for this fellowship.
- (h) Write a list of five ideas that you would include in your personal statement if you were to apply for this fellowship.

## Extra Credit

These problems are not graded and are for your edification. You are strongly encouraged to explore and discuss these topics, especially if they are in a field of interest to you.

### 1 Black Hole Entropy and Dimensional Analysis

This problem is from Tony Zee's *Einstein Gravity in a Nutshell*. This is recommended for those interested in cosmology and condensed matter physics.

#### 1.1 The Planck Mass

In natural units, the Newton constant  $G$  has [mass] dimension  $[G] = -2$  (here we're using natural units—see HW1b, extra credit problem 2), so that we can define a mass scale

$$M_P = \frac{1}{\sqrt{G}} . \quad (1.1)$$

This is called the **Planck mass**. Restore the factors of  $\hbar$  and  $c$  to make this definition correct in 'unnatural' units where we keep track of length and time dimensions.

#### 1.2 Hawking Radiation

Black holes can evaporate by Hawking radiation. A cartoon picture of this process is as follows: quantum mechanics + special relativity tells us that the vacuum ('empty' space) is composed of virtual particle-anti-particle pairs. Near the event horizon of a black hole, one of these particles can fall into the black hole while the other radiates away as a physical particle. This means that black holes have a temperature.

Use dimensional analysis to determine how this **Hawking temperature** scales with the mass of the black hole. How does  $T_H$  scale with the combination  $GM$ ? Observe that the black hole gets *hotter* as it loses energy.

HINT: there's one subtlety. There are two mass scales in the problem: the black hole mass,  $M$ , and the Planck mass,  $M_P = 1/\sqrt{G}$ . (In natural units, of course). In order to be able to use dimensional analysis, the additional piece of information is that  $G$  is a gravitational coupling, so that gravitational effects should go like  $GM$ .

### 1.3 The Holographic Principle

Recall from thermodynamics that entropy,  $S$ , is related to energy  $E$  and temperature  $T$  by

$$\frac{dS}{dE} = \frac{1}{T} . \quad (1.2)$$

- (i) Identify the temperature with the Hawking temperature  $T = T_H$  and set the energy to be the mass of the black hole so that  $dE = dM$ . Integrate with respect to the black hole mass to find how entropy scales with mass,  $S \sim M^?$ .
- (ii) The radius of the black hole's event horizon scales like  $R = GM$ . How does the black hole's entropy scale with its characteristic length scale?
- (iii) Contrast the above result to the expected scaling of entropy in ordinary thermodynamics. Recall that entropy is an *extensive*<sup>3</sup> measure of the number of microstates in a system.

The solution to this problem is spelled out in the introduction to Zee's *Einstein's Gravity in a Nutshell*. The 'holographic principle' is the proposal that the properties of the black hole are encoded on its surface rather than its volume. This is analogous to how a hologram is a 3D image encoded onto a 2D surface. A manifestation of the holographic principle is the AdS/CFT correspondence, which posits that certain theories of strongly interacting systems in  $d$ -dimensions are mathematically identical to a weakly-coupled  $(d + 1)$ -dimensional gravitational theory.

## 2 Renormalization Group as Dimensional Analysis

This problem is recommended for particle/nuclear physicists and especially to theorists of any type.

Read the paper "Dimensional Analysis in Field Theory: An Elementary Introduction to Broken Scale Invariance and the Renormalization Group Equations" by Paul Stevenson<sup>4</sup>. This paper describes the phenomenon of dimensional transmutation in quantum/statistical field theory in a way that strips it of the mysticism that tends to appear when you first learn field theory. In particular, it explains why dimensionless 'coupling constants' are not really constant and are scale dependent. Understand the dimensional analysis theorem in the paper and then understand how that theorem is evaded in actual physics. Theorists should take time to understand this paper carefully.

## 3 Sturm–Liouville Operator

This question is based on Matthews & Walker, chapter 9–2 and/or Stone & Goldbart chapter 4.2. Consider the **Sturm–Liouville** differential operator

$$L = p_2(x) \left( \frac{d}{dx} \right)^2 + p_1(x) \frac{d}{dx} + p_0(x). \quad (3.1)$$

---

<sup>3</sup>An **extensive** property is one that adds for separate subsystem. Consider a system of two lazy cats. The mass of the combined system is equal to the sum of the masses of each cat. This is in contrast to temperature, which is **intensive**; the temperature of the multi-lazy-cat system is  $T_{\text{cat}}$  no matter how many cats there are. This example obviously breaks down at large numbers of cats.

<sup>4</sup>Annals Phys. **132** (1981) 383, [http://dx.doi.org/10.1016/0003-4916\(81\)90072-5](http://dx.doi.org/10.1016/0003-4916(81)90072-5)

Assume weight  $w(x) = 1$  and some domain  $x \in [a, b]$ .

### 3.1 Formal adjoint

Using integration by parts, show that the **formal adjoint** of this operator is

$$L^\dagger = \left( \frac{d}{dx} \right)^2 p_2(x) - \left( \frac{d}{dx} \right) p_1(x) + p_0(x) , \quad (3.2)$$

where we use the notation

$$\left[ \left( \frac{d}{dx} \right)^n p_n(x) \right] f(x) \equiv \left( \frac{d}{dx} \right)^n [p_n(x) f(x)] . \quad (3.3)$$

### 3.2 Self-Adjoint Case

Show that in order for  $L$  to be formally self-adjoint, that is  $L = L^\dagger$  (without considering boundary conditions), then the following conditions must hold:

$$p_1(x) = 2p_2'(x) - p_1(x) \quad (3.4)$$

$$p_0(x) = p_x''(x) - p_1(x) + p_0(x) . \quad (3.5)$$

Thus  $L$  is self-adjoint (Hermitian) if  $p_1(x) = p_2'(x)$ .  $L$  can be written succinctly as:

$$L = \frac{d}{dx} \left( p_2(x) \frac{d}{dx} \right) + p_0(x) . \quad (3.6)$$