

Statistical Signal Processing

Lecture 4: Inverse Problem

From Power Spectrum/Autocorrelation to Simulation Model

Carlas Smith & Peyman Mohajerin Esfahani

1

Overview

INVERSE Problems - Part I: Frequency Domain

- From Power Spectra (Frequency domain) to generating a stochastic process
- Chapter 7.1-7.3

INVERSE Problems - Part II: Time Domain

- From Autocorrelation (Time domain) to generating a stochastic process
- Chapter 7.4

Example: Kolmogorov Turbulence Model

“A Mathematician doing physics”




“... *combine*

*theoretical studies with the analysis of
experimental results ...”*

Example: Kolmogorov Turbulence Model

“A Mathematician doing Work on Turbulence modelling based on Experimental results (Power Spectra):



“... combine

theoretical studies with the analysis of experimental results ...”

$$P_{\phi}(f) = 0.023 r_0^{-\frac{5}{3}} f^{-\frac{11}{3}}$$

with r_0 - the Fried parameter
and f - the spatial frequency

Summary Lecture 3

Example Lecture 3: We work with Rational Transfer functions, such as:

$$H(z) = \underbrace{\frac{z}{1 - 0.9z}}_{\text{anti-causal } ([H(z)]_-)} + \underbrace{\frac{z^{-1}}{1 - 0.9z^{-1}}}_{\text{causal } ([H(z)]_+)} \quad \text{ROC}(H(z)) \supset \Gamma$$

with the pole(s) $z = \frac{1}{0.9}$ of $[H(z)]_-$ outside the unit circle and the pole(s) $z = 0.9$ of $[H(z)]_+$ inside the unit circle.


Summary Lecture 3

Example Lecture 3: We work with Rational Transfer functions, such as:

$$H(z) = \underbrace{\frac{z}{1 - 0.9z}}_{\text{anti-causal } ([H(z)]_-)} + \underbrace{\frac{z^{-1}}{1 - 0.9z^{-1}}}_{\text{causal } ([H(z)]_+)} \quad \text{ROC}(H(z)) \supset \Gamma$$

with the pole(s) $z = \frac{1}{0.9}$ of $[H(z)]_-$ outside the unit circle and the pole(s) $z = 0.9$ of $[H(z)]_+$ inside the unit circle.

ARMA(p,q) Lecture 3:



Then the Power spectrum $P_x(z)$ equals:

$$P_x(z) = \sigma_v^2 \frac{B_q(z)B_q^*(1/z^*)}{A_p(z)A_p^*(1/z^*)} \quad \text{Property : } P_x(z) = P_x^*(1/z^*)$$

For example $P_x(z)$ given as $\frac{1 - 2.5z^{-1} + z^{-2}}{1 - 2.05z^{-1} + z^{-2}}$.

A simple formulation of our first inverse problem

Given a “Power” spectrum as a rational function:

$$P_x(z) = \frac{1 - 2.5z^{-1} + z^{-2}}{1 - 2.05z^{-1} + z^{-2}}$$

Can you determine an ARMA model (including σ_v) with transfer function:

$$H(z) = \frac{B_q(z)}{A_p(z)}$$

such that,

$$P_x(z) = \sigma_v^2 \frac{B_q(z)B_q^*(1/z^*)}{A_p(z)A_p^*(1/z^*)}$$

Highly non-unique?

Today's Challenge: A simplified Inverse Problem

SPECTRAL FACTORIZATION PROBLEM: Assume that (1) $P_x(z)$ is *rational*
and (2) $P_x(z)$ is *positive real*, i.e.:

$$P_x(z) = P_x^*(1/z^*) \quad P_x(e^{j\omega}) \in \mathbb{R} \text{ and } P_x(e^{j\omega}) > 0$$

Today's Challenge: A simplified Inverse Problem

SPECTRAL FACTORIZATION PROBLEM: Assume that (1) $P_x(z)$ is *rational*
and (2) $P_x(z)$ is *positive real*, i.e.:

$$P_x(z) = P_x^*(1/z^*) \quad P_x(e^{j\omega}) \in \mathbb{R} \text{ and } P_x(e^{j\omega}) > 0$$

and (3) $\text{ROC}(P_x(z)) \supset \Gamma$, then find the rational **spectral factor** $Q(z)$ and the constant $\sigma \in \mathbb{R}_+$ such that:

Today's Challenge: A simplified Inverse Problem

SPECTRAL FACTORIZATION PROBLEM: Assume that (1) $P_x(z)$ is *rational* and (2) $P_x(z)$ is *positive real*, i.e.:

$$P_x(z) = P_x^*(1/z^*) \quad P_x(e^{j\omega}) \in \mathbb{R} \text{ and } P_x(e^{j\omega}) > 0$$

and (3) $\text{ROC}(P_x(z)) \supset \Gamma$, then find the rational **spectral factor** $Q(z)$ and the constant $\sigma \in \mathbb{R}_+$ such that:

1. $P_x(z) = \sigma Q(z)Q^*(1/z^*)$
2. $Q(z)$ is causal and stable
3. $Q(z)^{-1}$ is also causal and stable

Remark 1: The property that the inverse of $Q(z)$ is causal and stable is indicated by $Q(z)$ being *minimum phase*.

Remark 2: $Q(z)$ is **unique** if $Q(\infty) = 1$.

Solving the Spectral Factorization problem

1. **Use of the Spectral Factor**
2. Illustrations of Key Definitions: A rational transfer function, a minimum phase transfer function, $Q(\infty) = 1$.
3. Special form of the power spectrum $P_x(z)$.
 - Constraints on its coefficients
 - Constraints on its poles and zeros
4. A constructive Solution
5. Illustration

Use of the Spectral Factor

Given the unique spectral factorization of the Power spectrum

$$P_x(z) = P_x^*(1/z^*), \text{ as } P_x(z) = \sigma Q(z)Q^*(1/z^*)$$

1. For a generated

ZMWN $\hat{v}(n)$ with variance $\sigma \in \mathbb{R}_+$, we can determine $\hat{x}(n)$ as,



such that, $P_{\hat{x}}(z) = P_x(z)$

Use of the Spectral Factor

Given the unique spectral factorization of the Power spectrum

$$P_x(z) = P_x^*(1/z^*), \text{ as } P_x(z) = \sigma Q(z)Q^*(1/z^*)$$

1. For a generated

ZMWN $\hat{v}(n)$ with variance $\sigma \in \mathbb{R}_+$, we can determine $\hat{x}(n)$ as,



such that, $P_{\hat{x}}(z) = P_x(z)$

2. If we have a realization of $x(n)$ we can use this to find the corresponding ZMWN that generated this realization,

Use of the Spectral Factor

Given the unique spectral factorization of the Power spectrum

$$P_x(z) = P_x^*(1/z^*), \text{ as } P_x(z) = \sigma Q(z)Q^*(1/z^*)$$


1. For a generated

ZMWN $\hat{v}(n)$ with variance $\sigma \in \mathbb{R}_+$, we can determine $\hat{x}(n)$ as,



such that, $P_{\hat{x}}(z) = P_x(z)$

2. If we have a realization of $x(n)$ we can use this to find the corresponding ZMWN that generated this realization,



This filter is called the “whitening” filter.

Solving the Spectral Factorization problem

1. Use of the Spectral Factor
2. **Illustrations of Key Definitions:** A rational transfer function, a minimum phase transfer function, $Q(\infty) = 1$.
3. Special form of the power spectrum $P_x(z)$.
 - Constraints on its coefficients
 - Constraints on its poles and zeros
4. A constructive Solution
5. Illustration

Illustrations of Key Definitions: minimum phase TF

When $Q(z)$ is a (causal, stable and) minimum-phase rational transfer function given as:

then
$$Q(z) = \frac{B(z)}{A(z)}$$

1. $Q(z)$ is causal and stable,

Illustrations of Key Definitions: minimum phase TF

When $Q(z)$ is a (causal, stable and) minimum-phase rational transfer function given as:

then
$$Q(z) = \frac{B(z)}{A(z)}$$

1. $Q(z)$ is causal and stable, i.e. the zeros of $A(z)$ are inside (NOT on!) the unit circle, and,
2. $Q(z)^{-1}$ is causal and stable,

Illustrations of Key Definitions: minimum phase TF

When $Q(z)$ is a (causal, stable and) minimum-phase rational transfer function given as:

then
$$Q(z) = \frac{B(z)}{A(z)}$$

1. $Q(z)$ is causal and stable, i.e. the zeros of $A(z)$ are inside (NOT on!) the unit circle, and,
2. $Q(z)^{-1}$ is causal and stable, i.e. the zeros of $B(z)$ are inside (NOT on!) the unit circle.

Example 2: A causal, rational $Q(z)$ that is stable and minimum-phase is:

$$Q(z) = \frac{1 - 0.5z^{-1}}{1 - 0.8z^{-1}} \quad \text{zero : } z = 0.5 \quad \text{pole : } z = 0.8$$

Remark: Such $Q(z)$ in Example 2 have $Q(\infty) = 1$.

Illustrations of Key Definitions: Rational TF

When $P_x(z)$ is a rational function, it can be written as:

$$P_x(z) = \frac{N(z)}{D(z)}$$

for $N(z)$ and $D(z)$ finite order polynomials in z .

Example 1: A rational $P_x(z)$ is given as:

$$P_x(z) = \frac{1 - 2.5z^{-1} + z^{-2}}{1 - 2.05z^{-1} + z^{-2}}$$

Remarks:

Illustrations of Key Definitions: Rational TF

When $P_x(z)$ is a rational function, it can be written as:

$$P_x(z) = \frac{N(z)}{D(z)}$$

for $N(z)$ and $D(z)$ finite order polynomials in z .

Example 1: A rational $P_x(z)$ is given as:

$$P_x(z) = \frac{1 - 2.5z^{-1} + z^{-2}}{1 - 2.05z^{-1} + z^{-2}}$$

Remarks: (1) When $P_x(z)$ is a rational function in z it has *poles* and *zeros* and

Illustrations of Key Definitions: Rational TF

When $P_x(z)$ is a rational function, it can be written as:

$$P_x(z) = \frac{N(z)}{D(z)}$$

for $N(z)$ and $D(z)$ finite order polynomials in z .

Example 1: A rational $P_x(z)$ is given as:

$$P_x(z) = \frac{1 - 2.5z^{-1} + z^{-2}}{1 - 2.05z^{-1} + z^{-2}}$$

Remarks: (1) When $P_x(z)$ is a rational function in z it has *poles* and *zeros* and (2) $P_x(e^{j\omega})$ does not contain discontinuities (due to harmonics): $x(n)$ is a regular RP.

Solving the Spectral Factorization problem

1. Use of the Spectral Factor
2. Illustrations of Key Definitions: A rational transfer function, a minimum phase transfer function, $Q(\infty) = 1$.
3. **Special form of the power spectrum $P_x(z)$.**
 - Constraints on its coefficients
 - Constraints on its poles and zeros
4. A constructive Solution
5. Illustration

Special form of the power spectrum $P_x(z)$

The special form of the Power Spectrum $P_x(z) = P_x^*(1/z^*)$ constraints its pole-zero location.

Example 3-b: Let $P_x(z) = az + b + cz^{-1}$, then $P_x(z) = P_x^*(1/z^*)$ yields $a^* = c$, $b^* = b$ and $P_x(z)$ is given as,

$$P_x(z) = az + b + a^*z^{-1} \quad b \in \mathbb{R}, a \in \mathbb{C} \quad \text{rootsPolReal.m}$$

Special form of the power spectrum $P_x(z)$ (II)

LEMMA: If $P_x(z) = \frac{N(z)}{D(z)}$, $N(z), D(z)$ coprime and $P_x(z) = P_x^*\left(\frac{1}{z^*}\right)$, then,

If p_0 is a pole of $P_x(z) \Rightarrow \frac{1}{p_0^*}$ is a pole of $P_x(z)$

If z_0 is a zero of $P_x(z) \Rightarrow \frac{1}{z_0^*}$ is a zero of $P_x(z)$

Special form of the power spectrum $P_x(z)$ (II)

LEMMA: If $P_x(z) = \frac{N(z)}{D(z)}$, $N(z), D(z)$ coprime and $P_x(z) = P_x^*\left(\frac{1}{z^*}\right)$, then,

If p_0 is a pole of $P_x(z) \Rightarrow \frac{1}{p_0^*}$ is a pole of $P_x(z)$

If z_0 is a zero of $P_x(z) \Rightarrow \frac{1}{z_0^*}$ is a zero of $P_x(z)$

Proof: Since $P_x^*\left(\frac{1}{z^*}\right) = \frac{N^*\left(\frac{1}{z^*}\right)}{D^*\left(\frac{1}{z^*}\right)} \Rightarrow \frac{N(z)}{D(z)} \frac{D^*\left(\frac{1}{z^*}\right)}{N^*\left(\frac{1}{z^*}\right)} = 1$

Therefore *perfect* pole-zero cancellation is necessary. Since $N(z)$ and $D(z)$ are coprime, and the order of $N(z)$ is identical to that of $N^*\left(\frac{1}{z^*}\right)$, \Rightarrow

Special form of the power spectrum $P_x(z)$ (II)

LEMMA: If $P_x(z) = \frac{N(z)}{D(z)}$, $N(z)$, $D(z)$ coprime and $P_x(z) = P_x^*\left(\frac{1}{z^*}\right)$, then,

If p_0 is a pole of $P_x(z) \Rightarrow \frac{1}{p_0^*}$ is a pole of $P_x(z)$

If z_0 is a zero of $P_x(z) \Rightarrow \frac{1}{z_0^*}$ is a zero of $P_x(z)$

Proof: Since $P_x^*\left(\frac{1}{z^*}\right) = \frac{N^*\left(\frac{1}{z^*}\right)}{D^*\left(\frac{1}{z^*}\right)} \Rightarrow \frac{N(z)}{D(z)} \frac{D^*\left(\frac{1}{z^*}\right)}{N^*\left(\frac{1}{z^*}\right)} = 1$

Therefore *perfect* pole-zero cancellation is necessary. Since $N(z)$ and $D(z)$ are coprime, and the order of $N(z)$ is identical to that of $N^*\left(\frac{1}{z^*}\right)$, \Rightarrow a zero of $N(z)$ has to be a zero of $N^*\left(\frac{1}{z^*}\right)$.

Hence,

Special form of the power spectrum $P_x(z)$ (II)

LEMMA: If $P_x(z) = \frac{N(z)}{D(z)}$, $N(z)$, $D(z)$ coprime and $P_x(z) = P_x^*(\frac{1}{z^*})$, then,

If p_0 is a pole of $P_x(z) \Rightarrow \frac{1}{p_0^*}$ is a pole of $P_x(z)$

If z_0 is a zero of $P_x(z) \Rightarrow \frac{1}{z_0^*}$ is a zero of $P_x(z)$

Proof: Since $P_x^*(\frac{1}{z^*}) = \frac{N^*(\frac{1}{z^*})}{D^*(\frac{1}{z^*})} \Rightarrow \frac{N(z)}{D(z)} \frac{D^*(\frac{1}{z^*})}{N^*(\frac{1}{z^*})} = 1$

Therefore *perfect* pole-zero cancellation is necessary. Since $N(z)$ and $D(z)$ are coprime, and the order of $N(z)$ is identical to that of $N^*(\frac{1}{z^*})$, \Rightarrow a zero of $N(z)$ has to be a zero of $N^*(\frac{1}{z^*})$.

Hence,

$$\text{If } \exists z_0 \in \mathbb{C} : N(z_0) = 0 \Rightarrow N^*(\frac{1}{z_0^*}) = 0 \Rightarrow N(\frac{1}{z_0^*}) = 0$$

Summary of the Properties of $P_x(z)$

From the properties:


1. $P_x(z) = P_x^*(1/z^*) \Rightarrow$ contains poles p_0 and $1/p_0^*$ and zeros z_0 and $1/z_0^*$.
2. $P_x(e^{j\omega}) > 0$ means that $P_x(z)$ does not contain zeros on the unit circle.
3. Since $\text{ROC}(P_x(z)) \supset \Gamma$, $P_x(z)$ does not contain poles on the unit circle.

Solving the Spectral Factorization problem

1. Use of the Spectral Factor
2. Illustrations of Key Definitions: A rational transfer function, a minimum phase transfer function, $Q(\infty) = 1$.
3. Special form of the power spectrum $P_x(z)$.
 - Constraints on its coefficients
 - Constraints on its poles and zeros
4. **A constructive Solution**
5. Illustration

A constructive Solution to the Spectral Factorization Problem


Pole-zero pattern $P_x(z)$



A constructive Solution to the Spectral Factorization Problem


Pole-zero pattern $P_x(z)$

Stable pole-zero part of $P_x(z)$:



$$Q(z) = \frac{(1 - z_1 z^{-1})(1 - z_2 z^{-1})}{(1 - p_0 z^{-1})(1 - p_0^* z^{-1})}$$


× pole
• zero




A constructive Solution to the Spectral Factorization Problem

Pole-zero pattern $P_x(z)$

× pole
• zero




Stable pole-zero part of $P_x(z)$:



$$Q(z) = \frac{(1 - z_1 z^{-1})(1 - z_2 z^{-1})}{(1 - p_0 z^{-1})(1 - p_0^* z^{-1})}$$

Unstable pole-zero part of $P_x(z)$:



$$Q^*\left(\frac{1}{z^*}\right) = \frac{(1 - z_1 z)(1 - z_2 z)}{(1 - p_0^* z)(1 - p_0 z)}$$

A constructive Spectral Factorization Method (Ct'd)

The finalize the calculations we need to fix the gain such that:

$$\lim_{z \rightarrow \alpha} P_x(z) = \sigma \lim_{z \rightarrow \alpha} Q(z) Q^*\left(\frac{1}{z^*}\right)$$

Example 2: Consider the case displayed in the last figure and take $\alpha = 1$:

$$\begin{aligned} \lim_{z \rightarrow 1} Q(z) &= \lim_{z \rightarrow 1} \frac{(1 - z_1 z^{-1})(1 - z_2 z^{-1})}{(1 - p_0 z^{-1})(1 - p_0^* z^{-1})} \\ &= \frac{(1 - z_1)(1 - z_2)}{(1 - p_0)(1 - p_0^*)} \\ \lim_{z \rightarrow 1} Q^*\left(\frac{1}{z^*}\right) &= \lim_{z \rightarrow 1} \frac{(1 - z_1 z)(1 - z_2 z)}{(1 - p_0^* z)(1 - p_0 z)} \\ &= \frac{(1 - z_1)(1 - z_2)}{(1 - p_0^*)(1 - p_0)} \end{aligned}$$

From this follows $\sigma \left(\frac{(1 - z_1)(1 - z_2)}{(1 - p_0)(1 - p_0^*)} \right)^2 = \lim_{z \rightarrow 1} P_x(z)$.

Solving the Spectral Factorization problem

1. Illustrations of Key Definitions: A rational transfer function, a minimum phase transfer function, $Q(\infty) = 1$.
2. Use of the Spectral Factor
3. Special form of the power spectrum $P_x(z)$.
 - Constraints on its coefficients
 - Constraints on its poles and zeros
4. A constructive Solution
5. **Illustration**

Illustration

$$P_x(z) = \frac{4}{(1 - 0.5z^{-1})(1 - 0.5z)}$$

This (rational) function has the poles 0.5 and 2. Therefore,

$$Q(z) = \frac{1}{1 - 0.5z^{-1}} \Rightarrow Q^*\left(\frac{1}{z^*}\right) = \frac{1}{1 - 0.5z}$$

With the spectral factorization given as:

$$P_x(z) = \sigma Q(z)Q^*(1/z^*)$$

we can find σ e.g. by considering the above equation for $z = 1$:

$$16 = \sigma 4 \Rightarrow \sigma = 4$$

Example 3

Problem: Given $P_x(\omega) = \frac{25-24\cos\omega}{26-10\cos\omega}$ for $\omega \in [0, \pi]$. Determine a realization of $x(n)$?

Solution:

1. Determine the z-transform $P_x(z)$.
2. Determine a spectral factorization of $P_x(z) = \sigma Q(z)Q^*(\frac{1}{z^*})$.
3. Then filtering a ZMWN(1) sequence $v(n)$ with the filter $\sqrt{\sigma}Q(z)$ delivers a signal that has the given spectrum.

Ex3_26.m

Generalization to non-rational Spectra (optional)

Generalization to non-rational Spectra

Recall the Kolmogorov turbulence spectra:

$$P_\phi(f) = 0.023 r_0^{-\frac{5}{3}} f^{-\frac{11}{3}}$$

In order to apply the constructive method for rational Power spectra, we approximate the given Power spectra by a rational function first.

Rational Approximation (Example)

Let the rational approximation be denoted as $\hat{P}_x(z) = \frac{N(z)}{D(z)}$, with e.g.

$$N(z) = az + b + a^*z^{-1} \quad D(z) = dz + e + d^*z^{-1}$$

Then the coefficients $a, d \in \mathbb{C}, b, e \in \mathbb{R}$ are found by solving the following optimization problem,

$$\min_{a,b,d,e} \frac{1}{2\pi} \int_{-\pi}^{\pi} \left| P_x(\omega) - \frac{N(e^{j\omega})}{D(e^{j\omega})} \right|^2 d\omega$$

Remark: This is a non-linear optimization problem.

[Intermezzo] Subspace Identification

The problem of approximating multivariable Power spectra where $P_x(e^{j\omega})$ is a matrix by a rational matrix function has been successfully addressed in our team.

`test.m`

More info provided in the course `sc42025`

The Wold Decomposition

Regular RPs

DEFINITION: A **regular** WSS random process has a continuous Power Spectrum $P(e^{j\omega})$ in ω and this power spectrum has a spectral factorization:

$$P_x(z) = \sigma_0^2 Q(z) Q^*\left(\frac{1}{z^*}\right)$$

The Wold Decomposition

WOLD DECOMPOSITION THEOREM: A general WSS random process can be written as,

$$x(n) = x_p(n) + x_r(n)$$

where $x_r(n)$ is a regular RP and $x_p(n)$ is a predictable process, with

$$E[x_r(m)x_p^*(n)] = 0 \quad \forall m, n \quad \text{i.e. } x_r(m), x_p(n) \text{ are orthogonal}$$

COROLLARY: For a general WSS RP $x(n)$:

$$P_x(e^{j\omega}) = \sigma_0^2 |Q(e^{j\omega})|^2 + \sum_{k=1}^N \alpha_k \delta(\omega - \omega_k)$$

Key message from Part I

1. If $P_x(z)$ is polynomial in z with the property:

$$P_x(z) = P_x^*\left(\frac{1}{z^*}\right)$$

then $P_x(z)$ is fully known from its causal part only.

Example 0: Let $P_x(z) = az + b + cz^{-1}$ then

$$a = c^* \quad b = b^*$$

Outline of Part II

1. **Problem formulation for ARMA(p,q)-model**
2. From $r_x(k)$ to the parameters $a(i)$, $|b(0)|$ of an AR(p)-model
3. Spectral factorization to determine a set of $b(j)$ parameters of an MA(q)-model.
4. From $r_x(k)$ to the parameters $a(i)$, $b(j)$ and the impulse response parameters $h(m)$ of an ARMA(p,q)-model:
 - The set of equations to determine its $a(i)$ -parameters.
 - Retrieving its $b(i)$ and impulse response parameters $h(m)$ using spectral factorization.

Problem Formulation for ARMA(p,q)

Consider the Yule-Walker Equations (Lecture 3) of an ARMA(p,q) model with impulse response parameters $h(m)$ (for $m = 0 : q$):

$$r_x(k) + \sum_{\ell=1}^p a(\ell)r_x(k-\ell) = \begin{cases} \sum_{\ell=k}^q b(\ell)h^*(\ell-k) & : 0 \leq k \leq q \\ 0 & : k > q \end{cases}$$

then given the Autocorrelation function $r_x(k)$

1. Determine the number N of samples of the Autocorrelation function $r_x(k)$ for $k = 0 : N$ necessary to determine the parameters $a(i), b(j)$.
2. Determine the parameters $a(i), b(j)$ for $i = 1 : p, j = 0 : q$ assuming $\sigma_v = 1$ such that the Autocorrelation from the derived ARMA(p,q) model equals $r_x(k)$.

The Yule-Walker equations for AR(p) models

Recall from Lecture 3 (slide 41) that for the AR(p) the

Yule-Walker equations reads:

$$r_x(k) + \sum_{\ell=1}^p a(\ell)r_x(k - \ell) = |b(0)|^2 \Delta(k) \quad k \geq 0$$

If we take $k = 1 : p$ we have p -equations:

$$\begin{bmatrix} r_x(1) & r_x(0) & r_x^*(1) & \cdots & r_x^*(p-1) \\ r_x(2) & r_x(1) & r_x(0) & \cdots & r_x^*(p-2) \\ \vdots & & & \ddots & \\ r_x(p) & r_x(p-1) & r_x(p-2) & \cdots & r_x(0) \end{bmatrix} \begin{bmatrix} 1 \\ a(1) \\ a(2) \\ \vdots \\ a(p) \end{bmatrix} = 0$$

The Yule-Walker equations for AR(p) models

Recall from Lecture 3 (slide 41) that for the AR(p) the

Yule-Walker equations reads:

$$r_x(k) + \sum_{\ell=1}^p a(\ell)r_x(k - \ell) = |b(0)|^2 \Delta(k) \quad k \geq 0$$

If we take $k = 1 : p$ we have p -equations:

$$\begin{bmatrix} r_x(0) & r_x^*(1) & \cdots & r_x^*(p-1) \\ r_x(1) & r_x(0) & \cdots & r_x^*(p-2) \\ \vdots & & \ddots & \\ r_x(p-1) & r_x(p-2) & \cdots & r_x(0) \end{bmatrix} \begin{bmatrix} a(1) \\ a(2) \\ \vdots \\ a(p) \end{bmatrix} = - \begin{bmatrix} r_x(1) \\ r_x(2) \\ \vdots \\ r_x(p) \end{bmatrix}$$

From which the parameters $a(i)$ for $i = 1 : p$ can be estimated provided the “system matrix” is invertible.

The Yule-Walker equations for AR(p) models

Recall from Lecture 3 (slide 43) that for the AR(p) the

Yule-Walker equations reads:

$$r_x(k) + \sum_{\ell=1}^p a(\ell)r_x(k - \ell) = |b(0)|^2 \Delta(k) \quad k \geq 0$$

Knowing the $a(i)$'s, we take $k = 0$ to yield the parameter $|b(0)|$ as,

$$|b(0)|^2 = r_x(0) + \sum_{\ell=1}^p a(\ell)r_x^*(\ell)$$

Example 1: AR(1) model

Consider the (real) Autocorrelation function given by:

$$r_x(k) = \frac{4}{3} \left(\frac{1}{2} \right)^{|k|}$$

then determine the parameter a and b in the AR(1) model:

$$x(n) + ax(n-1) = bv(n) \quad v(n) \text{ is ZMWN}(1)$$

Example 1: AR(1) model (Ct'd)

Consider the (real) Autocorrelation function given by:

$$r_x(k) = \frac{4}{3} \left(\frac{1}{2}\right)^{|k|}$$

For $k = 1$ in the Yule-Walker equation, we have the single equation:

$$r_x(1) + ar_x(0) = 0 \Rightarrow a = -\frac{r_x(1)}{r_x(0)}$$

For $k = 0$ in the Yule-Walker equation, we have:

$$r_x(0) + ar_x(1) = |b|^2 \Rightarrow |b|^2 = r_x(0) - \frac{r_x(1)^2}{r_x(0)} \Rightarrow |b| = \sqrt{\frac{r_x(0)^2 - r_x(1)^2}{r_x(0)}}$$

Example 1: AR(1) model (Ct'd)

Consider the (real) Autocorrelation function given by:

$$r_x(k) = \frac{4}{3} \left(\frac{1}{2} \right)^{|k|}$$

For $k = 1$ in the Yule-Walker equation, we have the single equation:

$$r_x(1) + ar_x(0) = 0 \Rightarrow a = -\frac{r_x(1)}{r_x(0)} = -\frac{1}{2}$$

For $k = 0$ in the Yule-Walker equation, we have:

$$r_x(0) + ar_x(1) = |b|^2 \Rightarrow |b|^2 = r_x(0) - \frac{r_x(1)^2}{r_x(0)} \Rightarrow |b| = \sqrt{\frac{r_x(0)^2 - r_x(1)^2}{r_x(0)}} = 1$$

Outline of Part II

1. Problem formulation for ARMA(p,q)-model
2. From $r_x(k)$ to the parameters $a(i)$, $|b(0)|$ of an AR(p)-model
3. **Spectral factorization to determine a set of $b(j)$ parameters of an MA(q)-model.**
4. From $r_x(k)$ to the parameters $a(i)$, $b(j)$ and the impulse response parameters $h(m)$ of an ARMA(p,q)-model:
 - The set of equations to determine its $a(i)$ -parameters.
 - Retrieving its $b(i)$ and impulse response parameters $h(m)$ using spectral factorization.

The Yule-Walker equations for MA(q) models

Recall from Lecture 3 (slide 39) that for the MA(q) the

Yule-Walker equations reads:

$$r_x(k) = \sum_{\ell=k}^q b(\ell)b^*(\ell - k) \quad 0 \leq k \leq q$$

(otherwise the $r_x(k)$'s are zero.). How to find from $r_x(k)$ a set of parameters $b(j)$ for $j = 1 : q$ such that the above equation holds?

The Yule-Walker equations for MA(q) models

Recall from Lecture 3 (slide 39) that for the MA(q) the

Yule-Walker equations reads:

$$r_x(k) = \sum_{\ell=k}^q b(\ell)b^*(\ell-k) \quad 0 \leq k \leq q$$

Define $r_\infty(k)$ and $b_\infty(j)$ as follows,

$$r_\infty(k) = \begin{cases} r_x(k) & k = -q : q \\ 0 & \text{otherwise} \end{cases} \quad b_\infty(j) = \begin{cases} b(j) & j = 0 : q \\ 0 & \text{otherwise} \end{cases}$$

Then,

$$P_\infty(z) = \sum_{k=-\infty}^{\infty} \sum_{\ell=-\infty}^{\infty} b_\infty(\ell)b_\infty^*(\ell-k)z^{-k} = \sum_{\ell=-\infty}^{\infty} b_\infty(\ell)z^{-\ell} \sum_{m=-\infty}^{\infty} b_\infty^*(m)z^m$$

for $m = \ell - k$. Or $P_\infty(z) = B_\infty(z)B_\infty^*(1/z^*)$.

The Yule-Walker equations for MA(q) models

Recall from Lecture 3 (slide 39) that for the MA(q) the

Yule-Walker equations reads:

$$r_x(k) = \sum_{\ell=k}^q b(\ell)b^*(\ell-k) \quad 0 \leq k \leq q$$

Define $r_\infty(k)$ and $b_\infty(j)$ as follows,

$$r_\infty(k) = \begin{cases} r_x(k) & k = -q : q \\ 0 & \text{otherwise} \end{cases} \quad b_\infty(j) = \begin{cases} b(j) & j = 0 : q \\ 0 & \text{otherwise} \end{cases}$$

Then,

$$P_\infty(z) = \sum_{k=-\infty}^{\infty} \sum_{\ell=-\infty}^{\infty} b_\infty(\ell)b_\infty^*(\ell-k)z^{-k} = \sum_{\ell=-\infty}^{\infty} b_\infty(\ell)z^{-\ell} \sum_{m=-\infty}^{\infty} b_\infty^*(m)z^m$$

for $m = \ell - k$. Or $P_x(z) = B(z)B^*(1/z^*)$.

The Yule-Walker equations for MA(q) models

Recall from Lecture 3 (slide 39) that for the MA(q) the

Yule-Walker equations reads:

$$r_x(k) = \sum_{\ell=k}^q b(\ell)b^*(\ell - k) \quad 0 \leq k \leq q$$

Therefore,

$$P_x(z) = B(z)B^*(1/z^*)$$

Knowing $r_x(k)$ we know $P_x(z)$ (a finite order polynomial in z) and the determination of $B(z)$ is done via a

The Yule-Walker equations for MA(q) models

Recall from Lecture 3 (slide 39) that for the MA(q) the

Yule-Walker equations reads:

$$r_x(k) = \sum_{\ell=k}^q b(\ell)b^*(\ell - k) \quad 0 \leq k \leq q$$

Therefore,

$$P_x(z) = B(z)B^*(1/z^*)$$

Knowing $r_x(k)$ we know $P_x(z)$ (a finite order polynomial in z) and the determination of $B(z)$ is done via a **Spectral Factorization**.

Example 2: MA(1) model

Consider the (real) Autocorrelation function given by:

$$r_x(-1) = -\frac{1}{2} \quad r_x(0) = \frac{5}{4} \quad r_x(1) = -\frac{1}{2}$$

Then $P_x(z) = -\frac{1}{2}z + \frac{5}{4} - \frac{1}{2}z^{-1}$ The roots of this polynomial are 2 and $\frac{1}{2}$. Therefore,

$$P_x(z) = \alpha(1 - \frac{1}{2}z^{-1})(1 - \frac{1}{2}z)$$

Considering $z = 1$ yields $\alpha = 1$.

Example 2: MA(1) model

Consider the (real) Autocorrelation function given by:

$$r_x(-1) = -\frac{1}{2} \quad r_x(0) = \frac{5}{4} \quad r_x(1) = -\frac{1}{2}$$

Then $P_x(z) = -\frac{1}{2}z + \frac{5}{4} - \frac{1}{2}z^{-1}$ The roots of this polynomial are 2 and $\frac{1}{2}$. Therefore,

$$P_x(z) = \alpha(1 - \frac{1}{2}z^{-1})(1 - \frac{1}{2}z)$$

Considering $z = 1$ yields $\alpha = 1$. Concluding: The parameter $b(0) = 1$ and $b(1) = -\frac{1}{2}$.

Outline Part II

1. Problem formulation for ARMA(p,q)-model
2. From $r_x(k)$ to the parameters $a(i)$, $|b(0)|$ of an AR(p)-model
3. Spectral factorization to determine a set of $b(j)$ parameters of an MA(q)-model.
4. **From $r_x(k)$ to the parameters $a(i)$, $b(j)$ and the impulse response parameters $h(m)$ of an ARMA(p,q)-model:**
 - The set of equations to determine its $a(i)$ -parameters.
 - Retrieving its $b(i)$ and impulse response parameters $h(m)$ using spectral factorization.

Estimating the $a(i)$'s of an ARMA(p,q) model

Recall the **Yule-Walker equations**

$$r_x(k) + \sum_{\ell=1}^p a(\ell)r_x(k-\ell) = \begin{cases} \sum_{\ell=k}^q b(\ell)h^*(\ell-k) & : 0 \leq k \leq q \\ 0 & : k > q \end{cases}$$

If we take $k = q+1, \dots, q+p$ we obtain the p equations:

$$\begin{bmatrix} r_x(q+1) & r_x(q) & \cdots & r_x(q-p+1) \\ \vdots & & \ddots & \\ r_x(q+p) & r_x(q+p-1) & \cdots & r_x(q) \end{bmatrix} \begin{bmatrix} 1 \\ a(1) \\ \vdots \\ a(p) \end{bmatrix} = 0$$

Estimating the $a(i)$'s of an ARMA(p,q) model

Recall the **Yule-Walker equations**

$$r_x(k) + \sum_{\ell=1}^p a(\ell)r_x(k-\ell) = \begin{cases} \sum_{\ell=k}^q b(\ell)h^*(\ell-k) & : 0 \leq k \leq q \\ 0 & : k > q \end{cases}$$

If we take $k = q + 1, \dots, q + p$ we obtain the p equations:

$$\begin{bmatrix} r_x(q) & \cdots & r_x(q-p+1) \\ \vdots & \ddots & \\ r_x(q+p-1) & \cdots & r_x(q) \end{bmatrix} \begin{bmatrix} a(1) \\ \vdots \\ a(p) \end{bmatrix} = - \begin{bmatrix} r_x(q+1) \\ \vdots \\ r_x(q+p) \end{bmatrix}$$

These are called the **modified Yule-Walker equations**.

Knowing the $a(i)$'s how to find $b(j)$'s?

Recall the **Yule-Walker equations**

$$r_x(k) + \sum_{\ell=1}^p a(\ell)r_x(k-\ell) = \sum_{\ell=k}^q b(\ell)h^*(\ell-k)$$

If we know the $a(i)$'s and take $k = 0 : q$,

Knowing the $a(i)$'s how to find $b(j)$'s?

Recall the **Yule-Walker equations**

$$c(k) = r_x(k) + \sum_{\ell=1}^p a(\ell)r_x(k-\ell) = \sum_{\ell=k}^q b(\ell)h^*(\ell-k)$$

If we know the $a(i)$'s and take $k = 0 : q$,

Knowing the $a(i)$'s how to find $b(j)$'s?

Recall the **Yule-Walker equations**

$$r_x(k) + \sum_{\ell=1}^p a(\ell)r_x(k - \ell) = \sum_{\ell=k}^q b(\ell)h^*(\ell - k) = c(k)$$

If we know the $a(i)$'s and take $k = 0 : q$, we can find the coefficients $c(k)$ for $k = 0 : q$ from,

$$\begin{bmatrix} r_x(0) & r_x^*(1) & \cdots & r_x^*(p) \\ r_x(1) & r_x(0) & \cdots & r_x^*(p-1) \\ \vdots & & \ddots & \\ r_x(q) & r_x(q+1) & \cdots & r_x(0) \end{bmatrix} \begin{bmatrix} 1 \\ a(1) \\ \vdots \\ a(p) \end{bmatrix} = \begin{bmatrix} c(0) \\ c(1) \\ \vdots \\ c(q) \end{bmatrix}$$

Finding the $b(j)$'s

Status: We know the coefficients $c(k)$ for $k = 0 : q$ and their expression:

$$c(k) = \sum_{\ell=k}^q b(\ell)h^*(\ell - k) \quad (1)$$

Finding the $b(j)$'s

Status: We know the coefficients $c(k)$ for $k = 0 : q$ and their expression:

$$c(k) = \sum_{\ell=k}^q b(\ell)h^*(\ell - k) \quad (1)$$

From this expression for $c(k)$ we find that for $k > q$ (i.e. $k = q + 1, q + 2, \dots, \infty$), $c(k) = 0$.

Finding the $b(j)$'s

Status: We know the coefficients $c(k)$ for $k = 0 : q$ and their expression:

$$c(k) = \sum_{\ell=k}^q b(\ell)h^*(\ell - k) \quad (1)$$

From this expression for $c(k)$ we find that for $k > q$ (i.e. $k = q + 1, q + 2, \dots, \infty$), $c(k) = 0$.

However from $C(z)$ only the causal part is known. The anti-causal part is unknown and (likely) non-zero. For example,

$$c(-1) = b(-1)h^*(0) + b(0)h^*(1) + \dots + b(q)h^*(q + 1)$$

Finding the $b(j)$'s

Status: We know the coefficients $c(k)$ for **only** for $k = 0 : \infty$ and their expression:

$$c(k) = \sum_{\ell=k}^q b(\ell) h^*(\ell - k) \quad (1)$$

Finding the $b(j)$'s

Status: We know the coefficients $c(k)$ for **only** for $k = 0 : \infty$ and their expression:

$$c(k) = \sum_{\ell=k}^q b(\ell)h^*(\ell - k) \quad (1)$$

From (1) and the following “embedding” (like in the MA(q)-case):

$$\{b(n)\}_{n=-\infty}^{\infty} = \begin{cases} 0 & n < 0 \\ b(n) & 0 \leq n \leq q \\ 0 & n > q \end{cases} \quad \{h(n)\}_{n=-\infty}^{\infty} = \begin{cases} 0 & n < 0 \\ h(n) & 0 \leq n \end{cases}$$

we find that $\boxed{C(z) = B(z)H^*(1/z^*)}$ (2)

But from $C(z)$ only the causal part is known.

Finding the $b(j)$'s

Status: Let $C(z)$ be denoted as,

$$C(z) = \sum_{k=-\infty}^{\infty} c(k)z^{-k} = \underbrace{\sum_{k=1}^{\infty} c(-k)z^k}_{[C(z)]_-} + \underbrace{\sum_{k=0}^{\infty} c(k)z^{-k}}_{[C(z)]_+}$$

with $[C(z)]_+$ known. Further a combination of the expression
 $C(z) = B(z)H^*(1/z^*)$

Finding the $b(j)$'s

Status: Let $C(z)$ be denoted as,

$$C(z) = \sum_{k=-\infty}^{\infty} c(k)z^{-k} = \underbrace{\sum_{k=1}^{\infty} c(-k)z^k}_{[C(z)]_-} + \underbrace{\sum_{k=0}^{\infty} c(k)z^{-k}}_{[C(z)]_+}$$

with $[C(z)]_+$ known. Further a combination of the expression $C(z) = B(z)H^*(1/z^*)$ with the definition $H(z) = \frac{B(z)}{A(z)}$, yields

Finding the $b(j)$'s

Status: Let $C(z)$ be denoted as,

$$C(z) = \sum_{k=-\infty}^{\infty} c(k)z^{-k} = \underbrace{\sum_{k=1}^{\infty} c(-k)z^k}_{[C(z)]_-} + \underbrace{\sum_{k=0}^{\infty} c(k)z^{-k}}_{[C(z)]_+}$$

with $[C(z)]_+$ known. Further a combination of the expression $C(z) = B(z)H^*(1/z^*)$ with the definition $H(z) = \frac{B(z)}{A(z)}$, yields

$$C(z) = B(z) \frac{B^*(1/z^*)}{A^*(1/z^*)} \Rightarrow C(z)A^*(1/z^*) = B(z)B^*(1/z^*) = P_y(z)$$

Finding the $b(j)$'s

Status: Let $C(z)$ be denoted as,

$$C(z) = \sum_{k=-\infty}^{\infty} c(k)z^{-k} = \underbrace{\sum_{k=1}^{\infty} c(-k)z^k}_{[C(z)]_-} + \underbrace{\sum_{k=0}^{\infty} c(k)z^{-k}}_{[C(z)]_+}$$

with $[C(z)]_+$ known. Further a combination of the expression $C(z) = B(z)H^*(1/z^*)$ with the definition $H(z) = \frac{B(z)}{A(z)}$, yields

$$C(z) = B(z) \frac{B^*(1/z^*)}{A^*(1/z^*)} \Rightarrow C(z)A^*(1/z^*) = B(z)B^*(1/z^*) = P_y(z)$$

Since $P_y(z)$ satisfies $P_y(z) = P_y^*(1/z^*)$ it is fully defined by its causal part. Can we find that causal part and subsequently determine $B(z)$ by **Spectral Factorization**?

The causal part of $P_y(z)$

With $P_y(z)$ equal to $C(z)A^*(1/z^*)$ we have,

$$\begin{aligned} P_y(z) &= C(z)A^*(1/z^*) \quad \text{with} \quad A^*(1/z^*) = 1 + a^*(1)z + \cdots + a^*(p)z^p \\ &= \left([C(z)]_+ + [C(z)]_- \right) A^*(1/z^*) \\ &= [C(z)]_+ A^*(1/z^*) + [C(z)]_- A^*(1/z^*) \end{aligned}$$

Therefore,

$$[P_y(z)]_+ = \left[[C(z)]_+ A^*(1/z^*) \right]_+$$

Let us further illustrate this calculation and the solution for $B(z)$ by an example.

Example 3 to find the Power spectrum $P_y(z)$ and its min-phase factor

Given: $r_x(0) = 26$ $r_x(1) = 7$ $r_x(2) = 3.5$, From the

Yule-Walker equations we find that for $p = 1$, $a(1) = -\frac{1}{2}$, and as such,

$$A(z) = 1 - \frac{1}{2}z^{-1}$$

Example 3 to find the Power spectrum $P_y(z)$ and its min-phase factor

Given: $r_x(0) = 26$ $r_x(1) = 7$ $r_x(2) = 3.5$, From the

Yule-Walker equations we find that for $p = 1$, $a(1) = -\frac{1}{2}$, and as such,

$$A(z) = 1 - \frac{1}{2}z^{-1}$$

Again from the Yule-Walker equations we find,

$$[C(z)]_+ = 22.5 - 6z^{-1} \Rightarrow$$

Example 3 to find the Power spectrum $P_y(z)$ and its min-phase factor

Given: $r_x(0) = 26$ $r_x(1) = 7$ $r_x(2) = 3.5$, From the

Yule-Walker equations we find that for $p = 1$, $a(1) = -\frac{1}{2}$, and as such,

$$A(z) = 1 - \frac{1}{2}z^{-1}$$

Again from the Yule-Walker equations we find,

$$\begin{aligned} [C(z)]_+ &= 22.5 - 6z^{-1} \Rightarrow \\ [P_y(z)]_+ &= \left[[C(z)]_+ A^*(1/z^*) \right]_+ = \left[\left(\frac{45}{2} - 6z^{-1} \right) \left(1 - \frac{1}{2}z \right) \right]_+ \\ &= \left[-\frac{45}{4}z + \left(\frac{45}{2} + \frac{6}{2} \right) - 6z^{-1} \right]_+ = \frac{51}{2} - 6z^{-1} \end{aligned}$$

and $P_y(z) = -6z + \frac{51}{2} - 6z^{-1}$. The spectral factorization of $P_y(z) = B(z)B^*(1/z^*)$ is,

$$P_y(z) = 24\left(1 - \frac{1}{4}z^{-1}\right)\left(1 - \frac{1}{4}z\right) \Rightarrow B(z) = 2\sqrt{6}\left(1 - \frac{1}{4}z^{-1}\right) \text{inv_ARMA.m}$$