

Phenotypically anchored mRNA and miRNA expression profiling in zebrafish reveals flame retardant chemical toxicity networks

Subham Dasgupta¹, Cheryl L. Dunham¹, Lisa Truong¹, Michael T. Simonich¹, Robyn L. Tanguay^{1*}

¹Department of Environmental and Molecular Toxicology, Sinnhuber Aquatic Research Laboratory, Oregon State University, Corvallis, Oregon, USA

Supplemental Tables- 3

Supplemental Figures- 1

Supplemental table legends:

Table S1- Phenotypic, behavioral, mRNA and miR data for the selected FRCs. Phenotyping and behavioral data represented as lowest effect levels (LELs, Table S1.1). mRNA and miR data represented as log₂ fold changes (Tables S1.2 and S1.3). Blank cells represent fold changes below statistical or fold change thresholds.

Table S2- TFmiR miR-TF-miR interaction data for all FRCs. Only experimentally validated data (Table S2.1), both experimentally validated and computationally predicted data (Table S2.2-2.9)

Table S3. Gene Ontology data for mRNA (Tables S3.1-3.10) and mRNA targets of miRs (Table S3.11)

Supplemental Figure legends

Figure S1. Heatmap representing log₂ fold changes of (A) all differentially expressed mRNAs and (B) all differentially expressed miRs across FRCs.

Figure S1

