

## HW 0

Kevin Lin

1/12/2026

### 1

- (a) Let  $x_1, \dots, x_n$  be real values. Then for the quadratic function  $f(\theta) = \sum_{i=1}^n w_i(x_i - \theta)^2$  where  $w_i > 0$  for all  $i$ , the optimal solution  $\theta^*$  denoted by  $\theta^* = \arg \min_{\theta} f(\theta)$  can be calculated as follows:

$$\begin{aligned}\frac{d}{d\theta} f(\theta) &= \frac{d}{d\theta} \sum_{i=1}^n w_i(x_i - \theta)^2 \\ &= \sum_{i=1}^n w_i \cdot 2(x_i - \theta) \cdot (-1) \\ &= -2 \sum_{i=1}^n w_i(x_i - \theta) \\ &= -2 \left( \sum_{i=1}^n w_i x_i - \theta \sum_{i=1}^n w_i \right)\end{aligned}$$

Setting the derivative to zero to find the minimum:

$$\begin{aligned}-2 \left( \sum_{i=1}^n w_i x_i - \theta \sum_{i=1}^n w_i \right) &= 0 \\ \sum_{i=1}^n w_i x_i - \theta \sum_{i=1}^n w_i &= 0 \\ \theta \sum_{i=1}^n w_i &= \sum_{i=1}^n w_i x_i \\ \theta^* &= \frac{\sum_{i=1}^n w_i x_i}{\sum_{i=1}^n w_i}\end{aligned}$$

Thus, the optimal solution is the weighted average of the  $x_i$ 's. If some weights are negative, the function may not be convex, and the solution may not correspond to a minimum.

- (b) (i) Given  $2n$  kids are randomly divided into two equal subgroups, the probability that the two tallest kids end up in the the same subgroup can be calculated as follows:

$$\begin{aligned}
P(\text{tallest in same group}) &= P(\text{both in group 1}) + P(\text{both in group 2}) \\
&= \frac{\binom{2n-2}{n-2}}{\binom{2n}{n}} + \frac{\binom{2n-2}{n-2}}{\binom{2n}{n}} \\
&= 2 \cdot \frac{\binom{2n-2}{n-2}}{\binom{2n}{n}} \\
&= 2 \cdot \frac{\frac{(2n-2)!}{(n-2)!(n)!}}{\frac{(2n)!}{(n)!(n)!}} \\
&= 2 \cdot \frac{(2n-2)!n!}{(n-2)!(2n)!} \\
&= 2 \cdot \frac{n(n-1)}{(2n)(2n-1)} \\
&= \frac{n(n-1)}{(2n-1)(n)} \\
&= \frac{n-1}{2(2n-1)}
\end{aligned}$$

- (ii) The probability that the two tallest kids end up in different subgroups is:

$$\begin{aligned}
P(\text{tallest in different groups}) &= 1 - P(\text{both tallest in same group}) \\
&= 1 - \frac{n-1}{2(2n-1)} \\
&= \frac{2(2n-1) - (n-1)}{2(2n-1)} \\
&= \frac{4n-2-n+1}{2(2n-1)} \\
&= \frac{3n-1}{2(2n-1)}
\end{aligned}$$

- (c) We know  $P(\text{knows answer}) = p$ , and  $P(\text{doesn't know answer}) = 1 - p$ . Also,  $P(\text{correct}|\text{knows answer}) = 0.99$  and  $P(\text{correct}|\text{doesn't know answer}) = 1/k$ . Then  $P(\text{knew answer}|\text{correct})$  can be calculated using Bayes' The-

orem:

$$\begin{aligned}
 P(\text{knew answer}|\text{correct}) &= \frac{P(\text{correct}|\text{knew answer}) \cdot P(\text{knew answer})}{P(\text{correct})} \\
 &= \frac{P(\text{correct}|\text{knew answer}) \cdot P(\text{knew answer})}{P(\text{correct}|\text{knew answer}) \cdot P(\text{knew answer}) + P(\text{correct}|\text{didn't know answer}) \cdot P(\text{didn't know answer})} \\
 &= \frac{0.99 \cdot p}{0.99 \cdot p + \frac{1}{k} \cdot (1 - p)}
 \end{aligned}$$

**2**

(a) a