

MATH 1530 Problem Set 3

Tanish Makadia

(Collaborated with Esmé, Marcos, Edward, and Kazuya)

February 2023

Problem 1. Please complete the mid-semester survey. Write “I have completed the mid-semester survey” and sign your name.

Problem 2. Let \mathbf{a} be an element of a group G . Prove that $\langle \mathbf{a}^m \rangle \cap \langle \mathbf{a}^n \rangle$ is cyclic, where n, m are integers. What is its generator?

Proof. Let $\mathbf{a}^k \in \langle \mathbf{a}^m \rangle \cap \langle \mathbf{a}^n \rangle$. We have that $\mathbf{a}^k \in \langle \mathbf{a}^m \rangle \implies \mathbf{a}^k = \mathbf{a}^{ms}$ where $s \in \mathbb{Z}$. We also have that $\mathbf{a}^k \in \langle \mathbf{a}^n \rangle \implies \mathbf{a}^k = \mathbf{a}^{nt}$ where $t \in \mathbb{Z}$. Together, we have

$$\mathbf{a}^k = \mathbf{a}^{ms} = \mathbf{a}^{nt} \implies k = ms = nt$$

In other words, k must be a common multiple of both m and n . Since every common multiple of m and n is itself a multiple of $\text{lcm}(m, n)$, we have that $\langle \mathbf{a}^m \rangle \cap \langle \mathbf{a}^n \rangle$ is equal to $\{(\mathbf{a}^{\text{lcm}(m, n)})^c \mid c \in \mathbb{Z}\} = \langle \mathbf{a}^{\text{lcm}(m, n)} \rangle$. \square

Problem 3. Let \mathbf{a} and \mathbf{b} belong to a group. If $|\mathbf{a}|$ and $|\mathbf{b}|$ are relatively prime, prove that $\langle \mathbf{a} \rangle \cap \langle \mathbf{b} \rangle = \{\mathbf{e}\}$.

Proof. Let G be a group containing elements \mathbf{a}, \mathbf{b} . Let $\mathbf{m} = |\mathbf{a}|$ and $\mathbf{n} = |\mathbf{b}|$. We can now express $\langle \mathbf{a} \rangle$ and $\langle \mathbf{b} \rangle$ as:

$$\langle \mathbf{a} \rangle = \{\mathbf{e}, \mathbf{a}^1, \dots, \mathbf{a}^{\mathbf{m}-1}\} \quad \langle \mathbf{b} \rangle = \{\mathbf{e}, \mathbf{b}^1, \dots, \mathbf{b}^{\mathbf{n}-1}\}$$

Because the identity element of G is unique, we have that $\mathbf{e} \in \langle \mathbf{a} \rangle \cap \langle \mathbf{b} \rangle$.

Next, we will show that for all $\mathbf{a}^k \in \langle \mathbf{a} \rangle$ such that $\mathbf{a}^k \neq \mathbf{e}$, we have that $\mathbf{a}^k \notin \langle \mathbf{b} \rangle$. By (Gallian, 4.2 Corollary 1), we know that if $\mathbf{a}^k \in \langle \mathbf{a} \rangle$, then $|\mathbf{a}^k|$ divides \mathbf{m} . Additionally, since $\mathbf{a}^k \neq \mathbf{e}$, we know $|\mathbf{a}^k| > 1$. If $\mathbf{a}^k \in \langle \mathbf{b} \rangle$, $|\mathbf{a}^k|$ must divide \mathbf{n} . But since $|\mathbf{a}|$ and $|\mathbf{b}|$ are relatively prime, we have that $\gcd(\mathbf{m}, \mathbf{n}) = 1$. Because $|\mathbf{a}^k| \neq 1$, we have shown that $\mathbf{a}^k \notin \langle \mathbf{b} \rangle$. The same process can be used to show that for all $\mathbf{b}^k \in \langle \mathbf{b} \rangle$ such that $\mathbf{b}^k \neq \mathbf{e}$, we have that $\mathbf{b}^k \notin \langle \mathbf{a} \rangle$.

Therefore, we have proven that $\langle \mathbf{a} \rangle \cap \langle \mathbf{b} \rangle = \{\mathbf{e}\}$. □

Problem 4. Let G be an Abelian group of order 77, and assume that for all $x \in G$, we have that $x^{77} = e$. Prove that G is cyclic.

Proof. For all $x \in G$, we have $x^{77} = e$ which implies that $|x|$ divides 77. Thus, for all $x \in G$, we have that $|x| \in \{1, 7, 11, 77\}$. To prove that G is cyclic, we must show that G has an element of order 77.

Suppose that every element of G besides $e \in G$ has order 7. Because G is closed, it must contain every cyclic subgroup generated by these elements. For all $x \in G$, we have $|x| = |\langle x \rangle|$ and $e \in \langle x \rangle$. This implies that each cyclic subgroup generated by an element of order 7 contains 6 distinct non-identity elements. Thus, the number of distinct elements in G is $|G| = 77 = 1 + 6n$ where $n \in \mathbb{Z}$. This is a contradiction since $6 \nmid 76$.

Alternatively, suppose every element of G besides $e \in G$ has order 11. We have that each cyclic subgroup generated by an element of order 11 contains 10 distinct non-identity elements. In this case, $|G| = 77 = 1 + 10n$ where $n \in \mathbb{Z}$. This is also a contradiction since $10 \nmid 76$.

We are therefore left with the following cases:

- **Case 1 (G has an element of order 7 and order 11):** Let $a, b \in G$ such that $|a| = 7$ and $|b| = 11$. Thus, $a^{77} = a^7 = e$ and $b^{77} = b^{11} = e$. This implies that $(ab)^{77} = a^{77}b^{77} = e^2 = e$. Hence, we have that $|ab|$ divides 77, giving us the following four cases:
 - **Case 1 ($|ab| = 1$):** This implies $ab = e$ which is a contradiction since a and b do not have the same order.
 - **Case 2 ($|ab| = 7$):** This implies $e = (ab)^7 = a^7b^7 = e \cdot b^7 = b^7$. This is a contradiction since $|b| = 11$.
 - **Case 3 ($|ab| = 11$):** This implies $e = (ab)^{11} = a^{11}b^{11} = a^{11} \cdot e = a^{11}$. This is a contradiction since $7 \nmid 11$.
 - **Case 4 ($|ab| = 77$):** By process of elimination, we have that $|ab| = 77$.
- **Case 2 (G has an element of order 77):** We have that G can be generated by this element and we are done.

Since both cases lead to the existence of an element of order 77 in G , we have proven that G is cyclic. \square