

MATH 1530 Problem Set 3

Tanish Makadia

(Collaborated with Esmé and Kazuya)

February 2023

Problem 1. Consider $\mathbb{U}(40)$. Find a subgroup which is cyclic of order 4. Find a subgroup which is noncyclic of order 4.

Proof. $\mathbb{U}(40)$ is defined as follows:

$$\mathbb{U}(40) = \{1, 3, 7, 9, 11, 13, 17, 19, 21, 23, 27, 29, 31, 33, 37, 39\}$$

- **Cyclic subgroup:** We will create a cyclic subgroup of order 4 using 7 as the generator.

$$\langle 7 \rangle = \{7, 9, 23, 1\} \quad (\text{Gallian, 3.4})$$

- **Non-cyclic subgroup:** We will show that $\{1, 9, 11, 19\}$ is a noncyclic subgroup.

		1	9	11	19
	1	1	9	11	19
Cayley Table:	9	9	1	19	11
	11	11	19	1	9
	19	19	11	9	1

Since $\{1, 9, 11, 19\}$ is finite and closed under multiplication mod 40, we have proven that it is subgroup of $\mathbb{U}(40)$. Additionally, since $|9| = |11| = |19| = 2$, there is no element in this group which can generate the entire group, proving that it is noncyclic.

□

Problem 2. If H and K are subgroups of a group G , prove that $H \cap K$ is a subgroup of G . If $H \not\subseteq K$ and $K \not\subseteq H$, prove that $H \cup K$ is never a subgroup of G .

1. $H \cap K$ is a subgroup of G .

Proof. We will use the two-step subgroup test.

- **Closed over inverses:** $x \in H \cap K \implies x \in H$ and $x \in K$. Since H and K are subgroups, we have the existence of $x^{-1} \in H$ and $x^{-1} \in K \implies x^{-1} \in H \cap K$.
- **Closed under group operation:** $a, b \in H \cap K \implies a, b \in H$ and $a, b \in K$. Since H and K are subgroups, we have that $ab \in H$ and $ab \in K \implies ab \in H \cap K$.

□

2. If $H \not\subseteq K$ and $K \not\subseteq H$, $H \cup K$ is never a subgroup of G .

Proof.

□

Problem 3. Prove that a group G is Abelian if and only if $G = Z(G)$.

Problem 4. Suppose G is a group with exactly 8 elements of order 3. how many subgroups of order 3 does G have?

Proof. Let $H \subset G$ be a subgroup of order 3:

$$H = \{e, a, b\}$$

Since e is unique, we have that $ab \neq a$ and $ab \neq b$. In order for H to be closed, the only remaining choice is $ab = e$. Thus, for any subgroup of order 3, the two elements besides the identity must be each other's inverse.

Now, we will show that $|a| = |b| = 3$. Consider $a^2 \in H$. Since the identity is unique, $a^2 \neq a$, and since b is the unique inverse of a , the only remaining choice is $a^2 = b$. Therefore,

$$\begin{aligned} a^3 &= a \cdot a^2 & b^3 &= (a^2)^3 \\ &= a \cdot b & &= (a^3)^2 \\ &= e & &= e \end{aligned}$$

We have that there are exactly 8 elements of G of order 3. Because a subgroup of order 3 requires two distinct elements of order 3, we can conclude that the number of distinct subgroups of order 3 in G is $8/2 = 4$. \square

Problem 5. Let G be a finite group with more than one element. Show that G has an element of prime order.