

MATHEMATICS II

(Analysis + Linear Algebra)

Partial Vols → Chp-10

sequences & series (of real numbers & complex numbers)

$\mathbb{R} \xrightarrow{f} \mathbb{R}$ $a \in \mathbb{R} \rightarrow f(a) \in f(\mathbb{R})$ $\rightarrow f \rightarrow \text{convergent (or) not}$ prove

Def: A function $f: \mathbb{N} \rightarrow \mathbb{R} (or \mathbb{C})$ is called an (infinite) sequence (real sequence / complex sequence)

Denote $\{f(n)\}_{n \in \mathbb{N}} = \{a_n\}_{n \in \mathbb{N}}$ $a_n = n^{\text{th}}$ term of sequence $a_n \in \mathbb{R} (or \mathbb{C})$

Examples: $\{2^n\}_{n \in \mathbb{N}}$ $\{\frac{1}{n}\}_{n \in \mathbb{N}}$ $\{\frac{1}{2^n}\}_{n \in \mathbb{N}}$

$a_1 = 1$ $a_n = a_{n-1} + a_{n-2}$ $n \geq 3$ Fibonacci series

Def: A (real (or) complex) sequence $\{a_n\}$ has a limit L ($L \in \mathbb{R} (or \mathbb{C})$) if for every positive number $\epsilon > 0$ there exists $N \in \mathbb{N}$ (N may depend on ϵ) such that

$|a_n - L| < \epsilon$ whenever $n \geq N$

Notations: $\lim_{n \rightarrow \infty} a_n = L$ or $(a_n \rightarrow L)$ as $(n \rightarrow \infty)$

$a_n = 1 + \frac{1}{n} + \frac{1}{n^2} + \dots + \frac{1}{n^i}$ $a_n \rightarrow \sum_{n=0}^{\infty} \frac{1}{n^i} = e$

if $\{a_n\}$ has a limit L then say that $\{a_n\}_{n \in \mathbb{N}}$ is convergent sequence else $\{a_n\}_{n \in \mathbb{N}}$ is divergent sequence.

a convergent sequence if $\exists L \in \mathbb{R}$ s.t. $\forall \epsilon > 0$

$\{a_n\}_{n \in \mathbb{N}}$ is convergent if $\exists N$ where $|a_n - L| < \epsilon$

$\exists N = N(\epsilon) \in \mathbb{N}$ with $n \geq N$ where $|a_n - L| < \epsilon$

$\{a_n\}_{n \in \mathbb{N}}$ is divergent if $\nexists L \in \mathbb{R}$ $\exists \epsilon > 0$ such that $\forall n \in \mathbb{N}$ $\exists n \geq N$ with $|a_n - L| \geq \epsilon$

divergent

Defn: $\lim_{n \rightarrow \infty} a_n = +\infty$ if given any $M \in \mathbb{Z}$ $\exists N \in \mathbb{N}$ such that

$$\forall n \geq N, a_n \geq M$$

$$\text{Ex: } a_n = n, a_n = 2^n, a_n = -n$$


$a_n \in \mathbb{R}, n \geq 0$ if given any $M \in \mathbb{Z}$ $\exists N \in \mathbb{N}$

s.t. $\forall n \geq N$ hence $a_n \in \mathbb{N}$



$$a_n = \lfloor \log(n) \rfloor, n \in \mathbb{N}$$

$L = \lim_{n \rightarrow \infty} a_n$ if given $\epsilon > 0$ $\exists N \in \mathbb{N}$ s.t. $|a_n - L| < \epsilon$ whenever $n \geq N$



$a_n \in \mathbb{R}$ say that $\lim_{n \rightarrow \infty} a_n = \infty$ if $\lim_{n \rightarrow \infty} |a_n| = +\infty$

$f: \mathbb{N} \rightarrow \mathbb{R}^k$ $\{f(n)\}_{n \in \mathbb{N}} = \{a_n\}_{n \in \mathbb{N}}$

$a_n = (a_{n1}, a_{n2}, \dots, a_{nk})$, $i = 1, \dots, k$
 $\lim_{n \rightarrow \infty} a_n = L$ i.e. $\lim_{n \rightarrow \infty} a_{ni} = L_i$

Defn: $\lim_{n \rightarrow \infty} a_n = L$ if given any $\epsilon > 0$ $\exists N \in \mathbb{N}$ s.t. $|a_n - L| < \epsilon$ whenever $n \geq N$

$$\lim_{n \rightarrow \infty} \frac{1}{n} = 0 \quad (1 < 1)$$

$$\lim_{n \rightarrow \infty} x^n = 0$$

$$\lim_{n \rightarrow \infty} n^n = 1$$

$$\lim_{n \rightarrow \infty} \left(\frac{1}{n}\right)^n = 0$$

$$\lim_{n \rightarrow \infty} \left(\frac{1}{n}\right)^n = e^0$$

$$\lim_{n \rightarrow \infty} \frac{n!}{n^n} = 1$$

$$\lim_{n \rightarrow \infty} \frac{n!}{(1-p)^n} = 0$$

Defn: A real sequence $\{a_n\}$ is increasing (decreasing) if

$a_n \leq a_{n+1}$ ($a_n \geq a_{n+1}$) $\forall n$

bounded if $\exists M > 0$ s.t. $|a_n| < M$ $\forall n \in \mathbb{N}$

Ex: $(-1)^n$ is ldd but not convergent

Thm: Monotonic seq. converges \Leftrightarrow ldd

conv seq \Rightarrow ldd seq (doesn't require monotonic)

Given a (real/complex) $\{a_n\}_{n \in \mathbb{N}}$ define $S_n = a_1 + a_2 + \dots + a_n$

S_n is partial sum of $\{a_n\}_{n \in \mathbb{N}}$. $\{S_n\}_{n \in \mathbb{N}}$ is

called an (infinite) series denoted by $\sum_{n=1}^{\infty} a_n$

if $\lim_{n \rightarrow \infty} S_n = S$ exists ($S \in \mathbb{R}$ or \mathbb{C}) then we say that the

series $\sum_{n=1}^{\infty} a_n$ is convergent & denote $\sum_{n=1}^{\infty} a_n = S$

converges to S

divergent series $\left(\sum_{n=1}^{\infty} \frac{1}{2^n} = 1\right)$

$\left(\sum_{n=1}^{\infty} \frac{1}{n} = \infty\right)$

$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$

Sanj dhera given

Thm 10.1 Let $\alpha_1, \dots, \alpha_n$ and $\beta_1, \dots, \beta_n \in \text{conv}(\text{real}(\alpha_i) \text{ complex})$
 satisfy $\alpha_i \neq \beta_i \in \mathbb{R} \cap (\alpha_i, \beta_i)$. Then $\sum_{i=1}^n (\alpha_i + \beta_i) \lambda_i$ is a convex
 & is equal to $\sum_{i=1}^n \alpha_i \lambda_i + \sum_{i=1}^n \beta_i \lambda_i$ (proof) (proof)

Thm 10.3) S_n converges, S_n diverges $\Rightarrow (S_n + b_n)$ diverges

Thm 104) $\{a_n\}$ $\{b_n\}$ 2 sequences (of reals) converging to \lim

$s.t. \quad a_n = b_n - b_{n+1} \quad \forall n \in \mathbb{N}$
 Then $\sum_{n=1}^{\infty} a_n$ converges \Leftrightarrow the seq $\{b_n\}_{n \in \mathbb{N}}$ converges
 & in such a case $\sum_{n=1}^{\infty} a_n = b_1 - \lim_{n \rightarrow \infty} b_n$

Thus $\{a_n\}$ is odd seq. \Rightarrow a subsequence of $\{a_n\}$ which converges. (B-W Thm) - proof

Subsequences $a_1, a_2, \dots, a_{n_1}, \dots$ If n_k is of a then consequently sub-sequence is a with order as given

⑧ $\{a_n\}_{n \in \mathbb{N}}$ a seq. of rational number $\{ \exists m \in \mathbb{N}$

S.t. $\{a_n\}_{n \in \mathbb{N}}$ is rational (Not works for above Thm)
 $(x_2 = 1.4142 \dots)$ is something extra that it converges to irrational

$$a_1 \quad a_2 \quad a_3 \quad a_4 \quad \dots \quad a_n \quad \text{dist.}$$
$$\lim_{n \rightarrow \infty} a_n = \sqrt{2}$$

then every subseq. $\{a_{n_k}\}$ of $\{a_n\}$ also converges to L .

$\{a_n\}$, $\{b_n\}$ are convergent then $\{a_n + b_n\}$ and $\{a_n - b_n\}$ are convergent. Proof

convergent $\left\{ \begin{array}{l} \text{series} \\ \{a_n\} \end{array} \right\} \rightarrow \text{convergent}$
 $\left\{ \begin{array}{l} \perp \\ a_n \neq 0 \ \& \ \underline{a \neq 0} \end{array} \right\}$

$$pmbn = k$$

Taken Σ (an+b) - average

$\sum (a_n b_n) + (-a_n) \rightarrow$ contradiction

as given Ans over

leaf of Thun 10.4)

$$\sum_{n=1}^{\infty} a_n = \lim_{n \rightarrow \infty} (s_n) = \lim_{n \rightarrow \infty} (b_n - b_{n+1})$$
$$\dim (b_1 - b_{n+1}) \text{ exists} \Rightarrow \dim (b_{n+1} - b_1) \text{ exists}$$

Alle (M(b) exists or M(b) not)

3. 1. 2. 3. 4. 5. 6. 7. 8. 9. 10. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 45. 46. 47. 48. 49. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60. 61. 62. 63. 64. 65. 66. 67. 68. 69. 70. 71. 72. 73. 74. 75. 76. 77. 78. 79. 80. 81. 82. 83. 84. 85. 86. 87. 88. 89. 90. 91. 92. 93. 94. 95. 96. 97. 98. 99. 100. 101. 102. 103. 104. 105. 106. 107. 108. 109. 110. 111. 112. 113. 114. 115. 116. 117. 118. 119. 120. 121. 122. 123. 124. 125. 126. 127. 128. 129. 130. 131. 132. 133. 134. 135. 136. 137. 138. 139. 140. 141. 142. 143. 144. 145. 146. 147. 148. 149. 150. 151. 152. 153. 154. 155. 156. 157. 158. 159. 160. 161. 162. 163. 164. 165. 166. 167. 168. 169. 170. 171. 172. 173. 174. 175. 176. 177. 178. 179. 180. 181. 182. 183. 184. 185. 186. 187. 188. 189. 190. 191. 192. 193. 194. 195. 196. 197. 198. 199. 200. 201. 202. 203. 204. 205. 206. 207. 208. 209. 210. 211. 212. 213. 214. 215. 216. 217. 218. 219. 220. 221. 222. 223. 224. 225. 226. 227. 228. 229. 230. 231. 232. 233. 234. 235. 236. 237. 238. 239. 240. 241. 242. 243. 244. 245. 246. 247. 248. 249. 250. 251. 252. 253. 254. 255. 256. 257. 258. 259. 260. 261. 262. 263. 264. 265. 266. 267. 268. 269. 270. 271. 272. 273. 274. 275. 276. 277. 278. 279. 280. 281. 282. 283. 284. 285. 286. 287. 288. 289. 290. 291. 292. 293. 294. 295. 296. 297. 298. 299. 300. 301. 302. 303. 304. 305. 306. 307. 308. 309. 310. 311. 312. 313. 314. 315. 316. 317. 318. 319. 320. 321. 322. 323. 324. 325. 326. 327. 328. 329. 330. 331. 332. 333. 334. 335. 336. 337. 338. 339. 340. 341. 342. 343. 344. 345. 346. 347. 348. 349. 350. 351. 352. 353. 354. 355. 356. 357. 358. 359. 360. 361. 362. 363. 364. 365. 366. 367. 368. 369. 370. 371. 372. 373. 374. 375. 376. 377. 378. 379. 380. 381. 382. 383. 384. 385. 386. 387. 388. 389. 390. 391. 392. 393. 394. 395. 396. 397. 398. 399. 400. 401. 402. 403. 404. 405. 406. 407. 408. 409. 410. 411. 412. 413. 414. 415. 416. 417. 418. 419. 420. 421. 422. 423. 424. 425. 426. 427. 428. 429. 430. 431. 432. 433. 434. 435. 436. 437. 438. 439. 440. 441. 442. 443. 444. 445. 446. 447. 448. 449. 450. 451. 452. 453. 454. 455. 456. 457. 458. 459. 460. 461. 462. 463. 464. 465. 466. 467. 468. 469. 470. 471. 472. 473. 474. 475. 476. 477. 478. 479. 480. 481. 482. 483. 484. 485. 486. 487. 488. 489. 490. 491. 492. 493. 494. 495. 496. 497. 498. 499. 500. 501. 502. 503. 504. 505. 506. 507. 508. 509. 510. 511. 512. 513. 514. 515. 516. 517. 518. 519. 520. 521. 522. 523. 524. 525. 526. 527. 528. 529. 530. 531. 532. 533. 534. 535. 536. 537. 538. 539. 540. 541. 542. 543. 544. 545. 546. 547. 548. 549. 550. 551. 552. 553. 554. 555. 556. 557. 558. 559. 560. 561. 562. 563. 564. 565. 566. 567. 568. 569. 570. 571. 572. 573. 574. 575. 576. 577. 578. 579. 580. 581. 582. 583. 584. 585. 586. 587. 588. 589. 590. 591. 592. 593. 594. 595. 596. 597. 598. 599. 600. 601. 602. 603. 604. 605. 606. 607. 608. 609. 610. 611. 612. 613. 614. 615. 616. 617. 618. 619. 620. 621. 622. 623. 624. 625. 626. 627. 628. 629. 630. 631. 632. 633. 634. 635. 636. 637. 638. 639. 640. 641. 642. 643. 644. 645. 646. 647. 648. 649. 650. 651. 652. 653. 654. 655. 656. 657. 658. 659. 660. 661. 662. 663. 664. 665. 666. 667. 668. 669. 670. 671. 672. 673. 674. 675. 676. 677. 678. 679. 680. 681. 682. 683. 684. 685. 686. 687. 688. 689. 690. 691. 692. 693. 694. 695. 696. 697. 698. 699. 700. 701. 702. 703. 704. 705. 706. 707. 708. 709. 710. 711. 712. 713. 714. 715. 716. 717. 718. 719. 720. 721. 722. 723. 724. 725. 726. 727. 728. 729. 730. 731. 732. 733. 734. 735. 736. 737. 738. 739. 740. 741. 742. 743. 744. 745. 746. 747. 748. 749. 750. 751. 752. 753. 754. 755. 756. 757. 758. 759. 760. 761. 762. 763. 764. 765. 766. 767. 768. 769. 770. 771. 772. 773. 774. 775. 776. 777. 778. 779. 780. 781. 782. 783. 784. 785. 786. 787. 788. 789. 790. 791. 792. 793. 794. 795. 796. 797. 798. 799. 800. 801. 802. 803. 804. 805. 806. 807. 808. 809. 810. 811. 812. 813. 814. 815. 816. 817. 818. 819. 820. 821. 822. 823. 824. 825. 826. 827. 828. 829. 830. 831. 832. 833. 834. 835. 836. 837. 838. 839. 840.

$$\sum_{n \in \mathbb{N}} a_n = \lim_{n \rightarrow \infty} (b_n - b_{n+1}) = \lim_{n \rightarrow \infty} b_n - \lim_{n \rightarrow \infty} b_{n+1}$$
$$= b_1 - \lim_{n \rightarrow \infty} b_n$$

10.5.0 of 'x' is a complex number with $|x| < 1$ then

geometric series $\sum_{n=0}^{\infty} x^n$ converges \iff has sum $\frac{1}{1-x}$

this series $\sum_{n=0}^{\infty} x^n$ diverges

2

10

3. $\sum_{n=0}^{\infty} \left(\frac{1}{2} + \frac{1}{2}i\right)^n = \frac{1}{1 - \left(\frac{1}{2} + \frac{1}{2}i\right)}$

proofs $S_n = 1 + x + x^2 + \dots + x^n = \frac{1-x^{n+1}}{1-x}$ ($x \neq 1$)
 so ($x \neq 1$)

$(1+x+x^2+\dots+x^n)(1-x) = 1-x^{n+1} = 1-x^{n+1}$
 $= 1-x^{n+1}$

$S_n = \frac{1-x^{n+1}}{1-x}$ ($x \neq 1$)

lim S_n exists $\Leftrightarrow \left\{ \frac{x^{n+1}}{1-x} \right\}_{n \in \mathbb{N}}$ limit exists

$|x| < 1$
 $\{x^{n+1}\}_{n \in \mathbb{N}}$ $\{x^{n+1}\}_{n \in \mathbb{N}}$

we are looking at the sequence $\{a_n = x^{n+1}\}_{n \in \mathbb{N}}$

want to show $\lim_{n \rightarrow \infty} x^{n+1} = 0$

ETP given $\epsilon > 0$ $\exists N \in \mathbb{N}$ s.t. $\forall n \geq N$ we should have $|x^{n+1} - 0| = |x^{n+1}| < \epsilon$

$x \in \mathbb{C}$ $|x|^n = |x|^n$
 $|x|^{n+1} < \epsilon$

$|x| = |x|$
 $|x|^{n+1} < \epsilon$
 $|x| < 1$ $\exists N$

$|x| < 1$ for any $\epsilon > 0$

$n \geq N \Rightarrow \log \epsilon$
 $\log \epsilon$

$|x| < 1$ $\forall n \geq N$

$\sum_{n=0}^{\infty} x^n$ converges if equal to $\frac{1}{1-x}$ for all x s.t. $|x| < 1$
 $\sum_{n=0}^{\infty} x^n \Rightarrow \sum_{n=0}^{\infty} x^n$ diverges $S_n = 1 + x + \dots + x^n = \frac{1-x^{n+1}}{1-x}$
 $\sum_{n=0}^{\infty} x^n = \lim_{n \rightarrow \infty} \frac{1-x^{n+1}}{1-x} = \frac{1}{1-x} - \frac{x^{n+1}}{1-x}$

$\{x^{n+1}\}_{n \in \mathbb{N}}$ diverges.

$|x| > 1$ $|x^{n+1}| = |x|^{n+1} \rightarrow \infty$

proof: $\sum_{n=1}^{\infty} a_n$ converges ($a_n \in \mathbb{R}$ or \mathbb{C})
 $\Rightarrow a_n \rightarrow 0$ as $n \rightarrow \infty$

$S_{n+1} - S_n = a_{n+1}$
 $(a_1 + \dots + a_{n+1}) - (a_1 + a_2 + \dots + a_n)$

$\lim_{n \rightarrow \infty} (S_{n+1} - S_n) = \lim_{n \rightarrow \infty} S_{n+1} - \lim_{n \rightarrow \infty} S_n = 0$

$\sum_{n=0}^{\infty} \frac{2^n + (-2)^n}{3^{n+1} + (-2)^{n+1}} = a_n$ Does $\{a_n\}$ converge (or diverge)

$\lim_{n \rightarrow \infty} \frac{2^n (1 + (-2)^n)}{3^{n+1} (1 + (-2)^n)} = \lim_{n \rightarrow \infty} \left(\frac{1}{3} \right)^{n+1} \left[\frac{1 + (-2)^n}{1 + (-2)^n} \right]$

converges to $\frac{1}{3}$

$$\sum_{n=1}^{\infty} \frac{1}{n} = \infty \quad \text{if } \lim_{n \rightarrow \infty} (1 + \frac{1}{n} + \dots + \frac{1}{n}) = \infty$$

$$\log \frac{1}{x} = -\log x = -\int_x^1 \frac{1}{t} dt$$



Example: $a_n = \frac{1}{n+1} = \frac{1}{n} - \frac{1}{n+1} = b_n - b_{n+1}$

by then $\sum_{n=1}^N a_n = b_1 - b_{N+1} = 1 - \frac{1}{N+1} \rightarrow 1$

Example: x is not a negative integer $(x \neq -1, -2, \dots)$

$$\sum_{n=0}^{\infty} \frac{1}{(n+x)(n+x+1)} = \frac{1}{x} \left[\frac{1}{n+x} - \frac{1}{n+x+1} \right]$$

if $n \in \mathbb{N}$

if $x \in \mathbb{C}$ $\sum_{n=0}^{\infty} \frac{1}{(n+x)(n+x+1)}$ converges (or not / compare test)

if $x \in \mathbb{R}$ $\sum_{n=0}^{\infty} \frac{1}{(n+x)(n+x+1)}$ converges (or not / compare test)

Proof: $\sum_{n=1}^{\infty} S_{n+1} - S_n = a_{n+1}$ exist \S are equal

if $a_n \geq 0$ $\lim_{n \rightarrow \infty} a_n = 0$

if $a_n \geq 0$

$$1 + x + x^2 + \dots + x^n = \frac{1-x^{n+1}}{1-x} \quad \text{if } x \neq 1$$

$$1 + x + x^2 + \dots + x^n = \frac{1-x^{n+1}}{1-x} \quad \text{if } x \neq 1$$

$$1 + x + x^2 + \dots + x^n = \frac{1-x^{n+1}}{1-x} \quad \text{if } x \neq 1$$

$$1 + x + x^2 + \dots + x^n = \frac{1-x^{n+1}}{1-x} \quad \text{if } x \neq 1$$

$$1 + x + x^2 + \dots + x^n = \frac{1-x^{n+1}}{1-x} \quad \text{if } x \neq 1$$

$$1 + x + x^2 + \dots + x^n = \frac{1-x^{n+1}}{1-x} \quad \text{if } x \neq 1$$

$$1 + x + x^2 + \dots + x^n = \frac{1-x^{n+1}}{1-x} \quad \text{if } x \neq 1$$

$$1 + x + x^2 + \dots + x^n = \frac{1-x^{n+1}}{1-x} \quad \text{if } x \neq 1$$

$$1 + x + x^2 + \dots + x^n = \frac{1-x^{n+1}}{1-x} \quad \text{if } x \neq 1$$

$$1 + x + x^2 + \dots + x^n = \frac{1-x^{n+1}}{1-x} \quad \text{if } x \neq 1$$

$$1 + x + x^2 + \dots + x^n = \frac{1-x^{n+1}}{1-x} \quad \text{if } x \neq 1$$

$$1 + x + x^2 + \dots + x^n = \frac{1-x^{n+1}}{1-x} \quad \text{if } x \neq 1$$

$$1 + x + x^2 + \dots + x^n = \frac{1-x^{n+1}}{1-x} \quad \text{if } x \neq 1$$

$$1 + x + x^2 + \dots + x^n = \frac{1-x^{n+1}}{1-x} \quad \text{if } x \neq 1$$

$$1 + x + x^2 + \dots + x^n = \frac{1-x^{n+1}}{1-x} \quad \text{if } x \neq 1$$

$$1 + x + x^2 + \dots + x^n = \frac{1-x^{n+1}}{1-x} \quad \text{if } x \neq 1$$

Thm 10.3 Assume $a_n \geq 0$ & $n \in \mathbb{N}$. Then $\sum_{n=1}^{\infty} a_n$ converges $\Leftrightarrow \{a_n\}$ is a bounded above sequence.

Proof: It's obvious $a_n \geq 0$, $n \in \mathbb{N}$, $\sum_{n=1}^{\infty} a_n$ converges.

Thm 10.4 Assume $a_n \geq 0$, $n \in \mathbb{N}$, and $\sum_{n=1}^{\infty} a_n$ converges. Then $\sum_{n=1}^{\infty} a_n c_n$ converges.

Proof: $a_n = a_1 + \dots + a_n$, $c_n = c_1 + \dots + c_n$. $\sum_{n=1}^{\infty} a_n c_n$ converges.

Thm 10.5 Assume $a_n \geq 0$, $n \in \mathbb{N}$, and $\sum_{n=1}^{\infty} a_n$ converges. Then $\sum_{n=1}^{\infty} a_n c_n$ converges.

Proof: $a_n = a_1 + \dots + a_n$, $c_n = c_1 + \dots + c_n$. $\sum_{n=1}^{\infty} a_n c_n$ converges.

Thm 10.6 Assume $a_n \geq 0$, $n \in \mathbb{N}$, and $\sum_{n=1}^{\infty} a_n$ converges. Then $\sum_{n=1}^{\infty} a_n c_n$ converges.

Thm 10.7 Assume $a_n \geq 0$, $n \in \mathbb{N}$, and $\sum_{n=1}^{\infty} a_n$ converges. Then $\sum_{n=1}^{\infty} a_n c_n$ converges.

Ex: $\sum_{k=1}^{\infty} \frac{1}{k!} \leq \frac{1}{2 \times 1} = \frac{1}{2}$. $\sum_{k=1}^{\infty} \frac{1}{k!} \leq \frac{1}{2}$.

Ex: $\sum_{k=1}^{\infty} \frac{1}{k!} \leq \frac{1}{2}$. $\sum_{k=1}^{\infty} \frac{1}{k!} \leq \frac{1}{2}$.

Ex: $\sum_{k=1}^{\infty} \frac{1}{k!} \leq \frac{1}{2}$. $\sum_{k=1}^{\infty} \frac{1}{k!} \leq \frac{1}{2}$.

Ex: $\sum_{k=1}^{\infty} \frac{1}{k!} \leq \frac{1}{2}$. $\sum_{k=1}^{\infty} \frac{1}{k!} \leq \frac{1}{2}$.

Ex: $\sum_{k=1}^{\infty} \frac{1}{k!} \leq \frac{1}{2}$. $\sum_{k=1}^{\infty} \frac{1}{k!} \leq \frac{1}{2}$.

Thm 10.8 Assume $a_n \geq 0$, $n \in \mathbb{N}$, and $\sum_{n=1}^{\infty} a_n$ converges. Then $\sum_{n=1}^{\infty} a_n c_n$ converges.

Thm 10.9 Assume $a_n \geq 0$, $n \in \mathbb{N}$, and $\sum_{n=1}^{\infty} a_n$ converges. Then $\sum_{n=1}^{\infty} a_n c_n$ converges.

Thm 10.10 Assume $a_n \geq 0$, $n \in \mathbb{N}$, and $\sum_{n=1}^{\infty} a_n$ converges. Then $\sum_{n=1}^{\infty} a_n c_n$ converges.

Thm 10.11 Assume $a_n \geq 0$, $n \in \mathbb{N}$, and $\sum_{n=1}^{\infty} a_n$ converges. Then $\sum_{n=1}^{\infty} a_n c_n$ converges.

Thm 10.12 Assume $a_n \geq 0$, $n \in \mathbb{N}$, and $\sum_{n=1}^{\infty} a_n$ converges. Then $\sum_{n=1}^{\infty} a_n c_n$ converges.

Thm 10.13 Assume $a_n \geq 0$, $n \in \mathbb{N}$, and $\sum_{n=1}^{\infty} a_n$ converges. Then $\sum_{n=1}^{\infty} a_n c_n$ converges.

Thm 10.14 Assume $a_n \geq 0$, $n \in \mathbb{N}$, and $\sum_{n=1}^{\infty} a_n$ converges. Then $\sum_{n=1}^{\infty} a_n c_n$ converges.

Thm 10.15 Assume $a_n \geq 0$, $n \in \mathbb{N}$, and $\sum_{n=1}^{\infty} a_n$ converges. Then $\sum_{n=1}^{\infty} a_n c_n$ converges.

Thm 10.16 Assume $a_n \geq 0$, $n \in \mathbb{N}$, and $\sum_{n=1}^{\infty} a_n$ converges. Then $\sum_{n=1}^{\infty} a_n c_n$ converges.

Thm 10.17 Assume $a_n \geq 0$, $n \in \mathbb{N}$, and $\sum_{n=1}^{\infty} a_n$ converges. Then $\sum_{n=1}^{\infty} a_n c_n$ converges.

Thm 10.18 Assume $a_n \geq 0$, $n \in \mathbb{N}$, and $\sum_{n=1}^{\infty} a_n$ converges. Then $\sum_{n=1}^{\infty} a_n c_n$ converges.

Thm 10.19 Assume $a_n \geq 0$, $n \in \mathbb{N}$, and $\sum_{n=1}^{\infty} a_n$ converges. Then $\sum_{n=1}^{\infty} a_n c_n$ converges.

Thm 10.20 Assume $a_n \geq 0$, $n \in \mathbb{N}$, and $\sum_{n=1}^{\infty} a_n$ converges. Then $\sum_{n=1}^{\infty} a_n c_n$ converges.

Thm 10.21 Assume $a_n \geq 0$, $n \in \mathbb{N}$, and $\sum_{n=1}^{\infty} a_n$ converges. Then $\sum_{n=1}^{\infty} a_n c_n$ converges.

Example 5: $\sum_{n=1}^{\infty} \frac{1}{(\log n)^n}$ converges.

$a_n = \frac{1}{(\log n)^n}$

$(a_n)^{1/n} = \frac{1}{\log n} \rightarrow \frac{1}{\log n} < 1$

Example 6: $\sum_{n=1}^{\infty} \left(\frac{n}{n+1}\right)^n$ converges.

$a_n = \left(\frac{n}{n+1}\right)^n = \frac{1}{\left(1+\frac{1}{n}\right)^n} \rightarrow \frac{1}{e} < 1$

Then ratio test $\left(\frac{a_{n+1}}{a_n}\right)^{1/(n+1)} > \frac{1}{e}$ is a series of the terms

1. $\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = L$ then

2. $L < 1 \Rightarrow \sum a_n$ conv

3. $L < 1 \Rightarrow \sum a_n$ div

4. $L > 1 \Rightarrow \sum a_n$ is nonconvergent

5. $L = 1 \Rightarrow$ test is inconclusive

6. $\frac{a_{n+1}}{a_n} \leq \frac{1}{n}$ div $\sum \frac{1}{n}$ conv

7. $\frac{a_{n+1}}{a_n} = \frac{n}{n+1} \rightarrow 1$ as $n \rightarrow \infty$

8. $\frac{a_{n+1}}{a_n} = \frac{n^2}{(n+1)^2} \rightarrow 1$ as $n \rightarrow \infty$

9. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n}} \rightarrow 1$ as $n \rightarrow \infty$

10. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n^2}} \rightarrow 1$ as $n \rightarrow \infty$

11. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n^3}} \rightarrow 1$ as $n \rightarrow \infty$

12. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n^4}} \rightarrow 1$ as $n \rightarrow \infty$

13. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n^5}} \rightarrow 1$ as $n \rightarrow \infty$

14. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n^6}} \rightarrow 1$ as $n \rightarrow \infty$

15. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n^7}} \rightarrow 1$ as $n \rightarrow \infty$

16. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n^8}} \rightarrow 1$ as $n \rightarrow \infty$

17. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n^9}} \rightarrow 1$ as $n \rightarrow \infty$

18. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n^{10}}} \rightarrow 1$ as $n \rightarrow \infty$

19. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n^{11}}} \rightarrow 1$ as $n \rightarrow \infty$

20. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n^{12}}} \rightarrow 1$ as $n \rightarrow \infty$

21. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n^{13}}} \rightarrow 1$ as $n \rightarrow \infty$

22. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n^{14}}} \rightarrow 1$ as $n \rightarrow \infty$

23. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n^{15}}} \rightarrow 1$ as $n \rightarrow \infty$

24. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n^{16}}} \rightarrow 1$ as $n \rightarrow \infty$

25. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n^{17}}} \rightarrow 1$ as $n \rightarrow \infty$

26. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n^{18}}} \rightarrow 1$ as $n \rightarrow \infty$

27. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n^{19}}} \rightarrow 1$ as $n \rightarrow \infty$

28. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n^{20}}} \rightarrow 1$ as $n \rightarrow \infty$

29. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n^{21}}} \rightarrow 1$ as $n \rightarrow \infty$

30. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n^{22}}} \rightarrow 1$ as $n \rightarrow \infty$

31. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n^{23}}} \rightarrow 1$ as $n \rightarrow \infty$

32. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n^{24}}} \rightarrow 1$ as $n \rightarrow \infty$

33. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n^{25}}} \rightarrow 1$ as $n \rightarrow \infty$

34. $\frac{a_{n+1}}{a_n} = \frac{1}{1+\frac{1}{n^{26}}} \rightarrow 1$ as $n \rightarrow \infty$

$$\lim_{n \rightarrow \infty} \delta_n = \delta^1 \quad \left\{ \begin{array}{l} \text{with these limits exist} \\ \lim_{n \rightarrow \infty} \delta_{2n+1} = \delta^0 \end{array} \right.$$

$$\delta_{2n+1} - \delta_{2n} = \delta_{2n+1}$$

$$\lim_{n \rightarrow \infty} \delta_{2n+1} - \lim_{n \rightarrow \infty} \delta_{2n} = \lim_{n \rightarrow \infty} \delta_{2n+1} = 0$$

$$\lim_{n \rightarrow \infty} \delta_{2n+1} = \lim_{n \rightarrow \infty} \delta_{2n}$$

$$\delta^1 = \delta^0$$

$$\lim_{n \rightarrow \infty} S_{2n+1} = S = \lim_{n \rightarrow \infty} S_{2n}$$

$$\lim_{n \rightarrow \infty} S_n = S$$

$$\epsilon > 0 \quad n \geq N_1 \Rightarrow |\delta_{2n+1} - \delta| < \epsilon \quad N = \max(N_1, N_2)$$

$$n \geq N_2 \Rightarrow |\delta_{2n} - \delta| < \epsilon$$

$$\left(\begin{array}{c} \delta - \epsilon \\ \delta \end{array} \right) \delta \epsilon$$

$$\Rightarrow \lim_{n \rightarrow \infty} S_n = \delta \quad \lim_{n \rightarrow \infty} S_n = \sum_{n=1}^{\infty} (-1)^{n+1} a_n = \delta$$

$$\delta_2 \quad \delta_4 \quad \delta_6 \quad \delta_{2n} \quad \delta_{2n+1} \quad \delta_5 \quad \delta_1$$

$$\delta_{2n} < \delta_{2n+1} < \delta < \delta_{2n+1} < \delta_{2n+1}$$

$$\Rightarrow 0 < \delta_{2n} \leq \delta_{2n+1} - \delta_{2n} = a_{2n+1}$$

$$\Rightarrow 0 < \delta_{2n+1} - \delta \leq \delta_{2n+1} - \delta_{2n} = a_{2n+1}$$

$$\Rightarrow 0 < (-1)^n (a_{2n} \delta_n) < a_{n+1}$$

Example
 $\sum_{n=1}^{\infty} (-1)^n \log n$ converges.

Leibniz rule $\log n \rightarrow 0$ as $n \rightarrow \infty$ & $\log(n)$ is a decreasing function

$$\lim_{n \rightarrow \infty} \frac{(\log n)^4}{n!} = 0 \quad \text{for } n \geq 20 \quad \text{with a calculator (on Friday)}$$

$$f(x) = \log x \quad \text{for } x \geq 20$$

$$f'(x) = \frac{1}{x} \log x < 0 \quad \text{if } x \geq 20$$

$$\Rightarrow f(n+1) < f(n) \quad \text{for } n \geq 20$$

Example: Define a real sequence $\{a_n\}_{n \in \mathbb{N}}$ as follows

$$a_{2n+1} = \frac{1}{n!}$$

$$a_{2n} = \int_0^1 \frac{t^{n-1} dt}{x}$$

$$a_2 = \int_0^1 \frac{t^2}{x} dt, a_4 = \int_0^1 \frac{t^4}{x} dt, a_6 = \int_0^1 \frac{t^6}{x} dt, \dots$$

consider a series $\sum_{n=1}^{\infty} (-1)^{n+1} a_n$

$$\delta_{2n+1} = a_1 - a_2 + a_3 - a_4 + \dots + a_{2n-1}$$

$$= \left(1 + \frac{1}{2} + \dots + \frac{1}{n} \right) - \left(\int_0^1 \frac{t^2}{x} dt \right)$$

$$(a_1 + a_3 + \dots + a_{2n-1}) - (a_2 + a_4 + \dots + a_{2n})$$

$$a_1 + a_3 + \dots + a_{2n-1} = \int_0^1 \frac{t^0}{x} dt + \int_0^1 \frac{t^2}{x} dt + \dots + \int_0^1 \frac{t^{2n-2}}{x} dt$$

$$= \int_0^1 \frac{t^{2n-2}}{x} dt$$

$$\delta_{2n+1} = 1 + \frac{1}{2} + \dots + \frac{1}{n} - \log n$$

$$a_n \rightarrow 0 \quad \text{as } n \rightarrow \infty \quad \text{and } \{a_n\} \text{ is a decreasing sequence}$$

$$a_{2n+1} > a_{2n} > a_{2n+1}$$

$$\frac{1}{n} > \frac{1}{n+1} > \frac{1}{n+1}$$

Leibniz rule applies: $\sum_{n=1}^{\infty} (-1)^{n+1} a_n$ converges.

$$\sum_{n=1}^{\infty} (-1)^{n+1} a_n = c \quad \text{hence } \delta_{2n+1} \rightarrow c \quad \text{as } n \rightarrow \infty$$

$$\lim_{n \rightarrow \infty} (1 + \frac{1}{2} + \dots + \frac{1}{n} - \log n) = c = \text{Euler's constant}$$

$$c = 0.5772156649$$

2.1 $\sum_{n=1}^{\infty} a_n$ converges $\Leftrightarrow \forall \varepsilon > 0 \exists N \in \mathbb{N}$ s.t. $\forall n \geq N, |a_n| < \varepsilon$

Then assume $\sum_{n=1}^{\infty} |a_n|$ converges. a_n are complex (a.s. real).
 Then $\sum_{n=1}^{\infty} a_n$ converges and $|\sum_{n=1}^{\infty} a_n| \leq \sum_{n=1}^{\infty} |a_n|$

Proofs
 (1) $\sum_{n=1}^{\infty} |a_n|$ conv \Rightarrow given any $\varepsilon > 0 \exists N \in \mathbb{N}$ s.t. $\forall n \geq N, |a_n| < \varepsilon$

$|a_{n+1}| + |a_{n+2}| + \dots + |a_m| < \varepsilon$

by triangle inequality
 $\Rightarrow \sum_{n=1}^{\infty} a_n$ conv

(2) case (1)': Assume $a_n \in \mathbb{R}$ & $n!$
 define $b_n = a_n + |a_n| = \begin{cases} 2|a_n| & \text{if } a_n \geq 0 \\ 0 & \text{if } a_n < 0 \end{cases}$

$\Rightarrow 0 \leq b_n \leq 2|a_n| \Rightarrow \sum b_n$ conv since $\sum |a_n|$ conv

$a_n = b_n - |a_n|$
 $\sum a_n$ conv $\Leftrightarrow \sum b_n - \sum |a_n|$

case (1)': general case $a_n = u_n + i v_n$

$|u_n|, |v_n| \leq |a_n| \Rightarrow \sum |a_n|$ conv $\Rightarrow \sum |u_n|$ conv
 $\Rightarrow \sum |v_n|$ conv

$\Rightarrow \sum u_n$ conv, $\sum v_n$ conv

$\Rightarrow \sum (u_n + i v_n) = \sum a_n$ conv

$\sum_{n=1}^{\infty} a_n$ (series of complex (a.s. real) no's) is absolutely convergent if $\sum_{n=1}^{\infty} |a_n|$ conv (Note: abs conv \Rightarrow conv)

conv \Rightarrow abs conv $\frac{a_n}{n} = a_n$
 $\frac{a_n}{n} = a_n$ is conv

conv but not abs conv = conditionally convergence

simple fact: $\sum_{n=1}^{\infty} a_n$ & $\sum_{n=1}^{\infty} b_n$ are abs conv $\Rightarrow \sum_{n=1}^{\infty} (a_n + b_n)$ conv

Proofs
 $|\sum_{k=1}^n a_k| \leq \sum_{k=1}^n |a_k|$ by triangle inequality

$|\sum_{k=1}^n a_k| \leq \sum_{k=1}^n |a_k|$

Now let $n \rightarrow \infty$ (n is then on the right)

$|\sum_{k=1}^n a_k| \leq \sum_{k=1}^n |a_k|$
 $b_n \rightarrow b$ $\Rightarrow \sum_{k=1}^n a_k \rightarrow \sum_{k=1}^{\infty} a_k$

$-c \leq b_n \leq c \quad \forall n \in \mathbb{N}$

$\rightarrow -c \leq \lim b_n = b \leq c$

$|b| \leq c$

Rearrangement of $\sum a_n$

$f: \mathbb{N} \rightarrow \mathbb{N}$ bijection $b_n = a_{f(n)}$

If $\sum a_n$ conv does $\sum b_n$ conv?

Example: $1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \frac{1}{6} + \frac{1}{7} - \frac{1}{8} + \frac{1}{9} - \frac{1}{10} + \dots$ $\sum a_n$

$1 - \frac{1}{2} - \frac{1}{2} + \frac{1}{3} + \frac{1}{3} - \frac{1}{4} + \frac{1}{4} + \frac{1}{4} - \frac{1}{5} + \frac{1}{5} + \frac{1}{5} - \frac{1}{6} + \frac{1}{6} + \frac{1}{6} - \frac{1}{7} + \frac{1}{7} + \frac{1}{7} - \frac{1}{8} + \frac{1}{8} + \frac{1}{8} - \frac{1}{9} + \frac{1}{9} + \frac{1}{9} - \frac{1}{10} + \frac{1}{10} + \frac{1}{10} - \dots$ $\sum b_n$

$$\lim_{n \rightarrow \infty} \frac{2n}{n+1} = 1 - 1 + 1 = 1$$

$$\left(\frac{1}{n+1} + \frac{1}{n+1} - \frac{1}{2n} \right) > 0$$

$$\frac{1}{n+1} > \frac{1}{2n}$$

Thus, $\sum_{n=1}^{\infty} \frac{1}{n^2}$ converges also to any rearrangement of $\sum_{n=1}^{\infty} \frac{1}{n^2}$

Thus, let $\sum_{n=1}^{\infty} a_n$ be a series of real numbers which converges conditionally.

$$\text{let } -a_n \leq a_n \leq a_n$$

Then \exists a rearrangement of $\sum_{n=1}^{\infty} a_n$ such that

$$\limsup_{n \rightarrow \infty} b_n = \infty \text{ and } \liminf_{n \rightarrow \infty} b_n = -\infty$$

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} = 0 \text{ for } \epsilon = 0$$

Proof: Given $\epsilon > 0$ to find $N \in \mathbb{N}$ such that $\frac{1}{n^2} < \epsilon$ for $n \geq N$

$$\frac{1}{n^2} < \epsilon \Leftrightarrow n > \frac{1}{\sqrt{\epsilon}}$$

Given $\epsilon > 0$ choose $N \in \mathbb{N}$ such that $n \geq N \Rightarrow \frac{1}{n^2} < \epsilon$

$$\text{S.t. } N > \frac{1}{\sqrt{\epsilon}}$$

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} = 0 \text{ for } \epsilon = 0$$

Proof: Given $\epsilon > 0$ to find $N \in \mathbb{N}$ s.t. $n \geq N \Rightarrow |\frac{1}{n^2} - 0| < \epsilon$

case (D): Assume $p > 1 \Rightarrow \frac{1}{n^p} > 1$ for $n \in \mathbb{N}$, define $a_n = \frac{1}{n^p} - 1 > 0$

$$\Rightarrow (1 + a_n)^n > \frac{1}{n^p} \Rightarrow (1 + a_n)^n = p$$

$$\text{N.B., } 1 + a_n \leq (1 + a_n)^n = p$$

$$\Rightarrow 0 < a_n \leq \frac{p-1}{n} \Rightarrow \lim_{n \rightarrow \infty} a_n = 0 \therefore 0 \leq \lim_{n \rightarrow \infty} a_n \leq \lim_{n \rightarrow \infty} \frac{p-1}{n} = 0$$

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} = 0 \text{ for } \epsilon = 0 \Rightarrow \lim_{n \rightarrow \infty} \frac{1}{n^2} = 0$$

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} = 1 \text{ (by above case, since } a_n > 0 \text{)}$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{1}{n^2} = 1$$

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} = 1$$

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} = 1$$

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} = 1$$

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} = 1$$

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} = 1$$

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} = 1$$

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} = 1$$

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} = 1$$

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} = 1$$

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} = 1$$

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} = 1$$

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} = 1$$

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} = 1$$

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} = 1$$

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} = 1$$

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} = 1$$

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} = 1$$

⑤ $|n| < 1$ then $\lim_{n \rightarrow \infty} x^n = 0$

Proof: choose $\epsilon > 0$ in (2)

$$0 = \lim_{n \rightarrow \infty} \frac{n!}{n!} = \lim_{n \rightarrow \infty} \frac{1}{n!} \quad (p=0)$$

$$\lim_{n \rightarrow \infty} \frac{1}{n!} \rightarrow 0 \text{ as } n \rightarrow \infty$$

⑥ $e = \sum_{n=0}^{\infty} \frac{1}{n!} = 1 + \frac{1}{1!} + \frac{1}{2!} + \dots$

Ratio test: $\frac{a_{n+1}}{a_n} = \frac{n!}{(n+1)!} = \frac{1}{n+1} \rightarrow 0$ as $n \rightarrow \infty$

$\Rightarrow e$ converges

So, $e = \sum_{n=0}^{\infty} \frac{1}{n!}$ makes sense

e is irrational but algebraic h_0

rational \Rightarrow algebraic

$\alpha = h_0 \Rightarrow q\alpha - p = 0$

non-algebraic h_0 is transcendental h_0

⑦ $\lim_{n \rightarrow \infty} (1 + \frac{1}{n})^n = e = \sum_{n=0}^{\infty} \frac{1}{n!}$

Proof: $s_n = \sum_{k=0}^n \frac{1}{k!}$

$t_n = (1 + \frac{1}{n})^n$

$= (1 + \frac{1}{n})^n = \sum_{k=0}^n \binom{n}{k} (\frac{1}{n})^k$

$\lim_{n \rightarrow \infty} t_n = \lim_{n \rightarrow \infty} \sum_{k=0}^n \binom{n}{k} (\frac{1}{n})^k$

$= \sum_{k=0}^{\infty} \lim_{n \rightarrow \infty} \binom{n}{k} (\frac{1}{n})^k$

$= \sum_{k=0}^{\infty} \frac{1}{k!}$

$< s_n \Rightarrow t_n < s_n \Rightarrow \lim_{n \rightarrow \infty} \sup t_n \leq \lim_{n \rightarrow \infty} \sup s_n = \lim_{n \rightarrow \infty} s_n = e$

def: n th partial sum

$s_n = 1 + \frac{1}{2!} + \frac{1}{3!} + \dots + \frac{1}{n!} = \sum_{k=0}^n \frac{1}{k!}$

def: n th partial sum, let $n \rightarrow \infty$

$\lim_{n \rightarrow \infty} s_n = e$

$\lim_{n \rightarrow \infty} \inf t_n = \lim_{n \rightarrow \infty} \sup t_n = \lim_{n \rightarrow \infty} s_n = e$

So, $e = \lim_{n \rightarrow \infty} (1 + \frac{1}{n})^n$

$s_n = 1 + \frac{1}{2!} + \dots + \frac{1}{n!}$

$e = \sum_{k=0}^{\infty} \frac{1}{k!}$

$e - s_n = \sum_{k=n+1}^{\infty} \frac{1}{k!}$

$= \frac{1}{(n+1)!} \left[1 + \frac{1}{n+2} + \frac{1}{(n+3)} + \dots \right]$

$< \frac{1}{(n+1)!} \left[1 + \frac{1}{n+1} + \frac{1}{(n+1)^2} + \dots \right]$

$= \frac{1}{(n+1)!} \cdot \frac{1}{1 - \frac{1}{n+1}} = \frac{1}{n!}$

$e - s_n < \frac{1}{n!}$

$n = 10 \Rightarrow s_{10} = 1 + \frac{1}{2!} + \frac{1}{3!} + \dots + \frac{1}{10!}$

$e - s_{10} < \frac{1}{10!} < \frac{1}{10^6}$ check

Thm 2: is irrational

Proof: Suppose π is rational, let $\pi = \frac{p}{q}$

$$p, q \in \mathbb{Z}, q \neq 0$$

$$0 < (x - \frac{p}{q}) < \frac{1}{q} \Rightarrow 0 < q(x - \frac{p}{q}) < 1$$

$$q(x - \frac{p}{q})$$

integer

$$q_1, q_2 = q_1(1 + \frac{1}{q_1} + \frac{1}{q_1^2} + \dots + \frac{1}{q_1^{n-1}})$$

q is also an integer

$$q_1 = q_1 \cdot \frac{p}{q} = (a_1) \cdot \frac{p}{q}$$

what we get is $q_1 - q_1 \cdot \frac{p}{q}$ is integer

by contradiction of these is no integer b/w

$$0 \leq 1/q$$

Q3: Does \exists a bijection from \mathbb{N} to \mathbb{Q} which is order preserving? i.e. $f: \mathbb{N} \rightarrow \mathbb{Q}$ $f(1) < f(2) < f(3) < \dots$

Sol: acc to dense prop, there is no such function

a bijection

Q4: (Axioms) Rudin, Spake, Munkres (calculus) (theorems)

lim inf & lim sup are not 0, this cause

f: $\mathbb{R} \rightarrow \mathbb{R}$ and consider a sequence of function $f_n(x) = \frac{1}{n}$ $f_n: \mathbb{R} \rightarrow \mathbb{R}$ $f_n(x) = \frac{1}{n}$ $f_n \rightarrow 0$ as $n \rightarrow \infty$

if $f_n \rightarrow f$ then $f_n(x) \rightarrow f(x)$ for all $x \in \mathbb{R}$

if $f_n \rightarrow f$ then $f_n(x) \rightarrow f(x)$ for all $x \in \mathbb{R}$

if $f_n \rightarrow f$ then $f_n(x) \rightarrow f(x)$ for all $x \in \mathbb{R}$

if $f_n \rightarrow f$ then $f_n(x) \rightarrow f(x)$ for all $x \in \mathbb{R}$

if $f_n \rightarrow f$ then $f_n(x) \rightarrow f(x)$ for all $x \in \mathbb{R}$

if $f_n \rightarrow f$ then $f_n(x) \rightarrow f(x)$ for all $x \in \mathbb{R}$

if $f_n \rightarrow f$ then $f_n(x) \rightarrow f(x)$ for all $x \in \mathbb{R}$

if $f_n \rightarrow f$ then $f_n(x) \rightarrow f(x)$ for all $x \in \mathbb{R}$

if $f_n \rightarrow f$ then $f_n(x) \rightarrow f(x)$ for all $x \in \mathbb{R}$

if $f_n \rightarrow f$ then $f_n(x) \rightarrow f(x)$ for all $x \in \mathbb{R}$

if $f_n \rightarrow f$ then $f_n(x) \rightarrow f(x)$ for all $x \in \mathbb{R}$

if $f_n \rightarrow f$ then $f_n(x) \rightarrow f(x)$ for all $x \in \mathbb{R}$

if $f_n \rightarrow f$ then $f_n(x) \rightarrow f(x)$ for all $x \in \mathbb{R}$

if $f_n \rightarrow f$ then $f_n(x) \rightarrow f(x)$ for all $x \in \mathbb{R}$

if $f_n \rightarrow f$ then $f_n(x) \rightarrow f(x)$ for all $x \in \mathbb{R}$

if $f_n \rightarrow f$ then $f_n(x) \rightarrow f(x)$ for all $x \in \mathbb{R}$

if $f_n \rightarrow f$ then $f_n(x) \rightarrow f(x)$ for all $x \in \mathbb{R}$

if $f_n \rightarrow f$ then $f_n(x) \rightarrow f(x)$ for all $x \in \mathbb{R}$

if $f_n \rightarrow f$ then $f_n(x) \rightarrow f(x)$ for all $x \in \mathbb{R}$

if $f_n \rightarrow f$ then $f_n(x) \rightarrow f(x)$ for all $x \in \mathbb{R}$

if $f_n \rightarrow f$ then $f_n(x) \rightarrow f(x)$ for all $x \in \mathbb{R}$

if $f_n \rightarrow f$ then $f_n(x) \rightarrow f(x)$ for all $x \in \mathbb{R}$

if $f_n \rightarrow f$ then $f_n(x) \rightarrow f(x)$ for all $x \in \mathbb{R}$

if $f_n \rightarrow f$ then $f_n(x) \rightarrow f(x)$ for all $x \in \mathbb{R}$

$$\lim_{n \rightarrow \infty} \int_a^b f_n(x) dx = \lim_{n \rightarrow \infty} \frac{1}{2} = \frac{1}{2}$$

$$\text{but } \int_a^b \lim_{n \rightarrow \infty} f_n(x) dx = \int_a^b f(x) dx = 0$$

pointwise conv:

$$f_n : E \rightarrow \mathbb{R} \text{ (or } \mathbb{C}) \quad f : E \rightarrow \mathbb{R} \text{ (or } \mathbb{C})$$

suppose $f_n \rightarrow f$ pointwise on E

let $x \in E$ then $\lim_{n \rightarrow \infty} f_n(x) = f(x)$ means for every $\epsilon > 0$
 $\exists N \in \mathbb{N}$ s.t. $\forall n \geq N$ we have $|f_n(x) - f(x)| < \epsilon$
 $N \in \mathbb{N}$

Defn: let $f_n : E \rightarrow \mathbb{R}$ (or \mathbb{C}) be a sequence of functions such that f_n converges to f pointwise i.e. $\lim_{n \rightarrow \infty} f_n(x) = f(x)$ with $x \in E$. f_n 's said to f uniformly on E if given $\epsilon > 0$ $\exists N \in \mathbb{N}$ (N depends only on ϵ but not index of $x \in E$)
 $\text{s.t. } \forall n \geq N \quad |f_n(x) - f(x)| < \epsilon \quad \forall x \in E$

Notation: $f_n \rightarrow f$ uniformly on E



We have $|f(x) - f_n(x)| < \epsilon$ for all $x \in E$

$$|f_n(x) - f(x)| < \epsilon$$

Thm: Suppose $f_n \rightarrow f$ uniformly on E, suppose f_n is

cont. $\forall n \in \mathbb{N}$
 $\Rightarrow f$ is cont. on E .

Proof: "Unit limit of cont. fns is cont."

Let f_n be a sequence of cont. fns on E. Suppose $f_n \rightarrow f$ uniformly on E. We want to show f is cont. on E.

Given $\epsilon > 0$ we have to find a $\delta > 0$ s.t. $|x - y| < \delta \Rightarrow |f(x) - f(y)| < \epsilon$

Given this $\epsilon > 0$, $\exists N \in \mathbb{N}$ s.t. $\forall n \geq N$ $\forall x \in E$ have

$$|f_n(x) - f(x)| < \epsilon/3$$

f_n is cont. hence $\exists \delta > 0$ s.t. $|x - y| < \delta \Rightarrow |f_n(x) - f_n(y)| < \epsilon/3$

$$|f(x) - f(y)| \leq |f(x) - f_n(x)| + |f_n(x) - f_n(y)| + |f_n(y) - f(y)|$$

$$\leq \epsilon/3 + \epsilon/3 + \epsilon/3 = \epsilon$$

$$|f(x) - f(y)| < \epsilon$$

$\Rightarrow f$ is cont. at $x \Rightarrow f$ is cont. on E