

Name : Tanjim Reza

Student ID : 20101065

Course : CSE330


Section : 10

Assignment no: 06

Ans: to the que no: A

1

Sketch from Desmos



From the values of the graph, the first positive value of intersection point is $(4.493, 4.493)$.

The positive value of x is in between $[4, 5]$ for the graph $y=x$, $y=\tan x$.

2

Given interval is $[4, 5]$. and the root for $f(x) = x - \tan x$ is in this interval.

Now, Using Bisection method:

Iteration 01:

$$\begin{aligned}c &= \frac{a+b}{2} \\&= \frac{4+5}{2} \\&= 4.5\end{aligned}$$

$$f(4.5) = -0.1373 < 0$$

New interval $[4, 4.5]$

Iteration 2:

$$\begin{aligned}c &= \frac{4+4.5}{2} \\&= 4.25\end{aligned}$$

$$f(4.25) = 2.2437 > 0$$

$\therefore [4.25, 4.5]$

Iteration 03:

$$c = \frac{4.25 + 4.5}{2}$$

$$= 4.375$$

$$f(4.375) = 1.5244 > 0$$

$$\text{New, interval} = [4.375, 4.5]$$

Iteration 4:

$$c = \frac{4.375 + 4.5}{2}$$

$$= 4.4375$$

$$f(4.4375) = 0.8918 > 0$$

$$\text{New interval} = [4.4375, 4.5]$$

Iteration 5:

$$c = \frac{4.4375 + 4.5}{2} = 4.46875$$

$$f(4.46875) = 0.4459 > 0$$

$$\text{New interval} = [4.46875, 4.5]$$

Iteration 06:

$$c = \frac{4.46875 + 4.5}{2} = 4.484375$$

$$P(4.484375) = 0.1749 > 0$$

$$\text{Interval} = [4.484375, 4.5]$$

Iteration 07:

$$c = \frac{4.484375 + 4.5}{2}$$

$$= 4.4921875$$

$$P(4.4921875) = 0.024570$$

$$\text{Interval now} = [4.4921875, 4.5]$$

Iteration 08:

$$c = \frac{4.4921875 + 4.5}{2}$$

$$= 4.49609375$$

$$P(4.49609375) = -0.0599 < 0$$

$$\therefore \text{Interval} = [4.4921875, 4.49609375]$$

Iteration 9

$$c = 4.494140625$$

$$f(4.494140625) = -0.0148 < 0$$

$$\therefore \text{Interval } [4.4921875, 4.494140625]$$

Iteration 10

$$c = \frac{a+b}{2}$$

$$= 4.493164063$$

\therefore Approximate first positive value = 4.493164063

Ans: to the que no: B

(1)

$$f(x) = x - \tan x$$

$$x \in I = [4, 5]$$

$$f(x) = 0$$

$$\therefore x - \tan x = 0$$

Now

$$g(x) = \tan(x)$$

$$\Rightarrow g(x) = \tan^{-1}(x)$$

$$g(x) = \sin^{-1}(x \cdot \cos(x))$$

Putting values 4, 5, 4.5 the solution can not reach to the solution. $\text{Ans} \notin I$.

$$\therefore g(x) = \pi + \tan^{-1}(x), \text{ over } [4, 5]$$

$$\therefore g(x) = \pi + \tan^{-1}(x)$$

2

Letting, $x_0 = 4.5$

$$\begin{aligned}x_1 &= g(4.5) \\ &= 4.49372\end{aligned}$$

$$x_2 = g(4.49372) = 4.49342$$

$$x_3 = g(4.49342) = 4.4931.$$