



Definite Integral

1.

Definite Integral As The Limit Of A Sum

$$\int_a^b f(x)dx = \lim_{h \rightarrow 0} [f(a) + f(a+h) + \dots + f(a+(n-1)h)]$$

$$h = \frac{b-a}{n} \rightarrow 0 \text{ as } n \rightarrow \infty$$

The above expression is known as the definite integral as the limit of a sum.

2.

Properties Of Definite Integrals

$$(i) \int_a^b f(x)dx = \int_a^b f(t)dt$$

$$(ii) \int_a^b f(x)dx = -\int_b^a f(x)dx \text{ in particular } \int_a^a f(x)dx = 0$$

$$(iii) \int_a^b f(x)dx = \int_a^c f(x)dx + \int_c^b f(x)dx \text{ where } a < c < b$$

$$(iv) \int_a^b f(x)dx = \int_a^b f(a+b-x)dx$$

$$(v) \int_0^a f(x)dx = \int_0^a f(a-x)dx$$

$$(vi) \int_{-a}^a f(x)dx = 2 \int_0^a f(x)dx \text{ if } f(x) \text{ is an even function}$$

$$(vii) \int_{-a}^a f(x)dx = 0, \text{ if } f(x) \text{ is an odd function}$$

$$(viii) \int_0^{2a} f(x)dx = \int_0^a f(x)dx + \int_0^a f(2a-x)dx$$

$$(ix) \int_0^{2a} f(x)dx = 2 \int_0^a f(x)dx \text{ if } f(2a-x) = f(x) = 0, \text{ if } f(2a-x) = -f(x)$$

3.

Walli's formula

$$\int_0^{\pi/2} \sin^n x dx = \int_0^{\pi/2} \cos^n x dx = \begin{cases} \frac{n-1}{n} \cdot \frac{n-3}{n-2} \cdot \frac{n-5}{n-4} \dots \frac{2}{3}, & \text{when } n \text{ is odd} \\ \frac{n-1}{n} \cdot \frac{n-3}{n-2} \cdot \frac{n-5}{n-4} \dots \frac{3}{4} \cdot \frac{1}{2} \cdot \frac{\pi}{2}, & \text{when } n \text{ is even} \end{cases}$$

$$\int_0^{\pi/2} \sin^m x \cos^n x dx = \frac{(m-1)(m-3)\dots(n-1)(n-3)\dots}{(m+n)(m+n-2)\dots(2 \text{ or } 1)}$$

[If m, n are both odd positive integers or one odd positive integer]

$$\int_0^{\pi/2} \sin^m x \cos^n x dx = \frac{(m-1)(m-3)\dots(n-1)(n-3)\dots}{(m+n)(m+n-2)\dots(2 \text{ or } 1)} \cdot \frac{\pi}{2}$$

[If m, n are both even positive integers]

4.

Periodic Properties

If f(x) is a periodic function with period T, then

$$01 \int_0^{nT} f(x)dx = n \int_0^T f(x)dx, n \in \mathbb{Z}$$

$$02 \int_a^{a+nT} f(x)dx = n \int_0^T f(x)dx, n \in \mathbb{Z}, a \in \mathbb{R}$$

$$03 \int_{mT}^{nT} f(x)dx = (n-m) \int_0^T f(x)dx, m, n \in \mathbb{Z}$$

$$04 \int_{nT}^{a+nT} f(x)dx = \int_0^a f(x)dx, n \in \mathbb{Z}, a \in \mathbb{R}$$

5.

Advance properties

$$\psi(x) \leq f(x) \leq \phi(x) \text{ for } a \leq x \leq b \text{ then } \int_a^b \psi(x)dx \leq \int_a^b f(x)dx \leq \int_a^b \phi(x)dx$$

$$\text{If } m \leq f(x) \leq M \text{ for } a \leq x \leq b, \text{ then } m(b-a) \leq \int_a^b f(x)dx \leq M(b-a)$$

01

$$\left| \int_a^b f(x)dx \right| \leq \int_a^b |f(x)|dx$$

02

$$\text{If } f(x) \geq 0 \text{ on } [a, b] \text{ then}$$

$$\int_a^b f(x)dx \geq 0$$

03

6.

Leibnitz Theorem

$$\text{if } F(x) = \int_{g(x)}^{h(x)} f(t)dt, \text{ then } \frac{dF(x)}{dx} = h'(x)f(h(x)) - g'(x)f(g(x))$$

Gamma function

$$\Gamma n = \int_0^\infty e^{-x} x^{n-1} dx$$

where n is a positive rational number

7.

Beta & Gama Function

Properties of gamma function

$$1) \Gamma(0) = \infty, \Gamma(1) = 1$$

$$2) \Gamma(n+1) = n\Gamma(n)$$

$$3) \Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$$

$$4) \Gamma(n)\Gamma(1-n) = \frac{\pi}{\sin n\pi}, 0 < n < 1$$

Beta function

$$\beta(m, n) = \int_0^1 x^{m-1} (1-x)^{n-1} dx$$

NOTE

The relationship between beta & gamma function will be

$$\beta(m, n) = \frac{\Gamma m \Gamma n}{\Gamma(m+n)}$$

**8.****Important results**

$$\sum_{r=1}^n r = \frac{n(n+1)}{2}$$

$$\sum_{r=1}^n r^2 = \frac{1}{6}n(n+1)(2n+1)$$

$$\sum_{n=1}^n r^3 = \frac{n^2(n+1)^2}{4}$$

In GP, sum of n terms, $S = S_n = \begin{cases} \frac{a(r^n - 1)}{r - 1}, & |r| > 1 \\ an, & r = 1 \\ \frac{a(1 - r^n)}{1 - r}, & |r| < 1 \end{cases}$

$$\cosh \theta = \frac{e^\theta + e^{-\theta}}{2}, \sinh \theta = \frac{e^\theta - e^{-\theta}}{2}$$

$$\sin \alpha + \sin(\alpha + \beta) + \sin(\alpha + 2\beta) + \dots + \sin(\alpha + (n-1)\beta) = \frac{\sin(n\beta/2)}{\sin(\beta/2)} \sin(\alpha + (n-1)\beta/2).$$

$$\cos \alpha + \cos(\alpha + \beta) + \dots + \cos(\alpha + (n-1)\beta) = \frac{\sin n\beta/2}{\sin \beta/2} \cdot \cos(\alpha + (n-1)\beta/2)$$

$$1 - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \frac{1}{5^2} - \frac{1}{6^2} + \dots \infty = \frac{\pi^2}{12}$$

$$1 + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots \infty = \frac{\pi^2}{6}$$

$$\frac{1}{2^2} + \frac{1}{4^2} + \frac{1}{6^2} + \dots \infty = \frac{\pi^2}{24}$$

$$\cos \theta = \frac{e^{i\theta} + e^{-i\theta}}{2}, \sin \theta = \frac{e^{i\theta} - e^{-i\theta}}{2i}$$

$$1/1^2 + 1/3^2 + 1/5^2 + 1/7^2 + \dots \infty = \pi^2 / 8$$

9**Average Value Theorem**

If f is a continuous function on $[a, b]$, then its average value on $[a, b]$ is given by the formula

$$f_{\text{AVG}[a,b]} = \frac{1}{b-a} \cdot \int_a^b f(x) dx$$

Application Of Integrals

1

$$\int_a^b f(x) dx \neq \text{Area under the curve } f(x) \text{ from } a \text{ to } b$$

$$\int_a^b f(x) dx = \text{Algebraic area under the curve } f(x) \text{ from } a \text{ to } b$$



2**POSITIVE AND NEGATIVE AREA**

Area is always taken as positive. If some part of the area lies in the positive side i.e., above x-axis and some part lies in the negative side i.e. below x-axis then the area of two parts should be calculated separately and then add their numerical values to get the desired area.



3

Area Under Simple Curves

(i) Area of the region bounded by a curve $y = f(x)$ and x -axis between the two ordinates




(ii) Area of the region bounded by a curve $x = f(y)$ and x -axis between the two ordinates



4


Area Under Different Curves

CASE-I




$$A = \int_a^c f(x) dx + \int_c^b g(x) dx$$

CASE-II




$$A = \int [g(x) - f(x)] dx$$

CASE-III




$$A = \int_a^b [g(x) - f(x)] dx$$

CASE-IV



$$A = \int_a^c [f(x) - g(x)] dx + \int_c^b [g(x) - f(x)] dx$$

CASE-V



$$A = \int_a^c (y_1 - y_2) dx + \int_c^d (y_2 - y_1) dx + \int_d^b (y_1 - y_2) dx$$