

# Discrete Mathematical Models

## Lecture 14

---

Kane Townsend

Semester 2, 2024

# Vectors

---

# Vectors and vector arithmetic

For any  $n \in \mathbb{N}$  an element  $x = (x_1, \dots, x_n) \in \mathbb{Q}^n$  will be called a **vector**.

# Vectors and vector arithmetic

For any  $n \in \mathbb{N}$  an element  $x = (x_1, \dots, x_n) \in \mathbb{Q}^n$  will be called a **vector**.

A vector  $x \in \mathbb{Q}^n$  can be viewed as  
an element of  $M_{1 \times n}(\mathbb{Q})$ ;  $x$  is then called a **row vector**

# Vectors and vector arithmetic

For any  $n \in \mathbb{N}$  an element  $x = (x_1, \dots, x_n) \in \mathbb{Q}^n$  will be called a **vector**.

A vector  $x \in \mathbb{Q}^n$  can be viewed as

an element of  $M_{1 \times n}(\mathbb{Q})$ ;  $x$  is then called a **row vector**

or as an element of  $M_{n \times 1}(\mathbb{Q})$ ;  $x$  is then called a **column vector**.

# Vectors and vector arithmetic

For any  $n \in \mathbb{N}$  an element  $x = (x_1, \dots, x_n) \in \mathbb{Q}^n$  will be called a **vector**.

A vector  $x \in \mathbb{Q}^n$  can be viewed as

an element of  $M_{1 \times n}(\mathbb{Q})$ ;  $x$  is then called a **row vector**

or as an element of  $M_{n \times 1}(\mathbb{Q})$ ;  $x$  is then called a **column vector**.

The **sum** of two vectors,  $x + y$ , is defined element-wise:

# Vectors and vector arithmetic

For any  $n \in \mathbb{N}$  an element  $x = (x_1, \dots, x_n) \in \mathbb{Q}^n$  will be called a **vector**.

A vector  $x \in \mathbb{Q}^n$  can be viewed as

an element of  $M_{1 \times n}(\mathbb{Q})$ ;  $x$  is then called a **row vector**

or as an element of  $M_{n \times 1}(\mathbb{Q})$ ;  $x$  is then called a **column vector**.

The **sum** of two vectors,  $x + y$ , is defined element-wise:

$$x + y = (x_1, \dots, x_n) + (y_1, \dots, y_n) = (x_1 + y_1, \dots, x_n + y_n).$$

# Vectors and vector arithmetic

For any  $n \in \mathbb{N}$  an element  $x = (x_1, \dots, x_n) \in \mathbb{Q}^n$  will be called a **vector**.

A vector  $x \in \mathbb{Q}^n$  can be viewed as

an element of  $M_{1 \times n}(\mathbb{Q})$ ;  $x$  is then called a **row vector**

or as an element of  $M_{n \times 1}(\mathbb{Q})$ ;  $x$  is then called a **column vector**.

The **sum** of two vectors,  $x + y$ , is defined element-wise:

$$x + y = (x_1, \dots, x_n) + (y_1, \dots, y_n) = (x_1 + y_1, \dots, x_n + y_n).$$

(When viewed as row or column vectors,  $x$  and  $y$  must be the same shape.)



# Vectors and vector arithmetic

For any  $n \in \mathbb{N}$  an element  $x = (x_1, \dots, x_n) \in \mathbb{Q}^n$  will be called a **vector**.

A vector  $x \in \mathbb{Q}^n$  can be viewed as

an element of  $M_{1 \times n}(\mathbb{Q})$ ;  $x$  is then called a **row vector**

or as an element of  $M_{n \times 1}(\mathbb{Q})$ ;  $x$  is then called a **column vector**.

The **sum** of two vectors,  $x + y$ , is defined element-wise:

$$x + y = (x_1, \dots, x_n) + (y_1, \dots, y_n) = (x_1 + y_1, \dots, x_n + y_n).$$

(When viewed as row or column vectors,  $x$  and  $y$  must be the same shape.)

There are a number of ways to define the product of two vectors (e.g the 'inner' and the 'outer' products) but we will not use them in this course.

# Vectors and vector arithmetic

For any  $n \in \mathbb{N}$  an element  $x = (x_1, \dots, x_n) \in \mathbb{Q}^n$  will be called a **vector**.

A vector  $x \in \mathbb{Q}^n$  can be viewed as

an element of  $M_{1 \times n}(\mathbb{Q})$ ;  $x$  is then called a **row vector**

or as an element of  $M_{n \times 1}(\mathbb{Q})$ ;  $x$  is then called a **column vector**.

The **sum** of two vectors,  $x + y$ , is defined element-wise:

$$x + y = (x_1, \dots, x_n) + (y_1, \dots, y_n) = (x_1 + y_1, \dots, x_n + y_n).$$

(When viewed as row or column vectors,  $x$  and  $y$  must be the same shape.)

There are a number of ways to define the product of two vectors (e.g the 'inner' and the 'outer' products) but we will not use them in this course.

However we do need to define the product of a number  $\lambda$  and a vector. In this context the number  $\lambda$  is referred to as a **scalar**, to distinguish it from a vector, and the product  $\lambda x$  is called a **scalar product**.

# Vectors and vector arithmetic

For any  $n \in \mathbb{N}$  an element  $x = (x_1, \dots, x_n) \in \mathbb{Q}^n$  will be called a **vector**.

A vector  $x \in \mathbb{Q}^n$  can be viewed as

an element of  $M_{1 \times n}(\mathbb{Q})$ ;  $x$  is then called a **row vector**

or as an element of  $M_{n \times 1}(\mathbb{Q})$ ;  $x$  is then called a **column vector**.

The **sum** of two vectors,  $x + y$ , is defined element-wise:

$$x + y = (x_1, \dots, x_n) + (y_1, \dots, y_n) = (x_1 + y_1, \dots, x_n + y_n).$$

(When viewed as row or column vectors,  $x$  and  $y$  must be the same shape.)

There are a number of ways to define the product of two vectors (e.g the 'inner' and the 'outer' products) but we will not use them in this course.

However we do need to define the product of a number  $\lambda$  and a vector. In this context the number  $\lambda$  is referred to as a **scalar**, to distinguish it from a vector, and the product  $\lambda x$  is called a **scalar product**. It is also defined element-wise:

$$\forall \lambda \in \mathbb{Q} \quad \lambda x = \lambda(x_1, \dots, x_n) = (\lambda x_1, \dots, \lambda x_n).$$

## Examples of vectors and vector arithmetic

- Let  $p = (p_1, p_2, p_3) \in \mathbb{Q}^3$  represent the state of an ecosystem with  $p_1, p_2, p_3$  being the sizes of the populations of three different species.

## Examples of vectors and vector arithmetic

- Let  $p = (p_1, p_2, p_3) \in \mathbb{Q}^3$  represent the state of an ecosystem with  $p_1, p_2, p_3$  being the sizes of the populations of three different species. If  $p_1$  increases by 10 individuals,  $p_2$  loses 20 individuals, and  $p_3$  gains 2, then the new state of the ecosystem is

$$p + c = (p_1, p_2, p_3) + (10, -20, 2)$$

## Examples of vectors and vector arithmetic

- Let  $\mathbf{p} = (p_1, p_2, p_3) \in \mathbb{Q}^3$  represent the state of an ecosystem with  $p_1, p_2, p_3$  being the sizes of the populations of three different species. If  $p_1$  increases by 10 individuals,  $p_2$  loses 20 individuals, and  $p_3$  gains 2, then the new state of the ecosystem is

$$\mathbf{p} + \mathbf{c} = (p_1, p_2, p_3) + (10, -20, 2)$$

- Let  $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{Q}^n$  represent the amplitudes  $a_k$ ,  $1 \leq k \leq n$ , of the harmonic frequencies  $kf$  of the fundamental frequency  $f$  of a note played on a violin.

## Examples of vectors and vector arithmetic

- Let  $\mathbf{p} = (p_1, p_2, p_3) \in \mathbb{Q}^3$  represent the state of an ecosystem with  $p_1, p_2, p_3$  being the sizes of the populations of three different species. If  $p_1$  increases by 10 individuals,  $p_2$  loses 20 individuals, and  $p_3$  gains 2, then the new state of the ecosystem is

$$\mathbf{p} + \mathbf{c} = (p_1, p_2, p_3) + (10, -20, 2)$$

- Let  $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{Q}^n$  represent the amplitudes  $a_k$ ,  $1 \leq k \leq n$ , of the harmonic frequencies  $ka$  of the fundamental frequency  $f$  of a note played on a violin. Then

$$3\mathbf{a} = 3(a_1, \dots, a_n),$$

represents to the same sound, but three times stronger.

# Addition and scalar multiplication of matrices

The same can be done with matrices.



# Addition and scalar multiplication of matrices

The same can be done with matrices.

For matrices  $A = (a_{i,j})$  and  $B = (b_{i,j})$  in  $M_n(\mathbb{Q})$ , and  $\lambda \in \mathbb{Q}$ , we define the **sum**  $A + B$  and **scalar product**  $\lambda A$  by

$$\begin{aligned} A + B &= (a_{i,j}) + (b_{i,j}) = (a_{i,j} + b_{i,j}). \\ \lambda A &= \lambda(a_{i,j}) = (\lambda a_{i,j}). \end{aligned}$$

# Addition and scalar multiplication of matrices

The same can be done with matrices.

For matrices  $A = (a_{i,j})$  and  $B = (b_{i,j})$  in  $M_n(\mathbb{Q})$ , and  $\lambda \in \mathbb{Q}$ , we define the **sum**  $A + B$  and **scalar product**  $\lambda A$  by

$$\begin{aligned} A + B &= (a_{i,j}) + (b_{i,j}) = (a_{i,j} + b_{i,j}). \\ \lambda A &= \lambda(a_{i,j}) = (\lambda a_{i,j}). \end{aligned}$$

Examples:

$$\begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} + \begin{pmatrix} 5 & 6 \\ 7 & 8 \end{pmatrix} = \begin{pmatrix} 6 & 8 \\ 10 & 12 \end{pmatrix}$$

# Addition and scalar multiplication of matrices

The same can be done with matrices.

For matrices  $A = (a_{i,j})$  and  $B = (b_{i,j})$  in  $M_n(\mathbb{Q})$ , and  $\lambda \in \mathbb{Q}$ , we define the **sum**  $A + B$  and **scalar product**  $\lambda A$  by

$$\begin{aligned} A + B &= (a_{i,j}) + (b_{i,j}) = (a_{i,j} + b_{i,j}). \\ \lambda A &= \lambda(a_{i,j}) = (\lambda a_{i,j}). \end{aligned}$$

Examples:

$$\begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} + \begin{pmatrix} 5 & 6 \\ 7 & 8 \end{pmatrix} = \begin{pmatrix} 6 & 8 \\ 10 & 12 \end{pmatrix}$$

$$5 \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} = \begin{pmatrix} 5 & 10 \\ 15 & 20 \end{pmatrix}.$$

# Linear Functions

---

# Linear functions

Definition: A function  $F : \mathbb{Q}^n \rightarrow \mathbb{Q}^n$  is called **linear** if and only if it satisfies the following two conditions:

- $F(x + y) = F(x) + F(y) \quad \forall x, y \in \mathbb{Q}^n.$
- $F(\lambda x) = \lambda F(x) \quad \forall x \in \mathbb{Q}^n \quad \forall \lambda \in \mathbb{Q}.$

# Linear functions

Definition: A function  $F : \mathbb{Q}^n \rightarrow \mathbb{Q}^n$  is called **linear** if and only if it satisfies the following two conditions:

- $F(x + y) = F(x) + F(y) \quad \forall x, y \in \mathbb{Q}^n.$
- $F(\lambda x) = \lambda F(x) \quad \forall x \in \mathbb{Q}^n \quad \forall \lambda \in \mathbb{Q}.$

Example: For  $n \in \mathbb{N}$  suppose  $(a_1, \dots, a_n) \in \mathbb{Q}^n$  represents the amplitudes of the different harmonics of a note played on a violin.

# Linear functions

Definition: A function  $F : \mathbb{Q}^n \rightarrow \mathbb{Q}^n$  is called **linear** if and only if it satisfies the following two conditions:

- $F(x + y) = F(x) + F(y) \quad \forall x, y \in \mathbb{Q}^n.$
- $F(\lambda x) = \lambda F(x) \quad \forall x \in \mathbb{Q}^n \quad \forall \lambda \in \mathbb{Q}.$

Example: For  $n \in \mathbb{N}$  suppose  $(a_1, \dots, a_n) \in \mathbb{Q}^n$  represents the amplitudes of the different harmonics of a note played on a violin. Thus  $a_n$  is the amplitude of frequency  $nf$ , where  $f$  is the fundamental frequency of the note.

# Linear functions

Definition: A function  $F : \mathbb{Q}^n \rightarrow \mathbb{Q}^n$  is called **linear** if and only if it satisfies the following two conditions:

- $F(x + y) = F(x) + F(y) \quad \forall x, y \in \mathbb{Q}^n.$
- $F(\lambda x) = \lambda F(x) \quad \forall x \in \mathbb{Q}^n \quad \forall \lambda \in \mathbb{Q}.$

Example: For  $n \in \mathbb{N}$  suppose  $(a_1, \dots, a_n) \in \mathbb{Q}^n$  represents the amplitudes of the different harmonics of a note played on a violin. Thus  $a_n$  is the amplitude of frequency  $nf$ , where  $f$  is the fundamental frequency of the note.

Then for  $m \in \mathbb{N}$  with  $m \leq n$  the function  $F$  specified by

$$\begin{aligned} F : \mathbb{Q}^n &\rightarrow \mathbb{Q}^n \\ (a_1, \dots, a_n) &\mapsto (a_1, \dots, a_m, 0, 0, \dots, 0) \end{aligned}$$

is called a **filter**. (It filters out the high frequencies).



# Linear functions

Definition: A function  $F : \mathbb{Q}^n \rightarrow \mathbb{Q}^n$  is called **linear** if and only if it satisfies the following two conditions:

- $F(x + y) = F(x) + F(y) \quad \forall x, y \in \mathbb{Q}^n.$
- $F(\lambda x) = \lambda F(x) \quad \forall x \in \mathbb{Q}^n \quad \forall \lambda \in \mathbb{Q}.$

Example: For  $n \in \mathbb{N}$  suppose  $(a_1, \dots, a_n) \in \mathbb{Q}^n$  represents the amplitudes of the different harmonics of a note played on a violin. Thus  $a_n$  is the amplitude of frequency  $nf$ , where  $f$  is the fundamental frequency of the note.

Then for  $m \in \mathbb{N}$  with  $m \leq n$  the function  $F$  specified by

$$\begin{aligned} F : \mathbb{Q}^n &\rightarrow \mathbb{Q}^n \\ (a_1, \dots, a_n) &\mapsto (a_1, \dots, a_m, 0, 0, \dots, 0) \end{aligned}$$

is called a **filter**. (It filters out the high frequencies).

Filters are linear functions. (Check!)

## Linear functions: another example

Let  $(p_n)_{n \in \mathbb{N}} \subseteq \mathbb{Q}^2$  represent the state of an ecosystem with two species at time  $n$ ; say  $p_n = (x_n, y_n)$ , where  $x_n$  is the size of the population of species 1, and  $y_n$  the size of the population of species 2.

## Linear functions: another example

Let  $(p_n)_{n \in \mathbb{N}} \subseteq \mathbb{Q}^2$  represent the state of an ecosystem with two species at time  $n$ ; say  $p_n = (x_n, y_n)$ , where  $x_n$  is the size of the population of species 1, and  $y_n$  the size of the population of species 2.

Assume that the ecosystem evolves as follows, due to a predator-prey relationship between the two species:

$$\forall n \in \mathbb{N} \quad \begin{cases} x_{n+1} = 4x_n - y_n, \\ y_{n+1} = y_n + 2x_n. \end{cases}$$

## Linear functions: another example

Let  $(p_n)_{n \in \mathbb{N}} \subseteq \mathbb{Q}^2$  represent the state of an ecosystem with two species at time  $n$ ; say  $p_n = (x_n, y_n)$ , where  $x_n$  is the size of the population of species 1, and  $y_n$  the size of the population of species 2.

Assume that the ecosystem evolves as follows, due to a predator-prey relationship between the two species:

$$\forall n \in \mathbb{N} \quad \begin{cases} x_{n+1} = 4x_n - y_n, \\ y_{n+1} = y_n + 2x_n. \end{cases}$$

Then  $p_{n+1} = F(p_n) \quad \forall n \in \mathbb{N}$ , where  $F(x, y) = (4x - y, 2x + y)$ .

The function  $F$  is linear. (Check!)

## Linear functions: another example

Let  $(p_n)_{n \in \mathbb{N}} \subseteq \mathbb{Q}^2$  represent the state of an ecosystem with two species at time  $n$ ; say  $p_n = (x_n, y_n)$ , where  $x_n$  is the size of the population of species 1, and  $y_n$  the size of the population of species 2.

Assume that the ecosystem evolves as follows, due to a predator-prey relationship between the two species:

$$\forall n \in \mathbb{N} \quad \begin{cases} x_{n+1} = 4x_n - y_n, \\ y_{n+1} = y_n + 2x_n. \end{cases}$$

Then  $p_{n+1} = F(p_n) \quad \forall n \in \mathbb{N}$ , where  $F(x, y) = (4x - y, 2x + y)$ .

The function  $F$  is linear. (Check!)

We will return to this example.

# Multiplying a vector by a matrix: motivation

We now explore the possibility of expressing the function

$$\begin{aligned} F : \mathbb{Q}^2 &\rightarrow \mathbb{Q}^2 \\ (x, y) &\mapsto (4x - y, 2x + y) \end{aligned}$$

using a matrix.

# Multiplying a vector by a matrix: motivation

We now explore the possibility of expressing the function

$$\begin{aligned} F : \mathbb{Q}^2 &\rightarrow \mathbb{Q}^2 \\ (x, y) &\mapsto (4x - y, 2x + y) \end{aligned}$$

using a matrix. We would like to write  $F(x, y) = M(x, y)$  for some matrix  $M$  and a suitable meaning for the 'product'  $M(x, y)$ .

# Multiplying a vector by a matrix: motivation

We now explore the possibility of expressing the function

$$\begin{aligned} F : \mathbb{Q}^2 &\rightarrow \mathbb{Q}^2 \\ (x, y) &\mapsto (4x - y, 2x + y) \end{aligned}$$

using a matrix. We would like to write  $F(x, y) = M(x, y)$  for some matrix  $M$  and a suitable meaning for the 'product'  $M(x, y)$ .

Writing  $(x, y)$  as the column vector  $x = \begin{pmatrix} x \\ y \end{pmatrix}$ , we want

$$M \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 4x - y \\ 2x + y \end{pmatrix},$$



# Multiplying a vector by a matrix: motivation

We now explore the possibility of expressing the function

$$\begin{aligned} F : \mathbb{Q}^2 &\rightarrow \mathbb{Q}^2 \\ (x, y) &\mapsto (4x - y, 2x + y) \end{aligned}$$

using a matrix. We would like to write  $F(x, y) = M(x, y)$  for some matrix  $M$  and a suitable meaning for the 'product'  $M(x, y)$ .

Writing  $(x, y)$  as the column vector  $x = \begin{pmatrix} x \\ y \end{pmatrix}$ , we want

$$M \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 4x - y \\ 2x + y \end{pmatrix},$$

and so the 'coefficient matrix'  $\begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix}$  looks like a good candidate for  $M$

# Multiplying a vector by a matrix: motivation

We now explore the possibility of expressing the function

$$\begin{aligned} F : \mathbb{Q}^2 &\rightarrow \mathbb{Q}^2 \\ (x, y) &\mapsto (4x - y, 2x + y) \end{aligned}$$

using a matrix. We would like to write  $F(x, y) = M(x, y)$  for some matrix  $M$  and a suitable meaning for the 'product'  $M(x, y)$ .

Writing  $(x, y)$  as the column vector  $x = \begin{pmatrix} x \\ y \end{pmatrix}$ , we want

$$M \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 4x - y \\ 2x + y \end{pmatrix},$$

and so the 'coefficient matrix'  $\begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix}$  looks like a good candidate for  $M$  provided that we define the product  $Mx$  so that

$$\begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 4x - y \\ 2x + y \end{pmatrix}.$$

# Multiplying a vector by a matrix: motivation

We now explore the possibility of expressing the function

$$\begin{aligned} F : \mathbb{Q}^2 &\rightarrow \mathbb{Q}^2 \\ (x, y) &\mapsto (4x - y, 2x + y) \end{aligned}$$

using a matrix. We would like to write  $F(x, y) = M(x, y)$  for some matrix  $M$  and a suitable meaning for the 'product'  $M(x, y)$ .

Writing  $(x, y)$  as the column vector  $x = \begin{pmatrix} x \\ y \end{pmatrix}$ , we want

$$M \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 4x - y \\ 2x + y \end{pmatrix},$$

and so the 'coefficient matrix'  $\begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix}$  looks like a good candidate for  $M$  provided that we define the product  $Mx$  so that

$$\begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 4x - y \\ 2x + y \end{pmatrix}.$$

# Multiplying a vector by a matrix: definition

For a matrix  $A = (a_{i,j}) \in M_n(\mathbb{Q})$  and a vector  $x = (x_1, \dots, x_n) \in \mathbb{Q}^n$  we define the **matrix-vector product**  $Ax$  as the vector given by

$$Ax = \left( \sum_{j=1}^n a_{i,j} x_j \right)_{1 \leq i \leq n} \in \mathbb{Q}^n.$$

# Multiplying a vector by a matrix: definition

For a matrix  $A = (a_{i,j}) \in M_n(\mathbb{Q})$  and a vector  $x = (x_1, \dots, x_n) \in \mathbb{Q}^n$  we define the **matrix-vector product**  $Ax$  as the vector given by

$$Ax = \left( \sum_{j=1}^n a_{i,j} x_j \right)_{1 \leq i \leq n} \in \mathbb{Q}^n.$$

By convention, in this context, we normally write the vectors as columns.

Thus

$$\begin{bmatrix} a_{1,1} & a_{1,2} & \cdots & a_{1,n} \\ a_{2,1} & a_{2,2} & \cdots & a_{2,n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n,1} & a_{n,2} & \cdots & a_{n,n} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} a_{1,1}x_1 + a_{1,2}x_2 + \cdots + a_{1,n}x_n \\ a_{2,1}x_1 + a_{2,2}x_2 + \cdots + a_{2,n}x_n \\ \vdots \\ a_{n,1}x_1 + a_{n,2}x_2 + \cdots + a_{n,n}x_n \end{bmatrix}$$

# Multiplying a vector by a matrix: definition

For a matrix  $A = (a_{i,j}) \in M_n(\mathbb{Q})$  and a vector  $x = (x_1, \dots, x_n) \in \mathbb{Q}^n$  we define the **matrix-vector product**  $Ax$  as the vector given by

$$Ax = \left( \sum_{j=1}^n a_{i,j} x_j \right)_{1 \leq i \leq n} \in \mathbb{Q}^n.$$

By convention, in this context, we normally write the vectors as columns.

Thus

$$\begin{bmatrix} a_{1,1} & a_{1,2} & \cdots & a_{1,n} \\ a_{2,1} & a_{2,2} & \cdots & a_{2,n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n,1} & a_{n,2} & \cdots & a_{n,n} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} a_{1,1}x_1 + a_{1,2}x_2 + \cdots + a_{1,n}x_n \\ a_{2,1}x_1 + a_{2,2}x_2 + \cdots + a_{2,n}x_n \\ \vdots \\ a_{n,1}x_1 + a_{n,2}x_2 + \cdots + a_{n,n}x_n \end{bmatrix}$$

Example: 
$$\begin{pmatrix} 2 & 0 & -1 \\ 0 & -1 & 2 \\ -1 & 2 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} = \begin{pmatrix} -1 \\ 4 \\ 3 \end{pmatrix}.$$

## Linear functions expressed using matrices

Example: 
$$\begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 4x - y \\ 2x + y \end{pmatrix} = F(x, y)$$

where, as we have seen, the function  $F : \mathbb{Q} \rightarrow \mathbb{Q}$  so defined is linear.

## Linear functions expressed using matrices

Example: 
$$\begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 4x - y \\ 2x + y \end{pmatrix} = F(x, y)$$

where, as we have seen, the function  $F : \mathbb{Q} \rightarrow \mathbb{Q}$  so defined is linear.

This is no coincidence.



# Linear functions expressed using matrices

Example: 
$$\begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 4x - y \\ 2x + y \end{pmatrix} = F(x, y)$$

where, as we have seen, the function  $F : \mathbb{Q} \rightarrow \mathbb{Q}$  so defined is linear.

This is no coincidence.

**Theorem** (proof omitted): To each linear function  $F : \mathbb{Q}^n \rightarrow \mathbb{Q}^n$  there is a matrix  $M \in M_n(\mathbb{Q})$  such that

$$F(x) = Mx \quad \forall x \in \mathbb{Q}^n.$$

Conversely, every function  $F : \mathbb{Q}^n \rightarrow \mathbb{Q}^n$  defined using a matrix in this way is linear.

# Matrix multiplication

---

# Matrix multiplication: motivation

Question: Given  $M \in M_n(\mathbb{Q})$ , how, if at all, should  $M^2$  be defined?

# Matrix multiplication: motivation

Question: Given  $M \in M_n(\mathbb{Q})$ , how, if at all, should  $M^2$  be defined?

Discussion: For any  $x$  in  $\mathbb{Q}^n$ ,  $Mx$  is also in  $\mathbb{Q}^n$  and so we can consider  $M(Mx)$ . Surely we would like this to equal to  $M^2x$ .

# Matrix multiplication: motivation

Question: Given  $M \in M_n(\mathbb{Q})$ , how, if at all, should  $M^2$  be defined?

Discussion: For any  $x$  in  $\mathbb{Q}^n$ ,  $Mx$  is also in  $\mathbb{Q}^n$  and so we can consider  $M(Mx)$ . Surely we would like this to equal to  $M^2x$ .

Example: For  $M = \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix}$ :

$$\begin{aligned} M(Mx) &= \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \left[ \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \right] \\ &= \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} 4x - y \\ 2x + y \end{pmatrix} \end{aligned}$$

# Matrix multiplication: motivation

Question: Given  $M \in M_n(\mathbb{Q})$ , how, if at all, should  $M^2$  be defined?

Discussion: For any  $x$  in  $\mathbb{Q}^n$ ,  $Mx$  is also in  $\mathbb{Q}^n$  and so we can consider  $M(Mx)$ . Surely we would like this to equal to  $M^2x$ .

Example: For  $M = \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix}$ :

$$\begin{aligned} M(Mx) &= \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \left[ \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \right] \\ &= \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} 4x - y \\ 2x + y \end{pmatrix} \\ &= \begin{pmatrix} 14x - 5y \\ 10x - y \end{pmatrix} = \begin{pmatrix} 14 & -5 \\ 10 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \end{aligned}$$

# Matrix multiplication: motivation

Question: Given  $M \in M_n(\mathbb{Q})$ , how, if at all, should  $M^2$  be defined?

Discussion: For any  $x$  in  $\mathbb{Q}^n$ ,  $Mx$  is also in  $\mathbb{Q}^n$  and so we can consider  $M(Mx)$ . Surely we would like this to equal to  $M^2x$ .

Example: For  $M = \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix}$ :

$$\begin{aligned} M(Mx) &= \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \left[ \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \right] \\ &= \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} 4x - y \\ 2x + y \end{pmatrix} \\ &= \begin{pmatrix} 14x - 5y \\ 10x - y \end{pmatrix} = \begin{pmatrix} 14 & -5 \\ 10 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \end{aligned}$$

So we want  $M^2 = \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} = \begin{pmatrix} 14 & -5 \\ 10 & -1 \end{pmatrix}$ .

# Matrix multiplication: definition

For matrices  $A = (a_{i,j})$  and  $B = (b_{i,j})$  in  $M_n(\mathbb{Q})$  the **product**  $AB = C = (c_{i,j}) \in M_n(\mathbb{Q})$  is defined by

$$c_{i,j} = \sum_{k=1}^n a_{i,k} b_{k,j} \quad \forall i, j \in \{1, \dots, n\}.$$



# Matrix multiplication: definition

For matrices  $A = (a_{i,j})$  and  $B = (b_{i,j})$  in  $M_n(\mathbb{Q})$  the **product**  $AB = C = (c_{i,j}) \in M_n(\mathbb{Q})$  is defined by

$$c_{i,j} = \sum_{k=1}^n a_{i,k} b_{k,j} \quad \forall i, j \in \{1, \dots, n\}.$$

Two Examples:

- (a) First, let's check that this formula produces what we were looking for with  $M^2$  on the previous slide:

$$\begin{aligned} M^2 &= \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 4 \times 4 + (-1) \times 2 & 4 \times (-1) + (-1) \times 1 \\ 2 \times 4 + 1 \times 2 & 2 \times (-1) + 1 \times 1 \end{pmatrix} = \begin{pmatrix} 14 & -5 \\ 10 & -1 \end{pmatrix}. \end{aligned}$$

# Matrix multiplication: definition

For matrices  $A = (a_{i,j})$  and  $B = (b_{i,j})$  in  $M_n(\mathbb{Q})$  the **product**  $AB = C = (c_{i,j}) \in M_n(\mathbb{Q})$  is defined by

$$c_{i,j} = \sum_{k=1}^n a_{i,k} b_{k,j} \quad \forall i, j \in \{1, \dots, n\}.$$

Two Examples:

- (a) First, let's check that this formula produces what we were looking for with  $M^2$  on the previous slide:

$$\begin{aligned} M^2 &= \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 4 \times 4 + (-1) \times 2 & 4 \times (-1) + (-1) \times 1 \\ 2 \times 4 + 1 \times 2 & 2 \times (-1) + 1 \times 1 \end{pmatrix} = \begin{pmatrix} 14 & -5 \\ 10 & -1 \end{pmatrix}. \end{aligned}$$

- (b) This example demonstrates the product formula more clearly:

$$\begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} a + 2c & b + 2d \\ 3a + 4c & 3b + 4d \end{pmatrix}.$$

# Identity matrices

Observe that the matrix  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$  acts as an 'identity' in the sense

that  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$  for any matrix  $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ .

# Identity matrices

Observe that the matrix  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$  acts as an 'identity' in the sense

that  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$  for any matrix  $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ .

More generally, for  $n \in \mathbb{N}$ , we define the  $n \times n$  **identity matrix**  $I_n$  by

$$I_n = (\delta_{i,j}) \in M_n(\mathbb{Q}) \text{ with } \delta_{i,j} = \begin{cases} 1 & \text{if } i = j, \\ 0 & \text{otherwise.} \end{cases}$$

# Identity matrices

Observe that the matrix  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$  acts as an 'identity' in the sense

that  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$  for any matrix  $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ .

More generally, for  $n \in \mathbb{N}$ , we define the  $n \times n$  **identity matrix**  $I_n$  by

$$I_n = (\delta_{i,j}) \in M_n(\mathbb{Q}) \text{ with } \delta_{i,j} = \begin{cases} 1 & \text{if } i = j, \\ 0 & \text{otherwise.} \end{cases}$$

So  $I_1 = [1]$ ,  $I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ ,  $I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ , etc.

# Identity matrices

Observe that the matrix  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$  acts as an 'identity' in the sense

that  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$  for any matrix  $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ .

More generally, for  $n \in \mathbb{N}$ , we define the  $n \times n$  **identity matrix**  $I_n$  by

$$I_n = (\delta_{i,j}) \in M_n(\mathbb{Q}) \text{ with } \delta_{i,j} = \begin{cases} 1 & \text{if } i = j, \\ 0 & \text{otherwise.} \end{cases}$$

So  $I_1 = [1]$ ,  $I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ ,  $I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ , etc.

By applying the matrix product formula we can immediately establish that, for any  $n \in \mathbb{N}$ , the identity matrix  $I_n$  does indeed have the identity property:

$$\forall n \in \mathbb{N}, \forall M \in M_n(\mathbb{Q}) \quad I_n M = M = M I_n.$$

Remark: When the value of  $n$  is clear from the context, we abbreviate  $I_n$  to just  $I$ .

## Matrix multiplication ....

**is not commutative:** *i.e.*  $MN \neq NM$  in general.

# Matrix multiplication ....

**is not commutative:** *I.e.*  $MN \neq NM$  in general.

$$E.g \text{ take } M = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \text{ and } N = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}.$$

$$\text{Then } MN = \begin{pmatrix} 1 & 2 \\ 1 & 2 \end{pmatrix} \text{ but } NM = \begin{pmatrix} 1 & 1 \\ 2 & 2 \end{pmatrix}.$$



# Matrix multiplication ....

**is not commutative:** *i.e.*  $MN \neq NM$  in general.

$$\text{E.g take } M = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \text{ and } N = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}.$$

$$\text{Then } MN = \begin{pmatrix} 1 & 2 \\ 1 & 2 \end{pmatrix} \text{ but } NM = \begin{pmatrix} 1 & 1 \\ 2 & 2 \end{pmatrix}.$$

**is associative:** *i.e.*  $\forall n \in \mathbb{N} \forall M, N, P \in M_n(\mathbb{Q}) \quad M(NP) = (MN)P$ .

# Matrix multiplication ....

**is not commutative:** *I.e.*  $MN \neq NM$  in general.

$$E.g \text{ take } M = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \text{ and } N = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}.$$

$$\text{Then } MN = \begin{pmatrix} 1 & 2 \\ 1 & 2 \end{pmatrix} \text{ but } NM = \begin{pmatrix} 1 & 1 \\ 2 & 2 \end{pmatrix}.$$

**is associative:** *I.e.*  $\forall n \in \mathbb{N} \forall M, N, P \in M_n(\mathbb{Q}) \quad M(NP) = (MN)P$ .

This follows from the one-to-one correspondence between matrices and linear functions, the way that matrix multiplication reflects function composition, and the fact that function composition is associative.

# Matrix multiplication ....

**is not commutative:** *I.e.*  $MN \neq NM$  in general.

$$E.g \text{ take } M = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \text{ and } N = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}.$$

$$\text{Then } MN = \begin{pmatrix} 1 & 2 \\ 1 & 2 \end{pmatrix} \text{ but } NM = \begin{pmatrix} 1 & 1 \\ 2 & 2 \end{pmatrix}.$$

**is associative:** *I.e.*  $\forall n \in \mathbb{N} \forall M, N, P \in M_n(\mathbb{Q}) \quad M(NP) = (MN)P$ .

This follows from the one-to-one correspondence between matrices and linear functions, the way that matrix multiplication reflects function composition, and the fact that function composition is associative.

**has 'zero divisors':** *I.e.*  $MN = O \not\Rightarrow (M = O) \vee (N = O)$ .

# Matrix multiplication ....

**is not commutative:** *I.e.*  $MN \neq NM$  in general.

*E.g.* take  $M = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$  and  $N = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$ .

Then  $MN = \begin{pmatrix} 1 & 2 \\ 1 & 2 \end{pmatrix}$  but  $NM = \begin{pmatrix} 1 & 1 \\ 2 & 2 \end{pmatrix}$ .

**is associative:** *I.e.*  $\forall n \in \mathbb{N} \forall M, N, P \in M_n(\mathbb{Q}) \quad M(NP) = (MN)P$ .

This follows from the one-to-one correspondence between matrices and linear functions, the way that matrix multiplication reflects function composition, and the fact that function composition is associative.

**has 'zero divisors':** *I.e.*  $MN = O \not\Rightarrow (M = O) \vee (N = O)$ .

*E.g.* take  $M = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$  and  $N = \begin{pmatrix} 1 & 1 \\ -1 & -1 \end{pmatrix}$ .

Then  $MN = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$  but  $M \neq \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \neq N$ .

## Inverses: motivation

In a certain chemical reaction between two reactants, R1 and R2:

- (a) The quantity of R1 should be twice the quantity of R2.

## Inverses: motivation

In a certain chemical reaction between two reactants, R1 and R2:

- (a) The quantity of R1 should be twice the quantity of R2.
- (b) The quantity of product is then equal to half the quantity of R1 plus one third of the quantity of R2.

## Inverses: motivation

In a certain chemical reaction between two reactants, R1 and R2:

- (a) The quantity of R1 should be twice the quantity of R2.
- (b) The quantity of product is then equal to half the quantity of R1 plus one third of the quantity of R2.

What quantities of R1 and R2 should be used to produce 5 units of the product?

## Inverses: motivation

In a certain chemical reaction between two reactants, R1 and R2:

- (a) The quantity of R1 should be twice the quantity of R2.
- (b) The quantity of product is then equal to half the quantity of R1 plus one third of the quantity of R2.

What quantities of R1 and R2 should be used to produce 5 units of the product?

Let  $x$  be the quantity of R1,  $y$  quantity of R2.

$$\begin{cases} \text{from (a): } x = 2y \\ \text{from (b): } \frac{x}{2} + \frac{y}{3} = 5 \end{cases} \iff \begin{cases} x - 2y = 0 \\ 3x + 2y = 30 \end{cases}$$



## Inverses: motivation

In a certain chemical reaction between two reactants, R1 and R2:

- (a) The quantity of R1 should be twice the quantity of R2.
- (b) The quantity of product is then equal to half the quantity of R1 plus one third of the quantity of R2.

What quantities of R1 and R2 should be used to produce 5 units of the product?

Let  $x$  be the quantity of R1,  $y$  quantity of R2.

$$\begin{cases} \text{from (a): } x = 2y \\ \text{from (b): } \frac{x}{2} + \frac{y}{3} = 5 \end{cases} \iff \begin{cases} x - 2y = 0 \\ 3x + 2y = 30 \end{cases}$$

We can solve these equations by elimination, but consider the equivalent matrix equation

$$\begin{pmatrix} 1 & -2 \\ 3 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 30 \end{pmatrix}.$$

Q: Can we solve this matrix equation, just using matrices?

# Inverses

Question: More generally, can we solve a matrix equation  $Ax = b$ , just using matrices?

# Inverses

Question: More generally, can we solve a matrix equation  $Ax = b$ , just using matrices? *i.e.* Can we “divide” by  $A$  to get  $x = \frac{b}{A}$ ?

# Inverses

Question: More generally, can we solve a matrix equation  $Ax = b$ , just using matrices? *i.e.* Can we “divide” by  $A$  to get  $x = \frac{b}{A}$ ?

Answer: Well, sort of, sometimes.

# Inverses

Question: More generally, can we solve a matrix equation  $Ax = b$ , just using matrices? *i.e.* Can we “divide” by  $A$  to get  $x = \frac{b}{A}$ ?

Answer: Well, sort of, sometimes.

In some cases there exists a matrix  $A^{-1}$  such that  $x = A^{-1}b$ .

# Inverses

Question: More generally, can we solve a matrix equation  $Ax = b$ , just using matrices? *i.e.* Can we “divide” by  $A$  to get  $x = \frac{b}{A}$ ?

Answer: Well, sort of, sometimes.

In some cases there exists a matrix  $A^{-1}$  such that  $x = A^{-1}b$ .

Our chemical reaction example is a case in point:

$$\begin{pmatrix} 1 & -2 \\ 3 & 2 \end{pmatrix}^{-1} = \frac{1}{8} \begin{pmatrix} 2 & 2 \\ -3 & 1 \end{pmatrix} \quad \begin{array}{l} \text{(we will see how} \\ \text{to get this later)} \end{array}$$

# Inverses

Question: More generally, can we solve a matrix equation  $Ax = b$ , just using matrices? *i.e.* Can we “divide” by  $A$  to get  $x = \frac{b}{A}$ ?

Answer: Well, sort of, sometimes.

In some cases there exists a matrix  $A^{-1}$  such that  $x = A^{-1}b$ .

Our chemical reaction example is a case in point:

$$\begin{pmatrix} 1 & -2 \\ 3 & 2 \end{pmatrix}^{-1} = \frac{1}{8} \begin{pmatrix} 2 & 2 \\ -3 & 1 \end{pmatrix} \quad \begin{matrix} \text{(we will see how} \\ \text{to get this later)} \end{matrix}$$

so  $\begin{pmatrix} x \\ y \end{pmatrix} = \frac{1}{8} \begin{pmatrix} 2 & 2 \\ -3 & 1 \end{pmatrix} \begin{pmatrix} 0 \\ 30 \end{pmatrix} = \begin{pmatrix} 7.5 \\ 3.75 \end{pmatrix}.$

# Inverses

Question: More generally, can we solve a matrix equation  $Ax = b$ , just using matrices? *i.e.* Can we “divide” by  $A$  to get  $x = \frac{b}{A}$ ?

Answer: Well, sort of, sometimes.

In some cases there exists a matrix  $A^{-1}$  such that  $x = A^{-1}b$ .

Our chemical reaction example is a case in point:

$$\begin{pmatrix} 1 & -2 \\ 3 & 2 \end{pmatrix}^{-1} = \frac{1}{8} \begin{pmatrix} 2 & 2 \\ -3 & 1 \end{pmatrix} \quad \begin{matrix} \text{(we will see how} \\ \text{to get this later)} \end{matrix}$$

so  $\begin{pmatrix} x \\ y \end{pmatrix} = \frac{1}{8} \begin{pmatrix} 2 & 2 \\ -3 & 1 \end{pmatrix} \begin{pmatrix} 0 \\ 30 \end{pmatrix} = \begin{pmatrix} 7.5 \\ 3.75 \end{pmatrix}.$

This matrix  $A^{-1}$  is an ‘inverse’ of  $A$  in the following sense:

An **inverse**, if one exists, of a matrix  $A \in M_n(\mathbb{Q})$  is a matrix  $A^{-1} \in M_n(\mathbb{Q})$  with the property that  $A^{-1}A = AA^{-1} = I_n$ .



# Inverses

Question: More generally, can we solve a matrix equation  $Ax = b$ , just using matrices? *i.e.* Can we “divide” by  $A$  to get  $x = \frac{b}{A}$ ?

Answer: Well, sort of, sometimes.

In some cases there exists a matrix  $A^{-1}$  such that  $x = A^{-1}b$ .

Our chemical reaction example is a case in point:

$$\begin{pmatrix} 1 & -2 \\ 3 & 2 \end{pmatrix}^{-1} = \frac{1}{8} \begin{pmatrix} 2 & 2 \\ -3 & 1 \end{pmatrix} \quad \begin{matrix} \text{(we will see how} \\ \text{to get this later)} \end{matrix}$$

so  $\begin{pmatrix} x \\ y \end{pmatrix} = \frac{1}{8} \begin{pmatrix} 2 & 2 \\ -3 & 1 \end{pmatrix} \begin{pmatrix} 0 \\ 30 \end{pmatrix} = \begin{pmatrix} 7.5 \\ 3.75 \end{pmatrix}.$

This matrix  $A^{-1}$  is an ‘inverse’ of  $A$  in the following sense:

An **inverse**, if one exists, of a matrix  $A \in M_n(\mathbb{Q})$  is a matrix  $A^{-1} \in M_n(\mathbb{Q})$  with the property that  $A^{-1}A = AA^{-1} = I_n$ .

Note that if an  $A^{-1}$  exists and  $Ax = b$  then

$$x = Ix = (A^{-1}A)x = A^{-1}(Ax) = A^{-1}b.$$

# Determinants

Not every square matrix has an inverse.

# Determinants

Not every square matrix has an inverse.

An obvious example is a zero matrix, but there are also plenty of non-zero examples.

# Determinants

Not every square matrix has an inverse.

An obvious example is a zero matrix, but there are also plenty of non-zero examples.

Question: How can we tell (determine) if  $A$  has an inverse?

# Determinants

Not every square matrix has an inverse.

An obvious example is a zero matrix, but there are also plenty of non-zero examples.

Question: How can we tell (determine) if  $A$  has an inverse?

As we shall see, one answer is that  $A$  has an inverse if and only if its 'determinant' is non-zero.

# Determinants

Not every square matrix has an inverse.

An obvious example is a zero matrix, but there are also plenty of non-zero examples.

Question: How can we tell (determine) if  $A$  has an inverse?

As we shall see, one answer is that  $A$  has an inverse if and only if its 'determinant' is non-zero.

In this course we only consider the  $2 \times 2$  case:

The **determinant** of a matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_2(\mathbb{Q})$  is the number

$$\det(A) = ad - bc.$$

# Determinants

Not every square matrix has an inverse.

An obvious example is a zero matrix, but there are also plenty of non-zero examples.

Question: How can we tell (determine) if  $A$  has an inverse?

As we shall see, one answer is that  $A$  has an inverse if and only if its 'determinant' is non-zero.

In this course we only consider the  $2 \times 2$  case:

The **determinant** of a matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_2(\mathbb{Q})$  is the number

$$\det(A) = ad - bc.$$

Example:  $\det \begin{pmatrix} 1 & -2 \\ 3 & 2 \end{pmatrix} = 1 \times 2 - (-2) \times 3 = 8.$

# Determinants

Not every square matrix has an inverse.

An obvious example is a zero matrix, but there are also plenty of non-zero examples.

Question: How can we tell (determine) if  $A$  has an inverse?

As we shall see, one answer is that  $A$  has an inverse if and only if its 'determinant' is non-zero.

In this course we only consider the  $2 \times 2$  case:

The **determinant** of a matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_2(\mathbb{Q})$  is the number

$$\det(A) = ad - bc.$$

Example:  $\det \begin{pmatrix} 1 & -2 \\ 3 & 2 \end{pmatrix} = 1 \times 2 - (-2) \times 3 = 8.$

Lemma: For any  $A, B \in M_2(\mathbb{Q})$ ,  $\det(AB) = \det(A) \det(B)$ .

Proof: Multiply out both sides.



# Calculating Inverses

A matrix  $A$  can have at most one inverse, because if  $B$  and  $C$  are both inverses then  $BA = I$  and  $AC = I$  and so

# Calculating Inverses

A matrix  $A$  can have at most one inverse, because if  $B$  and  $C$  are both inverses then  $BA = I$  and  $AC = I$  and so

$$B = BI$$

# Calculating Inverses

A matrix  $A$  can have at most one inverse, because if  $B$  and  $C$  are both inverses then  $BA = I$  and  $AC = I$  and so

$$B = BI = B(AC)$$

# Calculating Inverses

A matrix  $A$  can have at most one inverse, because if  $B$  and  $C$  are both inverses then  $BA = I$  and  $AC = I$  and so

$$B = BI = B(AC) = (BA)C$$

# Calculating Inverses

A matrix  $A$  can have at most one inverse, because if  $B$  and  $C$  are both inverses then  $BA = I$  and  $AC = I$  and so

$$B = BI = B(AC) = (BA)C = IC$$

# Calculating Inverses

A matrix  $A$  can have at most one inverse, because if  $B$  and  $C$  are both inverses then  $BA = I$  and  $AC = I$  and so

$$B = BI = B(AC) = (BA)C = IC = C.$$

# Calculating Inverses

A matrix  $A$  can have at most one inverse, because if  $B$  and  $C$  are both inverses then  $BA = I$  and  $AC = I$  and so

$$B = BI = B(AC) = (BA)C = IC = C.$$

How can we compute (the unique)  $A^{-1}$  when it does exist?

# Calculating Inverses

A matrix  $A$  can have at most one inverse, because if  $B$  and  $C$  are both inverses then  $BA = I$  and  $AC = I$  and so

$$B = BI = B(AC) = (BA)C = IC = C.$$

How can we compute (the unique)  $A^{-1}$  when it does exist?

Theorem: A matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_2(\mathbb{Q})$  has an inverse if and only if

$\det(A) \neq 0$  and in this case

$$A^{-1} = \frac{1}{\det(A)} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} \quad \text{e.g.} \quad \begin{pmatrix} 1 & -2 \\ 3 & 2 \end{pmatrix}^{-1} = \frac{1}{8} \begin{pmatrix} 2 & 2 \\ -3 & 1 \end{pmatrix}$$



# Calculating Inverses

A matrix  $A$  can have at most one inverse, because if  $B$  and  $C$  are both inverses then  $BA = I$  and  $AC = I$  and so

$$B = BI = B(AC) = (BA)C = IC = C.$$

How can we compute (the unique)  $A^{-1}$  when it does exist?

Theorem: A matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_2(\mathbb{Q})$  has an inverse if and only if  $\det(A) \neq 0$  and in this case

$$A^{-1} = \frac{1}{\det(A)} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} \quad \text{e.g.} \quad \begin{pmatrix} 1 & -2 \\ 3 & 2 \end{pmatrix}^{-1} = \frac{1}{8} \begin{pmatrix} 2 & 2 \\ -3 & 1 \end{pmatrix}$$

Proof: If  $A$  has an inverse then  
 $1 = \det(I_2)$

# Calculating Inverses

A matrix  $A$  can have at most one inverse, because if  $B$  and  $C$  are both inverses then  $BA = I$  and  $AC = I$  and so

$$B = BI = B(AC) = (BA)C = IC = C.$$

How can we compute (the unique)  $A^{-1}$  when it does exist?

Theorem: A matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_2(\mathbb{Q})$  has an inverse if and only if  $\det(A) \neq 0$  and in this case

$$A^{-1} = \frac{1}{\det(A)} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} \quad \text{e.g.} \quad \begin{pmatrix} 1 & -2 \\ 3 & 2 \end{pmatrix}^{-1} = \frac{1}{8} \begin{pmatrix} 2 & 2 \\ -3 & 1 \end{pmatrix}$$

Proof: If  $A$  has an inverse then

$$1 = \det(I_2) = \det(AA^{-1})$$

# Calculating Inverses

A matrix  $A$  can have at most one inverse, because if  $B$  and  $C$  are both inverses then  $BA = I$  and  $AC = I$  and so

$$B = BI = B(AC) = (BA)C = IC = C.$$

How can we compute (the unique)  $A^{-1}$  when it does exist?

Theorem: A matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_2(\mathbb{Q})$  has an inverse if and only if  $\det(A) \neq 0$  and in this case

$$A^{-1} = \frac{1}{\det(A)} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} \quad \text{e.g.} \quad \begin{pmatrix} 1 & -2 \\ 3 & 2 \end{pmatrix}^{-1} = \frac{1}{8} \begin{pmatrix} 2 & 2 \\ -3 & 1 \end{pmatrix}$$

Proof: If  $A$  has an inverse then

$$1 = \det(I_2) = \det(AA^{-1}) = \det(A) \det(A^{-1}),$$

# Calculating Inverses

A matrix  $A$  can have at most one inverse, because if  $B$  and  $C$  are both inverses then  $BA = I$  and  $AC = I$  and so

$$B = BI = B(AC) = (BA)C = IC = C.$$

How can we compute (the unique)  $A^{-1}$  when it does exist?

Theorem: A matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_2(\mathbb{Q})$  has an inverse if and only if  $\det(A) \neq 0$  and in this case

$$A^{-1} = \frac{1}{\det(A)} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} \quad \text{e.g.} \quad \begin{pmatrix} 1 & -2 \\ 3 & 2 \end{pmatrix}^{-1} = \frac{1}{8} \begin{pmatrix} 2 & 2 \\ -3 & 1 \end{pmatrix}$$

Proof: If  $A$  has an inverse then

$$1 = \det(I_2) = \det(AA^{-1}) = \det(A) \det(A^{-1}),$$

so  $\det(A)$  cannot be zero.

# Calculating Inverses

A matrix  $A$  can have at most one inverse, because if  $B$  and  $C$  are both inverses then  $BA = I$  and  $AC = I$  and so

$$B = BI = B(AC) = (BA)C = IC = C.$$

How can we compute (the unique)  $A^{-1}$  when it does exist?

Theorem: A matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_2(\mathbb{Q})$  has an inverse if and only if  $\det(A) \neq 0$  and in this case

$$A^{-1} = \frac{1}{\det(A)} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} \quad \text{e.g.} \quad \begin{pmatrix} 1 & -2 \\ 3 & 2 \end{pmatrix}^{-1} = \frac{1}{8} \begin{pmatrix} 2 & 2 \\ -3 & 1 \end{pmatrix}$$

Proof: If  $A$  has an inverse then

$$1 = \det(I_2) = \det(AA^{-1}) = \det(A) \det(A^{-1}),$$

so  $\det(A)$  cannot be zero. But if  $\det(A) \neq 0$  then multiplying out shows that  $AA^{-1} = A^{-1}A = I$  when  $A^{-1}$  is given by the formula.

# Calculating Inverses

A matrix  $A$  can have at most one inverse, because if  $B$  and  $C$  are both inverses then  $BA = I$  and  $AC = I$  and so

$$B = BI = B(AC) = (BA)C = IC = C.$$

How can we compute (the unique)  $A^{-1}$  when it does exist?

Theorem: A matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_2(\mathbb{Q})$  has an inverse if and only if

$\det(A) \neq 0$  and in this case

$$A^{-1} = \frac{1}{\det(A)} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} \quad \text{e.g.} \quad \begin{pmatrix} 1 & -2 \\ 3 & 2 \end{pmatrix}^{-1} = \frac{1}{8} \begin{pmatrix} 2 & 2 \\ -3 & 1 \end{pmatrix}$$

Proof: If  $A$  has an inverse then

$$1 = \det(I_2) = \det(AA^{-1}) = \det(A) \det(A^{-1}),$$

so  $\det(A)$  cannot be zero. But if  $\det(A) \neq 0$  then multiplying out shows that  $AA^{-1} = A^{-1}A = I$  when  $A^{-1}$  is given by the formula.

What about  $n > 2$ ?

# Calculating Inverses

A matrix  $A$  can have at most one inverse, because if  $B$  and  $C$  are both inverses then  $BA = I$  and  $AC = I$  and so

$$B = BI = B(AC) = (BA)C = IC = C.$$

How can we compute (the unique)  $A^{-1}$  when it does exist?

Theorem: A matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_2(\mathbb{Q})$  has an inverse if and only if  $\det(A) \neq 0$  and in this case

$$A^{-1} = \frac{1}{\det(A)} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} \quad \text{e.g.} \quad \begin{pmatrix} 1 & -2 \\ 3 & 2 \end{pmatrix}^{-1} = \frac{1}{8} \begin{pmatrix} 2 & 2 \\ -3 & 1 \end{pmatrix}$$

Proof: If  $A$  has an inverse then

$$1 = \det(I_2) = \det(AA^{-1}) = \det(A) \det(A^{-1}),$$

so  $\det(A)$  cannot be zero. But if  $\det(A) \neq 0$  then multiplying out shows that  $AA^{-1} = A^{-1}A = I$  when  $A^{-1}$  is given by the formula.

What about  $n > 2$ ? See Math1013 or Math1115.

## Interesting Examples

---



## Back to population dynamics

As a final example involving matrix multiplication and matrix inverses, we return to the simple ecosystem model

$$\forall n \in \mathbb{N} \quad \begin{cases} x_{n+1} = 4x_n - y_n, \\ y_{n+1} = y_n + 2x_n, \end{cases}$$

where  $x_n, y_n$  are the populations of two species after  $n$  time steps.

## Back to population dynamics

As a final example involving matrix multiplication and matrix inverses, we return to the simple ecosystem model

$$\forall n \in \mathbb{N} \quad \begin{cases} x_{n+1} = 4x_n - y_n, \\ y_{n+1} = y_n + 2x_n, \end{cases}$$

where  $x_n, y_n$  are the populations of two species after  $n$  time steps.

We can rewrite this in the form

$$\forall n \in \mathbb{N} \quad \begin{pmatrix} x_{n+1} \\ y_{n+1} \end{pmatrix} = \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x_n \\ y_n \end{pmatrix}.$$

## Back to population dynamics

As a final example involving matrix multiplication and matrix inverses, we return to the simple ecosystem model

$$\forall n \in \mathbb{N} \quad \begin{cases} x_{n+1} = 4x_n - y_n, \\ y_{n+1} = y_n + 2x_n, \end{cases}$$

where  $x_n, y_n$  are the populations of two species after  $n$  time steps.

We can rewrite this in the form

$$\forall n \in \mathbb{N} \quad \begin{pmatrix} x_{n+1} \\ y_{n+1} \end{pmatrix} = \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x_n \\ y_n \end{pmatrix}.$$

This is an implicit definition of a sequence of vectors. We will use mathematical induction to establish an explicit formula, stated and proved on the next slide.

## Back to population dynamics

As a final example involving matrix multiplication and matrix inverses, we return to the simple ecosystem model

$$\forall n \in \mathbb{N} \quad \begin{cases} x_{n+1} = 4x_n - y_n, \\ y_{n+1} = y_n + 2x_n, \end{cases}$$

where  $x_n, y_n$  are the populations of two species after  $n$  time steps.

We can rewrite this in the form

$$\forall n \in \mathbb{N} \quad \begin{pmatrix} x_{n+1} \\ y_{n+1} \end{pmatrix} = \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x_n \\ y_n \end{pmatrix}.$$

This is an implicit definition of a sequence of vectors. We will use mathematical induction to establish an explicit formula, stated and proved on the next slide. First two preliminary results

$$\text{R1: } \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 3 & 0 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \quad \left[ \begin{array}{l} \text{prove by multiplying} \\ \text{out the RHS} \end{array} \right]$$

## Back to population dynamics

As a final example involving matrix multiplication and matrix inverses, we return to the simple ecosystem model

$$\forall n \in \mathbb{N} \quad \begin{cases} x_{n+1} = 4x_n - y_n, \\ y_{n+1} = y_n + 2x_n, \end{cases}$$

where  $x_n, y_n$  are the populations of two species after  $n$  time steps.

We can rewrite this in the form

$$\forall n \in \mathbb{N} \quad \begin{pmatrix} x_{n+1} \\ y_{n+1} \end{pmatrix} = \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x_n \\ y_n \end{pmatrix}.$$

This is an implicit definition of a sequence of vectors. We will use mathematical induction to establish an explicit formula, stated and proved on the next slide. First two preliminary results

$$\text{R1: } \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 3 & 0 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \quad \left[ \begin{array}{l} \text{prove by multiplying} \\ \text{out the RHS} \end{array} \right]$$

$$\text{R2: } \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}^{-1} = \frac{1}{2-1} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} = \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \quad \left[ \begin{array}{l} \text{formula for} \\ \text{inverse of} \\ 2 \times 2 \text{ matrix} \end{array} \right]$$

**Claim:**  $\forall n \in \mathbb{N}^* \quad \begin{pmatrix} x_n \\ y_n \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 3^n & 0 \\ 0 & 2^n \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix}$

**Claim:**  $\forall n \in \mathbb{N}^* \quad \begin{pmatrix} x_n \\ y_n \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 3^n & 0 \\ 0 & 2^n \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix}$

**Proof:**

**Basis step:** When  $n = 0$  the RHS becomes (using R2)

$$\begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}^{-1} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix}$$

**Claim:**  $\forall n \in \mathbb{N}^* \quad \begin{pmatrix} x_n \\ y_n \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 3^n & 0 \\ 0 & 2^n \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix}$

**Proof:**

**Basis step:** When  $n = 0$  the RHS becomes (using R2)

$$\begin{aligned} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} &= \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}^{-1} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} = \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} = \text{LHS}. \end{aligned}$$



**Claim:**  $\forall n \in \mathbb{N}^* \quad \begin{pmatrix} x_n \\ y_n \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 3^n & 0 \\ 0 & 2^n \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix}$

**Proof:**

**Basis step:** When  $n = 0$  the RHS becomes (using R2)

$$\begin{aligned} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} &= \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}^{-1} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} = \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} = \text{LHS}. \end{aligned}$$

**Inductive step:** Assume the explicit formula holds up to and including some particular  $n$ , and consider the case  $n + 1$ .

**Claim:**  $\forall n \in \mathbb{N}^* \quad \begin{pmatrix} x_n \\ y_n \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 3^n & 0 \\ 0 & 2^n \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix}$

**Proof:**

**Basis step:** When  $n = 0$  the RHS becomes (using R2)

$$\begin{aligned} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} &= \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}^{-1} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} = \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} = \text{LHS}. \end{aligned}$$

**Inductive step:** Assume the explicit formula holds up to and including some particular  $n$ , and consider the case  $n + 1$ . Then, using the implicit definition, preliminary results R1 and R2, and the inductive assumption,

**Claim:**  $\forall n \in \mathbb{N}^* \quad \begin{pmatrix} x_n \\ y_n \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 3^n & 0 \\ 0 & 2^n \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix}$

**Proof:**

**Basis step:** When  $n = 0$  the RHS becomes (using R2)

$$\begin{aligned} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} &= \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}^{-1} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} = \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} = \text{LHS}. \end{aligned}$$

**Inductive step:** Assume the explicit formula holds up to and including some particular  $n$ , and consider the case  $n + 1$ . Then, using the implicit definition, preliminary results R1 and R2, and the inductive assumption,

$$\begin{pmatrix} x_{n+1} \\ y_{n+1} \end{pmatrix} = \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x_n \\ y_n \end{pmatrix}$$

**Claim:**  $\forall n \in \mathbb{N}^* \quad \begin{pmatrix} x_n \\ y_n \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 3^n & 0 \\ 0 & 2^n \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix}$

**Proof:**

**Basis step:** When  $n = 0$  the RHS becomes (using R2)

$$\begin{aligned} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} &= \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}^{-1} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} = \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} = \text{LHS}. \end{aligned}$$

**Inductive step:** Assume the explicit formula holds up to and including some particular  $n$ , and consider the case  $n + 1$ . Then, using the implicit definition, preliminary results R1 and R2, and the inductive assumption,

$$\begin{pmatrix} x_{n+1} \\ y_{n+1} \end{pmatrix} = \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x_n \\ y_n \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 3 & 0 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_n \\ y_n \end{pmatrix}$$

**Claim:**  $\forall n \in \mathbb{N}^* \quad \begin{pmatrix} x_n \\ y_n \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 3^n & 0 \\ 0 & 2^n \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix}$

**Proof:**

**Basis step:** When  $n = 0$  the RHS becomes (using R2)

$$\begin{aligned} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} &= \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}^{-1} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} = \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} = \text{LHS}. \end{aligned}$$

**Inductive step:** Assume the explicit formula holds up to and including some particular  $n$ , and consider the case  $n + 1$ . Then, using the implicit definition, preliminary results R1 and R2, and the inductive assumption,

$$\begin{aligned} \begin{pmatrix} x_{n+1} \\ y_{n+1} \end{pmatrix} &= \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x_n \\ y_n \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 3 & 0 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_n \\ y_n \end{pmatrix} \\ &= \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 3 & 0 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}^{-1} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 3^n & 0 \\ 0 & 2^n \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} \end{aligned}$$

**Claim:**  $\forall n \in \mathbb{N}^* \quad \begin{pmatrix} x_n \\ y_n \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 3^n & 0 \\ 0 & 2^n \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix}$

**Proof:**

**Basis step:** When  $n = 0$  the RHS becomes (using R2)

$$\begin{aligned} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} &= \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}^{-1} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} = \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} = \text{LHS.} \end{aligned}$$

**Inductive step:** Assume the explicit formula holds up to and including some particular  $n$ , and consider the case  $n + 1$ . Then, using the implicit definition, preliminary results R1 and R2, and the inductive assumption,

$$\begin{aligned} \begin{pmatrix} x_{n+1} \\ y_{n+1} \end{pmatrix} &= \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x_n \\ y_n \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 3 & 0 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_n \\ y_n \end{pmatrix} \\ &= \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 3 & 0 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}^{-1} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 3^n & 0 \\ 0 & 2^n \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 3^{n+1} & 0 \\ 0 & 2^{n+1} \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} \quad \text{and hence the formula} \\ &\quad \text{also holds for } n + 1. \end{aligned}$$

**Claim:**  $\forall n \in \mathbb{N}^* \quad \begin{pmatrix} x_n \\ y_n \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 3^n & 0 \\ 0 & 2^n \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix}$

**Proof:**

**Basis step:** When  $n = 0$  the RHS becomes (using R2)

$$\begin{aligned} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} &= \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}^{-1} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} = \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} = \text{LHS.} \end{aligned}$$

**Inductive step:** Assume the explicit formula holds up to and including some particular  $n$ , and consider the case  $n + 1$ . Then, using the implicit definition, preliminary results R1 and R2, and the inductive assumption,

$$\begin{aligned} \begin{pmatrix} x_{n+1} \\ y_{n+1} \end{pmatrix} &= \begin{pmatrix} 4 & -1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x_n \\ y_n \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 3 & 0 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_n \\ y_n \end{pmatrix} \\ &= \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 3 & 0 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}^{-1} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 3^n & 0 \\ 0 & 2^n \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 3^{n+1} & 0 \\ 0 & 2^{n+1} \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} \quad \text{and hence the formula} \\ &\quad \text{also holds for } n + 1. \end{aligned}$$

# Fibonacci Sequence

Recall the Fibonacci sequence  $(F_n)_{n \geq 0} = (1, 1, 2, 3, 5, 8, 13, 21, \dots)$

It has implicit formula,  $F_0 = 1, F_1 = 1$  and  $F_n = F_{n-1} + F_{n-2}, \forall n \geq 2$ .



# Fibonacci Sequence

Recall the Fibonacci sequence  $(F_n)_{n \geq 0} = (1, 1, 2, 3, 5, 8, 13, 21, \dots)$

It has implicit formula,  $F_0 = 1, F_1 = 1$  and  $F_n = F_{n-1} + F_{n-2}, \forall n \geq 2$ .

For all  $n \geq 1$  we have:

$$\begin{pmatrix} F_{n+1} \\ F_n \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} F_n \\ F_{n-1} \end{pmatrix}$$

# Fibonacci Sequence

Recall the Fibonacci sequence  $(F_n)_{n \geq 0} = (1, 1, 2, 3, 5, 8, 13, 21, \dots)$

It has implicit formula,  $F_0 = 1, F_1 = 1$  and  $F_n = F_{n-1} + F_{n-2}, \forall n \geq 2$ .

For all  $n \geq 1$  we have:

$$\begin{aligned} \begin{pmatrix} F_{n+1} \\ F_n \end{pmatrix} &= \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} F_n \\ F_{n-1} \end{pmatrix} \\ &= \frac{1}{\sqrt{5}} \begin{pmatrix} \varphi & -\varphi^{-1} \\ 1 & 1 \end{pmatrix} \begin{pmatrix} \varphi & 0 \\ 0 & -\varphi^{-1} \end{pmatrix} \begin{pmatrix} 1 & \varphi^{-1} \\ -1 & \varphi \end{pmatrix} \begin{pmatrix} F_n \\ F_{n-1} \end{pmatrix} \end{aligned}$$

# Fibonacci Sequence

Recall the Fibonacci sequence  $(F_n)_{n \geq 0} = (1, 1, 2, 3, 5, 8, 13, 21, \dots)$

It has implicit formula,  $F_0 = 1, F_1 = 1$  and  $F_n = F_{n-1} + F_{n-2}, \forall n \geq 2$ .

For all  $n \geq 1$  we have:

$$\begin{aligned}\begin{pmatrix} F_{n+1} \\ F_n \end{pmatrix} &= \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} F_n \\ F_{n-1} \end{pmatrix} \\ &= \frac{1}{\sqrt{5}} \begin{pmatrix} \varphi & -\varphi^{-1} \\ 1 & 1 \end{pmatrix} \begin{pmatrix} \varphi & 0 \\ 0 & -\varphi^{-1} \end{pmatrix} \begin{pmatrix} 1 & \varphi^{-1} \\ -1 & \varphi \end{pmatrix} \begin{pmatrix} F_n \\ F_{n-1} \end{pmatrix} \\ &= \frac{1}{\sqrt{5}} \begin{pmatrix} \varphi & -\varphi^{-1} \\ 1 & 1 \end{pmatrix} \begin{pmatrix} \varphi^n & 0 \\ 0 & -\varphi^{-n} \end{pmatrix} \begin{pmatrix} 1 & \varphi^{-1} \\ -1 & \varphi \end{pmatrix} \begin{pmatrix} F_1 \\ F_0 \end{pmatrix}\end{aligned}$$

where  $\varphi = \frac{1+\sqrt{5}}{2}$ . We can extract an explicit formula for the Fibonacci sequence!