

Discrete Mathematical Models

Lecture 21

Kane Townsend

Semester 2, 2024

C3: Markov Processes (cont.)

Markov Process Definition (Clearer)

A (discrete) **Markov process** is a system that has:

- a finite number k of states,
- a sequence of time steps $n \in \mathbb{N}^*$,
- probabilities of moving from a state to another state (including itself) that depends only on your current state.

Hence, probabilities of being in a particular state at time $n \geq 1$ depend on

- (i) its state at the $(n - 1)$ -th time step, and
- (ii) a fixed stochastic matrix $T \in M_k(\mathbb{Q}_+)$ called the **transition matrix** of the process.

Markov Process Definition (Clearer)

A (discrete) **Markov process** is a system that has:

- a finite number k of states,
- a sequence of time steps $n \in \mathbb{N}^*$,
- probabilities of moving from a state to another state (including itself) that depends only on your current state.

Hence, probabilities of being in a particular state at time $n \geq 1$ depend on

- (i) its state at the $(n - 1)$ -th time step, and
- (ii) a fixed stochastic matrix $T \in M_k(\mathbb{Q}_+)$ called the **transition matrix** of the process.

The (i, j) -entry T_{ij} of the transition matrix T specifies the probability that the system will be in the i -th state at any time step $n \geq 1$, given that it was in the j -th state at time step $n - 1$.

Markov Process Definition (Clearer)

A (discrete) **Markov process** is a system that has:

- a finite number k of states,
- a sequence of time steps $n \in \mathbb{N}^*$,
- probabilities of moving from a state to another state (including itself) that depends only on your current state.

Hence, probabilities of being in a particular state at time $n \geq 1$ depend on

- (i) its state at the $(n - 1)$ -th time step, and
- (ii) a fixed stochastic matrix $T \in M_k(\mathbb{Q}_+)$ called the **transition matrix** of the process.

The (i, j) -entry T_{ij} of the transition matrix T specifies the probability that the system will be in the i -th state at any time step $n \geq 1$, given that it was in the j -th state at time step $n - 1$.

A **transition diagram** is a complete weighted directed graph with k vertices representing the states of the system and the edge from the j -th vertex to the i -th vertex labelled with the probability T_{ij} .

Example 2

Another example: weather in 'Oz'

We investigate the weather on the island of Oz.

Another example: weather in 'Oz'

We investigate the weather on the island of Oz.

(This, it will be seen, is a rather simplified example, to illustrate the principles without too much heavy calculation.)

Another example: weather in 'Oz'

We investigate the weather on the island of Oz.

(This, it will be seen, is a rather simplified example, to illustrate the principles without too much heavy calculation.)

Weather in Oz is not great, e.g. it never has two fine days in a row.

Another example: weather in 'Oz'

We investigate the weather on the island of Oz.

(This, it will be seen, is a rather simplified example, to illustrate the principles without too much heavy calculation.)

Weather in Oz is not great, *e.g.* it never has two fine days in a row.

If the weather on a particular day is known, we cannot predict exactly what the weather will be the next day, but we can give probabilities of various kinds of weather as follows:

Another example: weather in 'Oz'

We investigate the weather on the island of Oz.

(This, it will be seen, is a rather simplified example, to illustrate the principles without too much heavy calculation.)

Weather in Oz is not great, *e.g.* it never has two fine days in a row.

If the weather on a particular day is known, we cannot predict exactly what the weather will be the next day, but we can give probabilities of various kinds of weather as follows:

There are only three kinds: fine (F), cloudy (C) and rain (R).

Another example: weather in 'Oz'

We investigate the weather on the island of Oz.

(This, it will be seen, is a rather simplified example, to illustrate the principles without too much heavy calculation.)

Weather in Oz is not great, e.g. it never has two fine days in a row.

If the weather on a particular day is known, we cannot predict exactly what the weather will be the next day, but we can give probabilities of various kinds of weather as follows:

There are only three kinds: fine (F), cloudy (C) and rain (R).

Here is the behaviour:

- After F, the weather is equally likely to be C or R.

Another example: weather in 'Oz'

We investigate the weather on the island of Oz.

(This, it will be seen, is a rather simplified example, to illustrate the principles without too much heavy calculation.)

Weather in Oz is not great, e.g. it never has two fine days in a row.

If the weather on a particular day is known, we cannot predict exactly what the weather will be the next day, but we can give probabilities of various kinds of weather as follows:

There are only three kinds: fine (F), cloudy (C) and rain (R).

Here is the behaviour:

- After F, the weather is equally likely to be C or R.
- After C, the probabilities are $\frac{1}{4}$ for F, $\frac{1}{4}$ for C and $\frac{1}{2}$ for R.

Another example: weather in 'Oz'

We investigate the weather on the island of Oz.

(This, it will be seen, is a rather simplified example, to illustrate the principles without too much heavy calculation.)

Weather in Oz is not great, e.g. it never has two fine days in a row.

If the weather on a particular day is known, we cannot predict exactly what the weather will be the next day, but we can give probabilities of various kinds of weather as follows:

There are only three kinds: fine (F), cloudy (C) and rain (R).

Here is the behaviour:

- After F, the weather is equally likely to be C or R.
- After C, the probabilities are $\frac{1}{4}$ for F, $\frac{1}{4}$ for C and $\frac{1}{2}$ for R.
- After R, the probabilities are $\frac{1}{4}$ For F, $\frac{1}{2}$ for C and $\frac{1}{4}$ for R.

Displaying the Oz weather system

Here are three ways to show the probabilities:

Displaying the Oz weather system

Here are three ways to show the probabilities:

As a table:

Probabilities of weather tomorrow are:

Given that the weather today is:	\curvearrowright	fine	cloudy	rain
	fine	0	$\frac{1}{2}$	$\frac{1}{2}$
	cloudy	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{2}$
	rain	$\frac{1}{4}$	$\frac{1}{2}$	$\frac{1}{4}$

Displaying the Oz weather system


Here are three ways to show the probabilities:

As a table: Probabilities of weather tomorrow are:

τ	fine	cloudy	rain
fine	0	$\frac{1}{2}$	$\frac{1}{2}$
cloudy	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{2}$
rain	$\frac{1}{4}$	$\frac{1}{2}$	$\frac{1}{4}$

Given
that the
weather
today is:

As a diagram:



Displaying the Oz weather system

Here are three ways to show the probabilities:


As a table: Probabilities of weather tomorrow are:

Given that the weather today is:

ω	fine	cloudy	rain
fine	0	$\frac{1}{2}$	$\frac{1}{2}$
cloudy	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{2}$
rain	$\frac{1}{4}$	$\frac{1}{2}$	$\frac{1}{4}$

As a matrix:

$$T = \begin{bmatrix} 0 & 1/4 & 1/4 \\ 1/2 & 1/4 & 1/2 \\ 1/2 & 1/2 & 1/4 \end{bmatrix}$$



Note: T is a stochastic matrix but not positive. However, since a power of T is positive, Perron-Frobenius still applies and we have a unique steady state vector and our guessing method still works.

Next day in Oz, via transition matrix

- Given probabilities on Day n , we can find probabilities on Day $n + 1$.

Next day in Oz, via transition matrix

- Given probabilities on Day n , we can find probabilities on Day $n + 1$.
- We need a state vector. Let us call the probabilities on day n
 - Prob x that it will be fine,
 - Prob y that it will be cloudy, and
 - Prob z that it will rain.

Next day in Oz, via transition matrix

- Given probabilities on Day n , we can find probabilities on Day $n + 1$.
- We need a state vector. Let us call the probabilities on day n
 - Prob x that it will be fine,
 - Prob y that it will be cloudy, and
 - Prob z that it will rain.

- Define $x_n = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$

Next day in Oz, via transition matrix

- Given probabilities on Day n , we can find probabilities on Day $n + 1$.
- We need a state vector. Let us call the probabilities on day n
 - Prob x that it will be fine,
 - Prob y that it will be cloudy, and
 - Prob z that it will rain.

- Define $x_n = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$

- Then, according to the Markov process theorem:

$$\begin{aligned} x_{n+1} &= T x_n \\ &= \begin{bmatrix} 0 & 1/4 & 1/4 \\ 1/2 & 1/4 & 1/2 \\ 1/2 & 1/2 & 1/4 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} (1/4)y + (1/4)z \\ (1/2)x + (1/4)y + (1/2)z \\ (1/2)x + (1/2)y + (1/4)z \end{bmatrix} \end{aligned}$$

Next week in Oz

Suppose we look outside
and see that today is fine.
Then we can say that

$$x_0 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}.$$

Next week in Oz

Suppose we look outside
and see that today is fine.
Then we can say that

$$x_0 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}.$$

That is, the probability of fine is 1, the probability of cloudy is 0 and the probability of rain is 0 today.

Next week in Oz

Suppose we look outside
and see that today is fine.
Then we can say that

$$x_0 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}.$$

That is, the probability of fine is 1, the probability of cloudy is 0 and the probability of rain is 0 today.

Then the probabilities
one week from today
(Day 7) are given by

$$x_7 = T^7 x_0.$$

Next week in Oz

Suppose we look outside
and see that today is fine.
Then we can say that

$$x_0 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}.$$

That is, the probability of fine is 1, the probability of cloudy is 0 and the probability of rain is 0 today.

Then the probabilities
one week from today
(Day 7) are given by

$$x_7 = T^7 x_0.$$

Using a computer to calculate the 7th power of the matrix, we get

$$x_7 = \begin{bmatrix} 0 & 1/4 & 1/4 \\ 1/2 & 1/4 & 1/2 \\ 1/2 & 1/2 & 1/4 \end{bmatrix}^7 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 819/4096 \\ 3277/8192 \\ 3277/8192 \end{bmatrix}.$$

Next week in Oz

Suppose we look outside
and see that today is fine.

Then we can say that

$$x_0 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}.$$

That is, the probability of fine is 1, the probability of cloudy is 0 and the probability of rain is 0 today.

Then the probabilities

one week from today

(Day 7) are given by

$$x_7 = T^7 x_0.$$

Using a computer to calculate the 7th power of the matrix, we get

$$x_7 = \begin{bmatrix} 0 & 1/4 & 1/4 \\ 1/2 & 1/4 & 1/2 \\ 1/2 & 1/2 & 1/4 \end{bmatrix}^7 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 819/4096 \\ 3277/8192 \\ 3277/8192 \end{bmatrix}.$$

Perhaps decimals would be more illuminating?

Days 1 through 10 in Oz

Computer calculations give:

$$x_1 = \begin{bmatrix} 0 \\ .5 \\ .5 \end{bmatrix}, \quad x_2 = \begin{bmatrix} .250 \\ .375 \\ .375 \end{bmatrix}, \quad x_3 = \begin{bmatrix} .18750 \\ .40625 \\ .40625 \end{bmatrix}, \quad x_4 = \begin{bmatrix} .2031250 \\ .3984375 \\ .3984375 \end{bmatrix},$$

$$x_5 = \begin{bmatrix} .199218750 \\ .400390625 \\ .400390625 \end{bmatrix}, \quad x_6 = \begin{bmatrix} .1999511719 \\ .4000244141 \\ .4000244141 \end{bmatrix}, \quad x_7 = \begin{bmatrix} .19995511719 \\ .4000244141 \\ .4000244141 \end{bmatrix},$$

$$x_8 = \begin{bmatrix} .2000122070 \\ .3999938965 \\ .3999938965 \end{bmatrix}, \quad x_9 = \begin{bmatrix} .1999969438 \\ .4000015260 \\ .4000015260 \end{bmatrix}, \quad x_{10} = \begin{bmatrix} .2000007629 \\ .3999996185 \\ .3999996185 \end{bmatrix}.$$

A steady state for the weather in Oz

These values seem to be
converging to a long-term
steady state of

$$v = \begin{bmatrix} .2 \\ .4 \\ .4 \end{bmatrix},$$

i.e. a probability of 0.2 of fine weather, a probability of 0.4 of cloudy weather and a probability of 0.4 of rainy weather.

A steady state for the weather in Oz

These values seem to be
converging to a long-term
steady state of

$$v = \begin{bmatrix} .2 \\ .4 \\ .4 \end{bmatrix},$$

i.e. a probability of 0.2 of fine weather, a probability of 0.4 of cloudy weather and a probability of 0.4 of rainy weather.

As they must, these probabilities sum to 1.

A steady state for the weather in Oz

These values seem to be converging to a long-term steady state of

$$v = \begin{bmatrix} .2 \\ .4 \\ .4 \end{bmatrix},$$

i.e. a probability of 0.2 of fine weather, a probability of 0.4 of cloudy weather and a probability of 0.4 of rainy weather.

As they must, these probabilities sum to 1.

To check that this v really is a normalised steady state vector, we also calculate

$$Tv = \begin{bmatrix} 0 & 0.25 & 0.25 \\ 0.50 & 0.25 & 0.50 \\ 0.50 & 0.50 & 0.25 \end{bmatrix} \begin{bmatrix} 0.2 \\ 0.4 \\ 0.4 \end{bmatrix} = \begin{bmatrix} 0.1 + 0.1 \\ 0.1 + 0.1 + 0.2 \\ 0.1 + 0.2 + 0.1 \end{bmatrix} = \begin{bmatrix} 0.2 \\ 0.4 \\ 0.4 \end{bmatrix}.$$

A steady state for the weather in Oz

These values seem to be converging to a long-term steady state of

$$v = \begin{bmatrix} .2 \\ .4 \\ .4 \end{bmatrix},$$

i.e. a probability of 0.2 of fine weather, a probability of 0.4 of cloudy weather and a probability of 0.4 of rainy weather.

As they must, these probabilities sum to 1.

To check that this v really is a normalised steady state vector, we also calculate

$$Tv = \begin{bmatrix} 0 & 0.25 & 0.25 \\ 0.50 & 0.25 & 0.50 \\ 0.50 & 0.50 & 0.25 \end{bmatrix} \begin{bmatrix} 0.2 \\ 0.4 \\ 0.4 \end{bmatrix} = \begin{bmatrix} 0.1 + 0.1 \\ 0.1 + 0.1 + 0.2 \\ 0.1 + 0.2 + 0.1 \end{bmatrix} = \begin{bmatrix} 0.2 \\ 0.4 \\ 0.4 \end{bmatrix}.$$

Therefore

$$Tv = v. \quad \checkmark$$

Methods for solving the matrix equation

There are several ways to solve the equation

$$Tv = v \iff (T - I)v = 0$$

for v and determine when v normalised.

Methods for solving the matrix equation

There are several ways to solve the equation

$$Tv = v \iff (T - I)v = 0$$

for v and determine when v normalised.

1. Gaussian (or Gauss-Jordan) elimination.

Methods for solving the matrix equation

There are several ways to solve the equation

$$Tv = v \iff (T - I)v = 0$$

for v and determine when v normalised.

1. Gaussian (or Gauss-Jordan) elimination. However,
Gaussian elimination is not taught nor assumed in this course
2. A shortcut method that then can be solved for 2×2 inverse matrices,
or for larger matrices we can use computer applications such as:
Matrix Reshish: <https://matrix.reshish.com>
MatrixCalc: <https://matrixcalc.org/en/>

Methods for solving the matrix equation

There are several ways to solve the equation

$$Tv = v \iff (T - I)v = 0$$

for v and determine when v normalised.

1. Gaussian (or Gauss-Jordan) elimination. However,
Gaussian elimination is not taught nor assumed in this course
2. A shortcut method that then can be solved for 2×2 inverse matrices,
or for larger matrices we can use computer applications such as:
Matrix Reshish: <https://matrix.reshish.com>
MatrixCalc: <https://matrixcalc.org/en/>
or, less conveniently but more robustly,
WolframAlpha: <https://www.wolframalpha.com/>

Methods for solving the matrix equation

There are several ways to solve the equation

$$Tv = v \iff (T - I)v = 0$$

for v and determine when v normalised.

1. Gaussian (or Gauss-Jordan) elimination. However,
Gaussian elimination is not taught nor assumed in this course
2. A shortcut method that then can be solved for 2×2 inverse matrices,
or for larger matrices we can use computer applications such as:
Matrix Reshish: <https://matrix.reshish.com>
MatrixCalc: <https://matrixcalc.org/en/>
or, less conveniently but more robustly,
WolframAlpha: <https://www.wolframalpha.com/>

A shortcut

A shortcut to this process is to take the augmented matrix $[T - I|0]$ as below (this is just representing $(T - I)v = 0$ in a more succinct way),

$$\left[\begin{array}{ccc|c} -1 & 1/4 & 1/4 & 0 \\ 1/2 & -3/4 & 1/2 & 0 \\ 1/2 & 1/2 & -3/4 & 0 \end{array} \right]$$

A shortcut

A shortcut to this process is to take the augmented matrix $[T - I|0]$ as below (this is just representing $(T - I)v = 0$ in a more succinct way),

$$\left[\begin{array}{ccc|c} -1 & 1/4 & 1/4 & 0 \\ 1/2 & -3/4 & 1/2 & 0 \\ 1/2 & 1/2 & -3/4 & 0 \end{array} \right]$$

throw away the last row and replace it with $[1 \dots 1|1]$, as in

$$\left[\begin{array}{ccc|c} -1 & 1/4 & 1/4 & 0 \\ 1/2 & -3/4 & 1/2 & 0 \\ 1 & 1 & 1 & 1 \end{array} \right]$$

A shortcut

A shortcut to this process is to take the augmented matrix $[T - I|0]$ as below (this is just representing $(T - I)v = 0$ in a more succinct way),

$$\left[\begin{array}{ccc|c} -1 & 1/4 & 1/4 & 0 \\ 1/2 & -3/4 & 1/2 & 0 \\ 1/2 & 1/2 & -3/4 & 0 \end{array} \right]$$

throw away the last row and replace it with $[1 \dots 1|1]$, as in

$$\left[\begin{array}{ccc|c} -1 & 1/4 & 1/4 & 0 \\ 1/2 & -3/4 & 1/2 & 0 \\ 1 & 1 & 1 & 1 \end{array} \right]$$

and solve this new system to directly obtain the unique solution for v .

Solving by Computer (using Reshish)

The system of equations

$$\left[\begin{array}{ccc|c} -1 & 1/4 & 1/4 & 0 \\ 1/2 & -3/4 & 1/2 & 0 \\ 1 & 1 & 1 & 1 \end{array} \right]$$

is entered into the

Reshish Matrix Calculator, using the “Gauss-Jordan Elimination” Tool :

Solving by Computer (using Reshish)

The system of equations

$$\left[\begin{array}{ccc|c} -1 & 1/4 & 1/4 & 0 \\ 1/2 & -3/4 & 1/2 & 0 \\ 1 & 1 & 1 & 1 \end{array} \right]$$

is entered into the

Reshish Matrix Calculator, using the “Gauss-Jordan Elimination” Tool :

Note that I have chosen to use “fractional” coefficients, to ensure an exact solution.

The screenshot shows the 'Matrix input' window of the Reshish Matrix Calculator. It features a title bar with a close button. Below the title bar are two buttons: 'Insert matrix' and 'Restore matrix'. A checkbox labeled 'Complex numbers (more)' is present. A dropdown menu is set to 'Fractional', and an information icon is visible. The main area contains a table with 3 rows and 5 columns. The columns are labeled X_1 , X_2 , X_3 , and b . The rows are numbered 1, 2, and 3. The values entered in the cells are: Row 1: -1, 1/4, 1/4, 0; Row 2: 1/2, -3/4, 1/2, 0; Row 3: 1, 1, 1, 1. At the bottom, there are buttons for 'Clear', 'Fill empty cells with zero', a checkbox for 'Very detailed solution', and a 'Solve' button.

	X_1	X_2	X_3	b
1	-1	1/4	1/4	0
2	1/2	-3/4	1/2	0
3	1	1	1	1

Solving by Computer (using Reshish)

The system of equations

$$\left[\begin{array}{ccc|c} -1 & 1/4 & 1/4 & 0 \\ 1/2 & -3/4 & 1/2 & 0 \\ 1 & 1 & 1 & 1 \end{array} \right]$$

is entered into the

Reshish Matrix Calculator, using the “Gauss-Jordan Elimination” Tool :

Note that I have chosen to use “fractional” coefficients, to ensure an exact solution.

The screenshot shows the 'Matrix input' window of the Reshish Matrix Calculator. It features a title bar with a close button. Below the title bar are two buttons: 'Insert matrix' and 'Restore matrix'. A checkbox labeled 'Complex numbers (more)' is present. A dropdown menu is set to 'Fractional', and an information icon is visible. The main area contains a table with 4 columns: an empty column, X_1 , X_2 , X_3 , and b . The rows are numbered 1, 2, and 3. The values entered are: Row 1: -1, 1/4, 1/4, 0; Row 2: 1/2, -3/4, 1/2, 0; Row 3: 1, 1, 1, 1. At the bottom, there are buttons for 'Clear', 'Fill empty cells with zero', a checkbox for 'Very detailed solution', and a 'Solve' button.

	X_1	X_2	X_3	b
1	-1	1/4	1/4	0
2	1/2	-3/4	1/2	0
3	1	1	1	1

Here is how Reshish responds:

Show solution

Solution set:

$$x_1 = 1/5$$

$$x_2 = 2/5$$

$$x_3 = 2/5$$

Back to the first example

We have seen that to find the steady state vector v for a Markov process with transition matrix T we need to solve the linear system that results from replacing the last equation in

$$(T - I)v = 0$$

by the equation that says that v is a probability vector.

For Cathy's employment process we had

$$T = \begin{bmatrix} 0.8 & 0.2 \\ 0.6 & 0.4 \end{bmatrix}$$

and, by a 'guess and check' method, we discovered that

$$v = \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0.75 \\ 0.25 \end{bmatrix}.$$

Solution by matrix inverse

Because T is 2×2 , and we have a formula for the inverse of a 2×2 matrix, we can find Cathy's steady state vector directly, without Gaussian elimination or computer. There are three steps:

1. Write out the matrix equation $(T - I)v = 0$:

$$\left(\begin{bmatrix} 0.8 & 0.6 \\ 0.2 & 0.4 \end{bmatrix} - \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

i.e.

$$\begin{bmatrix} -0.2 & 0.6 \\ 0.2 & -0.6 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

2. Replace the second equation by $x + y = 1$:

$$\begin{bmatrix} -0.2 & 0.6 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

Solution by matrix inverse (conclusion)

3. Solve this system using matrix inverse:

$$\begin{aligned}\begin{bmatrix} x \\ y \end{bmatrix} &= \begin{bmatrix} -0.2 & 0.6 \\ 1 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 0 \\ 1 \end{bmatrix} \\ &= \frac{1}{-0.2 - 0.6} \begin{bmatrix} 1 & -0.6 \\ -1 & -0.2 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} \\ &= \frac{1}{-0.8} \begin{bmatrix} -0.6 \\ -0.2 \end{bmatrix} = \begin{bmatrix} 6/8 \\ 2/8 \end{bmatrix} = \begin{bmatrix} 0.75 \\ 0.25 \end{bmatrix}\end{aligned}$$

Example 3

New Example — Colours of flowers

A species of flower (carnations say) has three colour varieties. The relevant genetics are as shown in the table:

Colour	Genotype
Red	RR
Pink	RW
White	WW

New Example — Colours of flowers

A species of flower (carnations say) has three colour varieties.

The relevant genetics are as shown in the table:

Colour	Genotype
Red	RR
Pink	RW
White	WW

At the nursery they are always crossed with the pink variety.
What will be the long term proportions of the three varieties?

New Example — Colours of flowers

A species of flower (carnations say) has three colour varieties.

The relevant genetics are as shown in the table:


Colour	Genotype
Red	RR
Pink	RW
White	WW

At the nursery they are always crossed with the pink variety.

What will be the long term proportions of the three varieties?

First, we need the transition probabilities.

Can you work them out?



New Example — Colours of flowers

A species of flower (carnations say) has three colour varieties. The relevant genetics are as shown in the table:


Colour	Genotype
Red	RR
Pink	RW
White	WW

At the nursery they are always crossed with the pink variety. What will be the long term proportions of the three varieties?

First, we need the transition probabilities. Can you work them out?

The transition matrix is

$$T = \begin{array}{cc} & \begin{array}{ccc} \text{Red} & \text{Pink} & \text{White} \end{array} \\ \begin{array}{c} \text{Red} \\ \text{Pink} \\ \text{White} \end{array} & \begin{bmatrix} 0.5 & 0.25 & 0 \\ 0.5 & 0.5 & 0.5 \\ 0 & 0.25 & 0.5 \end{bmatrix} \end{array} .$$



Finding the steady state

(a) $[T - I|0]$ is

$$\left[\begin{array}{ccc|c} -0.5 & 0.25 & 0 & 0 \\ 0.5 & -0.5 & 0.5 & 0 \\ 0 & 0.25 & -0.5 & 0 \end{array} \right]$$

(b) Replacing the bottom row with all 1's gives


$$\left[\begin{array}{ccc|c} -0.5 & 0.25 & 0 & 0 \\ 0.5 & -0.5 & 0.5 & 0 \\ 1 & 1 & 1 & 1 \end{array} \right]$$

Finding the steady state by computer

We can solve the system
using the computer.

Finding the steady state by computer

We can solve the system using the computer. For example, using Reshish:




The screenshot shows the Reshish software interface. At the top, there is a dropdown menu set to "Fractional" and an information icon. Below this is a table representing a linear system $AX = b$.

	X_1	X_2	X_3	b
1	$-1/2$	$1/4$	0	0
2	$1/2$	$-1/2$	$1/2$	0
3	1	1	1	1

Finding the steady state by computer

We can solve the system using the computer. For example, using Reshish:



The screenshot shows the Reshish software interface. At the top, there is a dropdown menu set to "Fractional" and an information icon. Below this is a table representing a linear system $AX = b$.


	X_1	X_2	X_3	b
1	$-1/2$	$1/4$	0	0
2	$1/2$	$-1/2$	$1/2$	0
3	1	1	1	1

Finding the steady state by computer

We can solve the system using the computer. For example, using Reshish:

Hence there is a unique steady state vector of

$$v = \begin{bmatrix} 1/4 \\ 1/2 \\ 1/4 \end{bmatrix}$$



The screenshot shows a software interface with a dark background. At the top, there is a dropdown menu labeled "Fractional" and a circular icon with an "i". Below these is a table with 4 columns and 4 rows. The columns are labeled x_1 , x_2 , x_3 , and b . The rows are labeled 1, 2, and 3. The values in the cells are as follows:

	x_1	x_2	x_3	b
1	-1/2	1/4	0	0
2	1/2	-1/2	1/2	0
3	1	1	1	1


Finding the steady state by computer

We can solve the system using the computer. For example, using Reshish:

Hence there is a unique steady state vector of

$$v = \begin{bmatrix} 1/4 \\ 1/2 \\ 1/4 \end{bmatrix}$$

So the species has a steady state in which 25% of the flowers are coloured red, 50% pink, and 25% white.



The screenshot shows a software interface with a dark background. At the top, there is a dropdown menu labeled 'Fractional' and a circular icon with the letter 'i'. Below these is a table representing a linear system. The table has four columns: an unlabeled column for row indices, and columns labeled x_1 , x_2 , x_3 , and b . The rows are indexed 1, 2, and 3.

	x_1	x_2	x_3	b
1	-1/2	1/4	0	0
2	1/2	-1/2	1/2	0
3	1	1	1	1

Checking the answer

Let's check $Tv = v$:

$$\begin{aligned}Tv &= \begin{bmatrix} 1/2 & 1/4 & 0 \\ 1/2 & 1/2 & 1/2 \\ 0 & 1/4 & 1/2 \end{bmatrix} \begin{bmatrix} 1/4 \\ 1/2 \\ 1/4 \end{bmatrix} \\&= \begin{bmatrix} 1/8 + 1/8 \\ 1/8 + 1/4 + 1/8 \\ 1/8 + 1/8 \end{bmatrix} \\&= \begin{bmatrix} 1/4 \\ 1/2 \\ 1/4 \end{bmatrix}\end{aligned}$$

Checking the answer

Let's check $Tv = v$:

$$\begin{aligned}Tv &= \begin{bmatrix} 1/2 & 1/4 & 0 \\ 1/2 & 1/2 & 1/2 \\ 0 & 1/4 & 1/2 \end{bmatrix} \begin{bmatrix} 1/4 \\ 1/2 \\ 1/4 \end{bmatrix} \\&= \begin{bmatrix} 1/8 + 1/8 \\ 1/8 + 1/4 + 1/8 \\ 1/8 + 1/8 \end{bmatrix} \\&= \begin{bmatrix} 1/4 \\ 1/2 \\ 1/4 \end{bmatrix}\end{aligned}$$

So yes, $Tv = v$. ✓

Will a Markov process always get to a steady state?

Will a Markov process always get to a steady state?

Not necessarily!

Example 4

Example: chemical compounds in transition

Consider a chemical compound whose molecule can exist in any one of five states, termed A , B , C , D and E .

Example: chemical compounds in transition


Consider a chemical compound whose molecule can exist in any one of five states, termed A , B , C , D and E .

Each molecule frequently undergoes transitions from one state to another, always to an 'adjacent' state, according to the probabilities shown in the transition diagram.

Example: chemical compounds in transition

Consider a chemical compound whose molecule can exist in any one of five states, termed A , B , C , D and E .


Each molecule frequently undergoes transitions from one state to another, always to an 'adjacent' state, according to the probabilities shown in the transition diagram.



Example: chemical compounds in transition

Consider a chemical compound whose molecule can exist in any one of five states, termed A, B, C, D and E .

Each molecule frequently undergoes transitions from one state to another, always to an 'adjacent' state, according to the probabilities shown in the transition diagram.



The transition matrix for this Markov Process is

$$T = \begin{bmatrix} 0 & 0.6 & 0 & 0 & 0 \\ 1 & 0 & 0.6 & 0 & 0 \\ 0 & 0.4 & 0 & 0.6 & 0 \\ 0 & 0 & 0.4 & 0 & 1 \\ 0 & 0 & 0 & 0.4 & 0 \end{bmatrix}$$

A beaker full of chemical

- Now suppose we have a beaker full of this chemical. We expect the proportions of A , B , C , D , E to relate to transition probabilities.

A beaker full of chemical

- Now suppose we have a beaker full of this chemical. We expect the proportions of A , B , C , D , E to relate to transition probabilities.
- What proportions of the compound will be in the various states?

A beaker full of chemical

- Now suppose we have a beaker full of this chemical. We expect the proportions of A , B , C , D , E to relate to transition probabilities.
- What proportions of the compound will be in the various states?
- To do a thorough analysis of all possible behaviours of this Markov Process, you need to study '*eigenvalues and eigenvectors*' – a reason to take a course or read a book on [Linear Algebra](#).

A beaker full of chemical

- Now suppose we have a beaker full of this chemical. We expect the proportions of A , B , C , D , E to relate to transition probabilities.
- What proportions of the compound will be in the various states?
- To do a thorough analysis of all possible behaviours of this Markov Process, you need to study '*eigenvalues and eigenvectors*' – a reason to take a course or read a book on [Linear Algebra](#).
- But let's see what we can figure out without those tools.

Chemical example — investigating with a computer

Suppose the beaker only contains form 'A' to start with, *i.e.*

$x_0 = [1, 0, 0, 0, 0]'$. Then by computer to 6dp we find:

$$\begin{aligned}x_{100} &= T^{100}x_0 \\&= [0.415383, 0.000000, 0.461538, 0.000000, 0.123077]^t \\x_{101} &= Tx_{100} \\&= [0.000000, 0.692308, 0.000000, 0.307692, 0.000000]^t\end{aligned}$$

Chemical example — investigating with a computer

Suppose the beaker only contains form 'A' to start with, *i.e.*

$x_0 = [1, 0, 0, 0, 0]'$. Then by computer to 6dp we find:

$$\begin{aligned}x_{100} &= T^{100}x_0 \\&= [0.415383, 0.000000, 0.461538, 0.000000, 0.123077]^t\end{aligned}$$

$$\begin{aligned}x_{101} &= Tx_{100} \\&= [0.000000, 0.692308, 0.000000, 0.307692, 0.000000]^t\end{aligned}$$

and continuing in the same manner

$$x_{102} = [0.415383, 0.000000, 0.461538, 0.000000, 0.123077]^t$$

$$x_{103} = [0.000000, 0.692308, 0.000000, 0.307692, 0.000000]^t$$

$$x_{104} = [0.415383, 0.000000, 0.461538, 0.000000, 0.123077]^t$$

$$x_{105} = [0.000000, 0.692308, 0.000000, 0.307692, 0.000000]^t$$

\vdots

\vdots

Chemical example — investigating with a computer

Suppose the beaker only contains form 'A' to start with, *i.e.*

$x_0 = [1, 0, 0, 0, 0]'$. Then by computer to 6dp we find:

$$\begin{aligned}x_{100} &= T^{100}x_0 \\&= [0.415383, 0.000000, 0.461538, 0.000000, 0.123077]^t\end{aligned}$$

$$\begin{aligned}x_{101} &= Tx_{100} \\&= [0.000000, 0.692308, 0.000000, 0.307692, 0.000000]^t\end{aligned}$$

and continuing in the same manner

$$x_{102} = [0.415383, 0.000000, 0.461538, 0.000000, 0.123077]^t$$

$$x_{103} = [0.000000, 0.692308, 0.000000, 0.307692, 0.000000]^t$$

$$x_{104} = [0.415383, 0.000000, 0.461538, 0.000000, 0.123077]^t$$

$$x_{105} = [0.000000, 0.692308, 0.000000, 0.307692, 0.000000]^t$$

\vdots

\vdots

It appears to alternate!

However starting with a beaker half full of A and half of B, *i.e.*

$x_0 = [0.5, 0.5, 0, 0, 0]^t$, and again using formulae

$$x_n = T^n x_0 \quad \text{and} \quad x_{n+1} = T x_n$$

repeatedly we get

$$x_{100} = [0.207692, 0.346154, 0.230769, 0.153846, 0.061539]^t$$

$$x_{101} = [0.207692, 0.346154, 0.230769, 0.153846, 0.061539]^t$$

$$x_{102} = [0.207692, 0.346154, 0.230769, 0.153846, 0.061539]^t$$

$$\vdots$$
$$\vdots$$

This looks like a steady state!

However starting with a beaker half full of A and half of B, *i.e.*

$x_0 = [0.5, 0.5, 0, 0, 0]^t$, and again using formulae

$$x_n = T^n x_0 \quad \text{and} \quad x_{n+1} = T x_n$$

repeatedly we get

$$x_{100} = [0.207692, 0.346154, 0.230769, 0.153846, 0.061539]^t$$

$$x_{101} = [0.207692, 0.346154, 0.230769, 0.153846, 0.061539]^t$$

$$x_{102} = [0.207692, 0.346154, 0.230769, 0.153846, 0.061539]^t$$

$$\vdots$$
$$\vdots$$

This looks like a steady state!

So this Markov Process is different to those we used to model employment, weather in Oz, and flower-colours because

eventual behaviour depends on where you start!

Steady state(s) for a beaker of chemical?

We can solve for the steady state to find out if it is unique.

Steady state(s) for a beaker of chemical?

We can solve for the steady state to find out if it is unique.

We need to solve

$$Tv = v$$

for $v = [x_1, x_2, x_3, x_4, x_5]^t$ subject to additional constraint

$$x_1 + x_2 + x_3 + x_4 + x_5 = 1.$$

Steady state(s) for a beaker of chemical?

We can solve for the steady state to find out if it is unique.

We need to solve

$$Tv = v$$

for $v = [x_1, x_2, x_3, x_4, x_5]^t$ subject to additional constraint

$$x_1 + x_2 + x_3 + x_4 + x_5 = 1.$$

We use the 'shortcut' method:

Steady state(s) for a beaker of chemical?

We can solve for the steady state to find out if it is unique.

We need to solve

$$Tv = v$$

for $v = [x_1, x_2, x_3, x_4, x_5]^t$ subject to additional constraint

$$x_1 + x_2 + x_3 + x_4 + x_5 = 1.$$

We use the 'shortcut' method:

(a) First construct $[T - I|0]$.

Steady state(s) for a beaker of chemical?

We can solve for the steady state to find out if it is unique.

We need to solve

$$Tv = v$$

for $v = [x_1, x_2, x_3, x_4, x_5]^t$ subject to additional constraint

$$x_1 + x_2 + x_3 + x_4 + x_5 = 1.$$

We use the 'shortcut' method:

- (a) First construct $[T - I|0]$.
- (b) Then replace the last row with all 1's.
- (c) Then solve by computer (Gaussian elimination).

Steady state(s) for a beaker of chemical?

(a) $[T - I \mid 0]$ is

$$\left[\begin{array}{ccccc|c} -1 & 0.6 & 0 & 0 & 0 & 0 \\ 1 & -1 & 0.6 & 0 & 0 & 0 \\ 0 & 0.4 & -1 & 0.6 & 0 & 0 \\ 0 & 0 & 0.4 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0.4 & -1 & 0 \end{array} \right]$$

(b) Replace the last row with all 1's


$$\left[\begin{array}{ccccc|c} -1 & 0.6 & 0 & 0 & 0 & 0 \\ 1 & -1 & 0.6 & 0 & 0 & 0 \\ 0 & 0.4 & -1 & 0.6 & 0 & 0 \\ 0 & 0 & 0.4 & -1 & 1 & 0 \\ 1 & 1 & 1 & 1 & 1 & 1 \end{array} \right]$$

Steady state for a beaker of chemical - by computer

Decimal 


	X_1	X_2	X_3	X_4	X_5	b
1	-1	0.6	0	0	0	0
2	1	-1	0.6	0	0	0
3	0	0.4	-1	0.4	0	0
4	0	0	0.4	-1	1	0
5	1	1	1	1	1	1

Steady state for a beaker of chemical - by computer

Decimal 

	X_1	X_2	X_3	X_4	X_5	b
1	-1	0.6	0	0	0	0
2	1	-1	0.6	0	0	0
3	0	0.4	-1	0.4	0	0
4	0	0	0.4	-1	1	0
5	1	1	1	1	1	1

Steady state for a beaker of chemical - by computer



A screenshot of a computer interface for solving a linear system. At the top, there is a 'Decimal' dropdown menu and a help icon. Below is a table with 7 columns: an index column, and columns for variables x_1 through x_5 , and a column for the vector b . The table contains 5 rows of data.

	x_1	x_2	x_3	x_4	x_5	b
1	-1	0.6	0	0	0	0
2	1	-1	0.6	0	0	0
3	0	0.4	-1	0.4	0	0
4	0	0	0.4	-1	1	0
5	1	1	1	1	1	1

This confirms there is a **unique** normalised steady state solution:

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = \begin{bmatrix} 27/130 \\ 45/130 \\ 15/65 \\ 10/65 \\ 4/65 \end{bmatrix} = \begin{bmatrix} 0.2077 \\ 0.3462 \\ 0.2308 \\ 0.1538 \\ 0.0615 \end{bmatrix}.$$

Steady state for a beaker of chemical - by computer

	x_1	x_2	x_3	x_4	x_5	b
1	-1	0.6	0	0	0	0
2	1	-1	0.6	0	0	0
3	0	0.4	-1	0.4	0	0
4	0	0	0.4	-1	1	0
5	1	1	1	1	1	1

This confirms there is a **unique** normalised steady state solution:

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = \begin{bmatrix} 27/130 \\ 45/130 \\ 15/65 \\ 10/65 \\ 4/65 \end{bmatrix} = \begin{bmatrix} 0.2077 \\ 0.3462 \\ 0.2308 \\ 0.1538 \\ 0.0615 \end{bmatrix}.$$

So the steady-state proportions of the five forms of the chemical are:

A: 20.77%, B: 34.62%,
C: 23.08%, D: 15.38%,
E: 6.15%.

A steady state for a beaker of chemical - conclusion

We found that **provided** the beaker reaches a **steady-state**, then proportions of the various forms of the chemical remain stable at

A : 20.77%, *B* : 34.62%, *C* : 23.08%, *D* : 15.38%, *E* : 6.15%.

A steady state for a beaker of chemical - conclusion

We found that **provided** the beaker reaches a **steady-state**, then proportions of the various forms of the chemical remain stable at

A : 20.77%, *B* : 34.62%, *C* : 23.08%, *D* : 15.38%, *E* : 6.15%.

Individual molecules DO change their form, but at a rate such that overall, proportions remain as above.

A steady state for a beaker of chemical - conclusion

We found that **provided** the beaker reaches a **steady-state**, then proportions of the various forms of the chemical remain stable at

A : 20.77%, *B* : 34.62%, *C* : 23.08%, *D* : 15.38%, *E* : 6.15%.

Individual molecules DO change their form, but at a rate such that overall, proportions remain as above.

END OF SECTION C3