

Discrete Mathematical Models

Lecture 17

Kane Townsend

Semester 2, 2024

Unordered Selection (cont.)

Stars and bars

[Multisets (Stars and Bars)] There are $\binom{r+n-1}{r}$ size- r multisets with members from a set of size n . That is, there are $\binom{r+n-1}{r}$ ways to arrange a list of r stars and $n - 1$ bars.

The connection between these two statements is definitely not obvious!

Stars and bars

[Multisets (Stars and Bars)] There are $\binom{r+n-1}{r}$ size- r multisets with members from a set of size n . That is, there are $\binom{r+n-1}{r}$ ways to arrange a list of r stars and $n - 1$ bars.

The connection between these two statements is definitely not obvious!

How does one think about multisets as an arrangement of stars and bars?

Firstly, we explain the second sentence

Stars and bars

[Multisets (Stars and Bars)] There are $\binom{r+n-1}{r}$ size- r multisets with members from a set of size n . That is, there are $\binom{r+n-1}{r}$ ways to arrange a list of r stars and $n - 1$ bars.

The connection between these two statements is definitely not obvious!

How does one think about multisets as an arrangement of stars and bars?

Firstly, we explain the second sentence

If the stars and bars were all distinct then there would be $(r + n - 1)!$ ways to permute them. However, we are overcounting, since there are r indistinguishable stars and $(n-1)$ indistinguishable bars the total number of arrangements is

$$\frac{(r + n - 1)!}{r! \times (n - 1)!}$$

This is exactly $\binom{r+n-1}{r}$.

Counting 'Multisets'

How many different size- r multisets can be formed from members of a set S of cardinality n ?

Counting 'Multisets'

How many different size- r multisets can be formed from members of a set S of cardinality n ?

Represent each multiset as a pattern of r stars and $n - 1$ bars.

Counting 'Multisets'

How many different size- r multisets can be formed from members of a set S of cardinality n ?

Represent each multiset as a pattern of r stars and $n - 1$ bars.

For example if $S = \{a, b, c, d\}$ and $m : S \rightarrow \mathbb{N}^*$, where $m(a) = 2, m(b) = 1, m(c) = 2, m(d) = 0$ then the multiset less formally written is

$$\{c, b, a, c, a\}$$

can be represented by

$$\star \star \mid \star \mid \star \star \mid$$

Counting 'Multisets'

How many different size- r multisets can be formed from members of a set S of cardinality n ?

Represent each multiset as a pattern of r stars and $n - 1$ bars.

For example if $S = \{a, b, c, d\}$ and $m : S \rightarrow \mathbb{N}^*$, where $m(a)=2, m(b)=1, m(c)=2, m(d)=0$ then the multiset less formally written is

$$\{c, b, a, c, a\}$$

can be represented by

$$\star\star \mid \star \mid \star\star \mid$$

There are $\binom{r+n-1}{r}$ size- r multisets with members from a set of size n .

Multisets example

If n is a positive integer, how many 5-tuples of integers from 1 through n can be formed in which the elements of the 5-tuple are written in decreasing order?

Multisets example

If n is a positive integer, how many 5-tuples of integers from 1 through n can be formed in which the elements of the 5-tuple are written in decreasing order?

For $n = 9$ some 5-tuples are $(8, 6, 4, 2, 1)$, $(9, 3, 3, 2, 2)$, $(6, 6, 6, 6, 6)$.

Multisets example

If n is a positive integer, how many 5-tuples of integers from 1 through n can be formed in which the elements of the 5-tuple are written in decreasing order?

For $n = 9$ some 5-tuples are $(8, 6, 4, 2, 1)$, $(9, 3, 3, 2, 2)$, $(6, 6, 6, 6, 6)$.

There is a bijection (one-to-one correspondence) between the set of all these 5-tuples and the set of all size-5 multisets chosen from $\{1, \dots, n\}$, because the r 'members' of the multiset can only be arranged in one way in decreasing order (there are no other arrangements).

Multisets example

If n is a positive integer, how many 5-tuples of integers from 1 through n can be formed in which the elements of the 5-tuple are written in decreasing order?

For $n = 9$ some 5-tuples are $(8, 6, 4, 2, 1)$, $(9, 3, 3, 2, 2)$, $(6, 6, 6, 6, 6)$.

There is a bijection (one-to-one correspondence) between the set of all these 5-tuples and the set of all size-5 multisets chosen from $\{1, \dots, n\}$, because the r 'members' of the multiset can only be arranged in one way in decreasing order (there are no other arrangements).

So by stars-and-bars, there are $\binom{5+n-1}{5} = \binom{n+4}{5}$ of these 5-tuples.

Multisets example

If n is a positive integer, how many 5-tuples of integers from 1 through n can be formed in which the elements of the 5-tuple are written in decreasing order?

For $n = 9$ some 5-tuples are $(8, 6, 4, 2, 1)$, $(9, 3, 3, 2, 2)$, $(6, 6, 6, 6, 6)$.

There is a bijection (one-to-one correspondence) between the set of all these 5-tuples and the set of all size-5 multisets chosen from $\{1, \dots, n\}$, because the r 'members' of the multiset can only be arranged in one way in decreasing order (there are no other arrangements).

So by stars-and-bars, there are $\binom{5+n-1}{5} = \binom{n+4}{5}$ of these 5-tuples.

For example for $n = 3$ there are $\binom{7}{5} = \binom{7}{2} = 21$ such 5-tuples:

33333 33332 33331 33322 33321 33311 33222 33221 33211 33111
32222 32221 32211 32111 31111 22222 22221 22211 22111 21111 11111

Another 'Stars and Bars' example

For how many integers from 1 through 99 999 is the sum of their digits equal to 10?

Another 'Stars and Bars' example

For how many integers from 1 through 99 999 is the sum of their digits equal to 10?

By inserting leading zeros if necessary, all these integers can be considered as 5-digit strings $abcde$ with $a+b+c+d+e=10$.

Another 'Stars and Bars' example

For how many integers from 1 through 99 999 is the sum of their digits equal to 10?

By inserting leading zeros if necessary, all these integers can be considered as 5-digit strings $abcde$ with $a+b+c+d+e=10$.

Each of these 5-digit strings can be represented as a length-14 pattern of 10 **stars** and 4 **bars**.

Another 'Stars and Bars' example

For how many integers from 1 through 99 999 is the sum of their digits equal to 10?

By inserting leading zeros if necessary, all these integers can be considered as 5-digit strings $abcde$ with $a+b+c+d+e=10$.

Each of these 5-digit strings can be represented as a length-14 pattern of 10 **stars** and 4 **bars**. For example:

$☆☆|☆☆☆|☆|☆☆|☆$ represents 24121.

Another 'Stars and Bars' example

For how many integers from 1 through 99 999 is the sum of their digits equal to 10?

By inserting leading zeros if necessary, all these integers can be considered as 5-digit strings $abcde$ with $a+b+c+d+e=10$.

Each of these 5-digit strings can be represented as a length-14 pattern of 10 **stars** and 4 **bars**. For example:

$\star\star | \star\star\star\star | \star | \star\star | \star$ represents 24 121.

$\star\star\star || \star\star\star\star | \star\star\star |$ represents 30 430.

Another 'Stars and Bars' example

For how many integers from 1 through 99 999 is the sum of their digits equal to 10?

By inserting leading zeros if necessary, all these integers can be considered as 5-digit strings $abcde$ with $a+b+c+d+e=10$.

Each of these 5-digit strings can be represented as a length-14 pattern of 10 **stars** and 4 **bars**. For example:

★★ ★★★★★ ★ ★★ ★	represents	24 121.
★★★ ★★★★★ ★★★	represents	30 430.
★★★★★★★★ ★	represents	00 091.

Another 'Stars and Bars' example

For how many integers from 1 through 99 999 is the sum of their digits equal to 10?

By inserting leading zeros if necessary, all these integers can be considered as 5-digit strings $abcde$ with $a+b+c+d+e=10$.

Each of these 5-digit strings can be represented as a length-14 pattern of 10 **stars** and 4 **bars**. For example:

$☆☆ ☆☆☆ ☆ ☆☆ ☆$	represents	24 121.
$☆☆☆ ☆☆☆ ☆☆☆ $	represents	30 430.
$ ☆☆☆☆☆☆☆ ☆$	represents	00 091.

There are $\binom{14}{4}$ of these 10-star-4-bar patterns.

Another 'Stars and Bars' example

For how many integers from 1 through 99 999 is the sum of their digits equal to 10?

By inserting leading zeros if necessary, all these integers can be considered as 5-digit strings $abcde$ with $a+b+c+d+e=10$.

Each of these 5-digit strings can be represented as a length-14 pattern of 10 **stars** and 4 **bars**. For example:

$☆☆ ☆☆☆ ☆ ☆☆ ☆$	represents	24 121.
$☆☆☆ ☆☆☆ ☆☆☆ $	represents	30 430.
$ ☆☆☆☆☆☆☆ ☆$	represents	00 091.

There are $\binom{14}{4}$ of these 10-star-4-bar patterns. But five of the patterns have ten stars in a row and so don't count (ten is not a digit).

Another 'Stars and Bars' example

For how many integers from 1 through 99 999 is the sum of their digits equal to 10?

By inserting leading zeros if necessary, all these integers can be considered as 5-digit strings $abcde$ with $a+b+c+d+e=10$.

Each of these 5-digit strings can be represented as a length-14 pattern of 10 **stars** and 4 **bars**. For example:

$\star\star | \star\star\star\star | \star | \star\star | \star$ represents 24 121.

$\star\star\star || \star\star\star\star | \star\star\star |$ represents 30 430.

$||| \star\star\star\star\star\star\star\star | \star$ represents 00 091.

There are $\binom{14}{4}$ of these 10-star-4-bar patterns. But five of the patterns have ten stars in a row and so don't count (ten is not a digit).

So the number of integers is $\binom{14}{4} - 5 = \frac{14 \times 13 \times 12 \times 11}{4 \times 3 \times 2 \times 1} - 5 = 996$.

Selecting summary

Select r from n	With repetition	Without repetition
Ordered	n^r	$\frac{n!}{(n-r)!}$
Unordered	$\binom{r+n-1}{r}$	$\binom{n}{r}$

Warning: Questions are often harder and more subtle than just applying these rules directly.

Question: What are the set and function interpretations of each?

Bit-String Example of Inclusion-Exclusion

How many bit-strings of length 8 can be constructed that start with '1'
OR end with '00'?

Bit-String Example of Inclusion-Exclusion

How many bit-strings of length 8 can be constructed that start with '1'
OR end with '00'?

Task 1: Construct a string of length 8 that starts with '1'.

Bit-String Example of Inclusion-Exclusion

How many bit-strings of length 8 can be constructed that start with '1'
OR end with '00'?

Task 1: Construct a string of length 8 that starts with '1'.

- There is one way to choose the first bit (1)
- There are two ways to choose the second bit (0 or 1)
- There are two ways to choose the third bit (0 or 1)
- \vdots
- There are two ways to choose the eighth bit (0 or 1)

Bit-String Example of Inclusion-Exclusion

How many bit-strings of length 8 can be constructed that start with '1'
OR end with '00'?

Task 1: Construct a string of length 8 that starts with '1'.

- There is one way to choose the first bit (1)
- There are two ways to choose the second bit (0 or 1)
- There are two ways to choose the third bit (0 or 1)
- \vdots
- There are two ways to choose the eighth bit (0 or 1)

Product Rule: Task 1 can be done in $1 \times 2^7 = 128$ ways.

Bit-String Example of Inclusion-Exclusion

Task 2: Construct a string of length 8 that ends with '00'.

Bit-String Example of Inclusion-Exclusion

Task 2: Construct a string of length 8 that ends with '00'.

- There are two ways to choose the first bit (0 or 1)
- There are two ways to choose the second bit (0 or 1)
- \vdots
- There are two ways to choose the sixth bit (0 or 1)
- There is one way to choose the seventh bit (0)
- There is one way to choose the eighth bit (0)

Bit-String Example of Inclusion-Exclusion

Task 2: Construct a string of length 8 that ends with '00'.

- There are two ways to choose the first bit (0 or 1)
- There are two ways to choose the second bit (0 or 1)
- \vdots
- There are two ways to choose the sixth bit (0 or 1)
- There is one way to choose the seventh bit (0)
- There is one way to choose the eighth bit (0)

Product Rule: Task 2 can be done in $2^6 \times 1^2 = 64$ ways.

Bit-String Example of Inclusion-Exclusion

Task: How many bit-strings of length 8 can be constructed that start with '1' OR end with '00'?

Is the answer $128 + 64 = 196$?

Bit-String Example of Inclusion-Exclusion

Task: How many bit-strings of length 8 can be constructed that start with '1' OR end with '00'?

Is the answer $128 + 64 = 196$? **NO!**

Bit-String Example of Inclusion-Exclusion

Task: How many bit-strings of length 8 can be constructed that start with '1' OR end with '00'?

Is the answer $128 + 64 = 196$? **NO!**

Bit-String Example of Inclusion-Exclusion

Task: How many bit-strings of length 8 can be constructed that start with '1' OR end with '00'?

Is the answer $128 + 64 = 196$? **NO!**

Task 3: Construct a string of length 8 that both starts with '1' and ends with '00'.

Bit-String Example of Inclusion-Exclusion

Task: How many bit-strings of length 8 can be constructed that start with '1' OR end with '00'?

Is the answer $128+64 = 196$? **NO!**

Task 3: Construct a string of length 8 that both starts with '1' and ends with '00'.

- There is one way to choose the first bit (1)
- There are two ways to choose each of the the second to sixth bit (0 or 1)
- There are two ways to choose the sixth bit (0 or 1)
- There is one way to choose the seventh bit (0)
- There is one way to choose the eighth bit (0)

Bit-String Example of Inclusion-Exclusion

Task: How many bit-strings of length 8 can be constructed that start with '1' OR end with '00'?

Is the answer $128+64 = 196$? **NO!**

Task 3: Construct a string of length 8 that both starts with '1' and ends with '00'.

- There is one way to choose the first bit (1)
- There are two ways to choose each of the the second to sixth bit (0 or 1)
- There are two ways to choose the sixth bit (0 or 1)
- There is one way to choose the seventh bit (0)
- There is one way to choose the eighth bit (0)

Product Rule: Task 3 can be done in $1 \times 2^5 \times 1^2 = 32$ ways.

Bit-String Example of Inclusion-Exclusion

Finally, the number of ways to construct a bit string of length 8 that starts with '1' or ends with '00' is equal to:

No. ways to do Task 1 + No. ways to do Task 2 – No. ways to do Task 3

That is:

$$128 + 64 - 32 = 160$$

Note: An alternative, and quite different, way to solve this problem is to use **complementary counting**;

Bit-String Example of Inclusion-Exclusion

Finally, the number of ways to construct a bit string of length 8 that starts with '1' or ends with '00' is equal to:

No. ways to do Task 1 + No. ways to do Task 2 – No. ways to do Task 3

That is:

$$128 + 64 - 32 = 160$$

Note: An alternative, and quite different, way to solve this problem is to use **complementary counting**; *i.e.* calculate $|S^c|$ where S is the set of strings we are interested in and U is the set of all (8-bit) strings.

Bit-String Example of Inclusion-Exclusion

Finally, the number of ways to construct a bit string of length 8 that starts with '1' or ends with '00' is equal to:

No. ways to do Task 1 + No. ways to do Task 2 – No. ways to do Task 3

That is:

$$128 + 64 - 32 = 160$$

Note: An alternative, and quite different, way to solve this problem is to use **complementary counting**; *i.e.* calculate $|S^c|$ where S is the set of strings we are interested in and U is the set of all (8-bit) strings. Then $|S| = |U| - |S^c| = 2^8 - 1 \times 2^5 \times 3 = 256 - 96 = 160$.

Bit-String Example of Inclusion-Exclusion

Finally, the number of ways to construct a bit string of length 8 that starts with '1' or ends with '00' is equal to:

No. ways to do Task 1 + No. ways to do Task 2 – No. ways to do Task 3

That is:

$$128 + 64 - 32 = 160$$

Note: An alternative, and quite different, way to solve this problem is to use **complementary counting**; i.e. calculate $|S^c|$ where S is the set of strings we are interested in and U is the set of all (8-bit) strings. Then $|S| = |U| - |S^c| = 2^8 - 1 \times 2^5 \times 3 = 256 - 96 = 160$.

Can you see how to get the $1 \times 2^5 \times 3$?

C2: Probability

Equal likelihood models

A thought experiment:



- Toss a coin.
- What are the possible outcomes?
- 'Heads' or 'Tails'
- What is the probability of 'Heads'?
- We say it is

$$\mathbb{P}(\text{Heads}) = \frac{1}{2}. \quad \text{Why?}$$

Methods of assigning probabilities

Method 1: Use relative frequencies

Method 1: Use *relative frequencies* (empirical experiment)

Methods of assigning probabilities

Method 1: Use **relative frequencies** (empirical experiment)

Method 2: Use a **model**

Methods of assigning probabilities

Method 1: Use **relative frequencies** (empirical experiment)

Method 2: Use a **model** (combination of prior knowledge, guessing, deduction)

- Eg. assume **equally likely outcomes**

Relative Frequencies

- For example, one may carry out the experiment of tossing a coin ten times and noting the results.

Relative Frequencies

- For example, one may carry out the experiment of tossing a coin ten times and noting the results.
- My experiment gave

T,T,H,H,T,T,T,H,T,H

Relative Frequencies

- For example, one may carry out the experiment of tossing a coin ten times and noting the results.
- My experiment gave

T,T,H,H,T,T,T,H,T,H

- Slightly fewer than half the coin-tosses resulted in 'H' (for 'Heads').

Relative Frequencies

- For example, one may carry out the experiment of tossing a coin ten times and noting the results.
- My experiment gave

T,T,H,H,T,T,T,H,T,H

- Slightly fewer than half the coin-tosses resulted in 'H' (for 'Heads').
- A 'longer run' may give different (better?) results.

Relative Frequencies

- For example, one may carry out the experiment of tossing a coin ten times and noting the results.
- My experiment gave

T,T,H,H,T,T,T,H,T,H

- Slightly fewer than half the coin-tosses resulted in 'H' (for 'Heads').
- A 'longer run' may give different (better?) results.
- There is much more to be said on 'relative frequencies', but for this course we will focus on making 'models'.

A model for coin tossing

- Observe a real coin: it has two sides – ‘Heads’ and ‘Tails’.

A model for coin tossing

- Observe a real coin: it has two sides – ‘Heads’ and ‘Tails’.
- Actually there is a third ‘side’: the rim.

A model for coin tossing

- Observe a real coin: it has two sides – ‘Heads’ and ‘Tails’.
- Actually there is a third ‘side’: the rim.
- Because the rim is small we **decide to ignore** this possibility.

A model for coin tossing

- Observe a real coin: it has two sides – ‘Heads’ and ‘Tails’.
- Actually there is a third ‘side’: the rim.
- Because the rim is small we **decide to ignore** this possibility.
- We **simplify** on purpose, to make the model *tractable*.

A model for coin tossing

- Observe a real coin: it has two sides – ‘Heads’ and ‘Tails’.
- Actually there is a third ‘side’: the rim.
- Because the rim is small we **decide to ignore** this possibility.
- We **simplify** on purpose, to make the model *tractable*.
- The ‘Heads’ and ‘Tails’ sides are so similar physically that we make an assumption:

A model for coin tossing

- Observe a real coin: it has two sides – ‘Heads’ and ‘Tails’.
- Actually there is a third ‘side’: the rim.
- Because the rim is small we **decide to ignore** this possibility.
- We **simplify** on purpose, to make the model *tractable*.
- The ‘Heads’ and ‘Tails’ sides are so similar physically that we make an assumption:

equal likelihood

for

Heads or Tails.

A model for coin tossing: equal likelihood

The two possibilities are just as likely as each other.


$$\mathbb{P}(\text{Heads}) = \frac{1}{2} \quad \mathbb{P}(\text{Tails}) = \frac{1}{2}$$

A model for coin tossing: equal likelihood

The two possibilities are just as likely as each other.

$$\mathbb{P}(\text{Heads}) = \frac{1}{2} \quad \mathbb{P}(\text{Tails}) = \frac{1}{2}$$

We can represent this situation graphically as



Fix some terminology

An **Experiment** observes a phenomenon that has one or more possible **outcomes**.

Fix some terminology

An **Experiment** observes a phenomenon that has one or more possible **outcomes**.

The **Sample space** of an experiment is the set of possible outcomes of the experiment.

Fix some terminology

An **Experiment** observes a phenomenon that has one or more possible **outcomes**.

The **Sample space** of an experiment is the *set* of possible outcomes of the experiment.

An **Event** is any *subset* of the sample space.

Fix some terminology

An **Experiment** observes a phenomenon that has one or more possible **outcomes**.

The **Sample space** of an experiment is the set of possible outcomes of the experiment.

An **Event** is any *subset* of the sample space.

The probability of an event E is denoted by $\mathbb{P}(E)$.

Fix some terminology

An **Experiment** observes a phenomenon that has one or more possible **outcomes**.

The **Sample space** of an experiment is the set of possible outcomes of the experiment.

An **Event** is any *subset* of the sample space.

The probability of an event E is denoted by $\mathbb{P}(E)$.

Example:

- **Experiment:** single toss of a standard die, noting upper face's number.

Fix some terminology

An **Experiment** observes a phenomenon that has one or more possible **outcomes**.

The **Sample space** of an experiment is the set of possible outcomes of the experiment.

An **Event** is any *subset* of the sample space.

The probability of an event E is denoted by $\mathbb{P}(E)$.

Example:

- **Experiment:** single toss of a standard die, noting upper face's number.
- One possible **outcome** would be '3'

Fix some terminology

An **Experiment** observes a phenomenon that has one or more possible **outcomes**.

The **Sample space** of an experiment is the set of possible outcomes of the experiment.


An **Event** is any *subset* of the sample space.

The probability of an event E is denoted by $\mathbb{P}(E)$.

Example:


- **Experiment:** single toss of a standard die, noting upper face's number.
- One possible **outcome** would be '3'
- **Sample space:** $\{1, 2, 3, 4, 5, 6\}$

An equal likelihood model for die-tossing



What's an event?

An equal likelihood model for die-tossing




What's an event? Any subset of the sample space.

Eg. an event is

$$\{3, 6\}$$

An equal likelihood model for die-tossing




What's an event? Any subset of the sample space.

Eg. an event is

$$\{3, 6\}$$

= the set of numbers divisible by 3 in sample space $\{1, 2, 3, 4, 5, 6\}$.

An equal likelihood model for die-tossing



What's an event? Any subset of the sample space.

Eg. an event is

$$\{3, 6\}$$

= the set of numbers divisible by 3 in sample space $\{1, 2, 3, 4, 5, 6\}$.

$$\mathbb{P}(\{3, 6\}) = \frac{|\{3, 6\}|}{|\{1, 2, 3, 4, 5, 6\}|} = \frac{2}{6} = \frac{1}{3}$$

Equal Likelihood for finite Sample Spaces

Generalising from the previous example we have:

- Let S be a finite sample space in which all outcomes are equally likely.
- Let E be an event in S .
- Then the probability of the event E is

$$\mathbb{P}(E) = \frac{|E|}{|S|}$$

Equal Likelihood for finite Sample Spaces

Generalising from the previous example we have:

- Let S be a finite sample space in which all outcomes are equally likely.
- Let E be an event in S .
- Then the probability of the event E is

$$\mathbb{P}(E) = \frac{|E|}{|S|}$$

where $|E|$ is the number of outcomes in E , and
 $|S|$ is the number of outcomes in S .

Properties of Probability

For any finite sample space:

- $\mathbb{P}(E)$ is a real number between 0 and 1, i.e.

$$0 \leq \mathbb{P}(E) \leq 1.$$

Properties of Probability

For any finite sample space:

- $\mathbb{P}(E)$ is a real number between 0 and 1, i.e.

$$0 \leq \mathbb{P}(E) \leq 1.$$

- Probability of the complement is one minus the probability of the event, i.e.

$$\mathbb{P}(\text{not } E) = 1 - \mathbb{P}(E).$$

Properties of Probability

For any finite sample space:

- $\mathbb{P}(E)$ is a real number between 0 and 1, i.e.

$$0 \leq \mathbb{P}(E) \leq 1.$$

- Probability of the complement is one minus the probability of the event, i.e.

$$\mathbb{P}(\text{not } E) = 1 - \mathbb{P}(E).$$

- The sum of the probabilities of all outcomes in the sample space is 1.

Properties of Probability

For any finite sample space:

- $\mathbb{P}(E)$ is a real number between 0 and 1, i.e.

$$0 \leq \mathbb{P}(E) \leq 1.$$

- Probability of the complement is one minus the probability of the event, i.e.

$$\mathbb{P}(\text{not } E) = 1 - \mathbb{P}(E).$$

- The sum of the probabilities of all outcomes in the sample space is 1.
- ' $\mathbb{P}(E) = 1$ ' implies E is certain to occur.*
- ' $\mathbb{P}(E) = 0$ ' implies E is impossible.*

*For infinite sets, this isn't necessarily true. 'Measure theory' explains why.

Previous example of tossing a die:

Probability of event of 'getting a number exactly divisible by 3' is one third, which satisfies

$$0 \leq \frac{1}{3} \leq 1.$$

Previous example of tossing a die:

Probability of event of 'getting a number exactly divisible by 3' is one third, which satisfies

$$0 \leq \frac{1}{3} \leq 1.$$

Probability of 'not E ' is

$$\mathbb{P}(\text{number not divisible by 3}) = 1 - \frac{1}{3} = \frac{2}{3}.$$

Previous example of tossing a die:

Probability of event of 'getting a number exactly divisible by 3' is one third, which satisfies

$$0 \leq \frac{1}{3} \leq 1.$$

Probability of 'not E ' is

$$\mathbb{P}(\text{number not divisible by 3}) = 1 - \frac{1}{3} = \frac{2}{3}.$$

The sum of the probabilities of all outcomes is

$$\mathbb{P}(\{1\}) + \dots + \mathbb{P}(\{6\}) = \frac{1}{6} + \frac{1}{6} + \frac{1}{6} + \frac{1}{6} + \frac{1}{6} + \frac{1}{6} = 1$$

Probability Rules

The Sum Rule

Sum Rule: If events E_1, \dots, E_n are mutually disjoint, i.e. $E_i \cap E_j = \emptyset$ for all $i \neq j$, then

$$\mathbb{P}(E_1 \cup \dots \cup E_n) = \mathbb{P}(E_1) + \dots + \mathbb{P}(E_n).$$

The Sum Rule

Sum Rule: If events E_1, \dots, E_n are mutually disjoint, i.e. $E_i \cap E_j = \emptyset$ for all $i \neq j$, then


$$\mathbb{P}(E_1 \cup \dots \cup E_n) = \mathbb{P}(E_1) + \dots + \mathbb{P}(E_n).$$

Disjoint events exclude one another in the sense that they cannot happen at the same time.

Sum Rule for probability: another die-tossing example

What is the probability that the outcome from a single toss of a die is an odd number?

The six possible outcomes are all disjoint (cannot occur simultaneously).



Thus the sum rule applies.

- We assign equal probabilities to each of these disjoint events (Why?)
- Six possible outcomes in total \rightarrow each has probability $\frac{1}{6}$ of occurring.
- The probability that the die lands with an odd number up is

$$\begin{aligned} & \Pr\left(\boxed{\cdot}\right) + \Pr\left(\boxed{\vdots}\right) + \Pr\left(\boxed{\ddots}\right) \\ &= \frac{1}{6} + \frac{1}{6} + \frac{1}{6} \\ &= \frac{1}{2} \end{aligned}$$

by the sum rule.

Sum Rule for probability. Example: Non-zero numbers

Let $R_n = \{-n, \dots, -2, -1, 0, 1, 2, \dots, n\}$.

What is the probability that a number chosen at random from R_n is non-zero?

Sum Rule for probability. Example: Non-zero numbers

Let $R_n = \{-n, \dots, -2, -1, 0, 1, 2, \dots, n\}$.

What is the probability that a number chosen at random from R_n is non-zero?

We assume that we are equally likely to choose any element of R_n .

Sum Rule for probability. Example: Non-zero numbers

Let $R_n = \{-n, \dots, -2, -1, 0, 1, 2, \dots, n\}$.

What is the probability that a number chosen at random from R_n is non-zero?

We assume that we are equally likely to choose any element of R_n .

- The probability that the number is negative is $\frac{|R_n^-|}{|R_n|} = \frac{n}{2n+1}$.

Sum Rule for probability. Example: Non-zero numbers

Let $R_n = \{-n, \dots, -2, -1, 0, 1, 2, \dots, n\}$.

What is the probability that a number chosen at random from R_n is non-zero?

We assume that we are equally likely to choose any element of R_n .

- The probability that the number is negative is $\frac{|R_n^-|}{|R_n|} = \frac{n}{2n+1}$.
- The probability that the number is positive is $\frac{|R_n^+|}{|R_n|} = \frac{n}{2n+1}$.

Therefore the probability of a number chosen at random from the set $\{-n, \dots, -2, -1, 0, 1, 2, \dots, n\}$ being non-zero is:

Sum Rule for probability. Example: Non-zero numbers

Let $R_n = \{-n, \dots, -2, -1, 0, 1, 2, \dots, n\}$.

What is the probability that a number chosen at random from R_n is non-zero?

We assume that we are equally likely to choose any element of R_n .

- The probability that the number is negative is $\frac{|R_n^-|}{|R_n|} = \frac{n}{2n+1}$.
- The probability that the number is positive is $\frac{|R_n^+|}{|R_n|} = \frac{n}{2n+1}$.

Therefore the probability of a number chosen at random from the set $\{-n, \dots, -2, -1, 0, 1, 2, \dots, n\}$ being non-zero is:

$$\begin{aligned} & \mathbb{P}(\text{the number is negative}) + \mathbb{P}(\text{the number is positive}) \\ &= \frac{n}{2n+1} + \frac{n}{2n+1} = \frac{2n}{2n+1}. \end{aligned}$$

The Product Rule

- **Product Rule:** If events E_1, \dots, E_n are 'independent' of each other; then the probability of composite event ' E_1 and E_2 and ... and E_n ' is

$$\mathbb{P}(E_1 \times E_2 \times \dots \times E_n) = \mathbb{P}(E_1) \times \mathbb{P}(E_2) \times \dots \mathbb{P}(E_n).$$

The Product Rule

- **Product Rule:** If events E_1, \dots, E_n are 'independent' of each other; then the probability of composite event ' E_1 and E_2 and ... and E_n ' is

$$\mathbb{P}(E_1 \times E_2 \times \dots \times E_n) = \mathbb{P}(E_1) \times \mathbb{P}(E_2) \times \dots \mathbb{P}(E_n).$$

- To see what we mean by 'independent', consider a procedure that can be broken down into successive tasks, each of which could be done in a number of ways. If the choice of the way to do any one task had no influence on the choice of ways to do any other of the tasks, then the tasks would be independent.


The Product Rule

- **Product Rule:** If events E_1, \dots, E_n are 'independent' of each other; then the probability of composite event ' E_1 and E_2 and ... and E_n ' is

$$\mathbb{P}(E_1 \times E_2 \times \dots \times E_n) = \mathbb{P}(E_1) \times \mathbb{P}(E_2) \times \dots \mathbb{P}(E_n).$$

- To see what we mean by 'independent', consider a procedure that can be broken down into successive tasks, each of which could be done in a number of ways. If the choice of the way to do any one task had no influence on the choice of ways to do any other of the tasks, then the tasks would be independent.
- A formal definition of independence will be given later.


Product Rule probability example: Tossing two dice



- What is the probability that the outcome from tossing a pair of dice is '4' for the first die and '5' for the second die i.e.



Product Rule probability example: Tossing two dice




- What is the probability that the outcome from tossing a pair of dice is '4' for the first die and '5' for the second die i.e.



- We assume that the outcomes for each die are **independent**, i.e that they don't influence one another at all.

Product Rule probability example: Tossing two dice



- What is the probability that the outcome from tossing a pair of dice is '4' for the first die and '5' for the second die i.e.



- We assume that the outcomes for each die are **independent**, i.e. that they don't influence one another at all.
- Hence the product rule applies.

$$\begin{aligned}
& \Pr\left(\boxed{\begin{smallmatrix} \cdot & \cdot \\ \cdot & \cdot \end{smallmatrix}} \quad \boxed{5}\right) \\
&= \Pr\left(\boxed{\begin{smallmatrix} \cdot & \cdot \\ \cdot & \cdot \end{smallmatrix}}\right) \times \Pr\left(\boxed{5}\right) \\
&= \frac{1}{6} \times \frac{1}{6} \\
&= \frac{1}{36}
\end{aligned}$$


by the Product Rule.

An example of the Sum and Product Rules used together

- Often we combine use of the Sum and Product rules in one problem.


An example of the Sum and Product Rules used together

- Often we combine use of the Sum and Product rules in one problem.
- For example, *what is the probability of getting an odd total when tossing a pair of dice?*



An example of the Sum and Product Rules used together

- Often we combine use of the Sum and Product rules in one problem.
- For example, *what is the probability of getting an odd total when tossing a pair of dice?*



- To obtain an odd total, either
 - the first die must give odd and the second die even; or
 - the first die must give even and the second die odd.
- These two possibilities are **disjoint**, so the sum rule applies:
$$\mathbb{P}(\text{odd total}) = \mathbb{P}(\text{1st odd, 2nd even}) + \mathbb{P}(\text{1st even, 2nd odd})$$

- But now consider $\mathbb{P}(1\text{st odd, 2nd even})$. The events
"1st odd" and "2nd even"
are **independent** of each other; they don't affect each other.

- But now consider $\mathbb{P}(1\text{st odd, 2nd even})$. The events
"1st odd" and "2nd even"
are **independent** of each other; they don't affect each other.
- Hence the **product rule** applies to this part of the problem:

$$\begin{aligned}\mathbb{P}(1\text{st odd, 2nd even}) &= \mathbb{P}(1\text{st odd}) \times \mathbb{P}(2\text{nd even}) \\ &= \frac{3}{6} \times \frac{3}{6} = \frac{1}{2} \times \frac{1}{2} = \frac{1}{4}\end{aligned}$$

- But now consider $\mathbb{P}(\text{1st odd, 2nd even})$. The events
“1st odd” and “2nd even”
are **independent** of each other; they don't affect each other.
- Hence the **product rule** applies to this part of the problem:

$$\begin{aligned}\mathbb{P}(\text{1st odd, 2nd even}) &= \mathbb{P}(\text{1st odd}) \times \mathbb{P}(\text{2nd even}) \\ &= \frac{3}{6} \times \frac{3}{6} = \frac{1}{2} \times \frac{1}{2} = \frac{1}{4}\end{aligned}$$

- Similarly, $\mathbb{P}(\text{1st even, 2nd odd}) = \frac{1}{4}$

- But now consider $\mathbb{P}(\text{1st odd, 2nd even})$. The events
“1st odd” and “2nd even”
are **independent** of each other; they don't affect each other.
- Hence the **product rule** applies to this part of the problem:

$$\begin{aligned}\mathbb{P}(\text{1st odd, 2nd even}) &= \mathbb{P}(\text{1st odd}) \times \mathbb{P}(\text{2nd even}) \\ &= \frac{3}{6} \times \frac{3}{6} = \frac{1}{2} \times \frac{1}{2} = \frac{1}{4}\end{aligned}$$

- Similarly, $\mathbb{P}(\text{1st even, 2nd odd}) = \frac{1}{4}$
- Putting it all together,

$$\begin{aligned}\mathbb{P}(\text{odd total}) &= \mathbb{P}(\text{1st odd, 2nd even}) + \mathbb{P}(\text{1st even, 2nd odd}) \\ &= \frac{1}{4} + \frac{1}{4} = \frac{1}{2}.\end{aligned}$$