

# Discrete Mathematical Models

## Lecture 30

---


Kane Townsend

Semester 2, 2024

# Wikipedia Example


---

# Wiki WebGraph




<https://en.wikipedia.org/wiki/PageRank> We will fill in  $\mathbb{P}\mathbb{R}$ . It will differ slightly to seen on wiki due to a small difference in our algorithm which is common to do for large  $n$ ).

## Example 5



At left is the Wikipedia example we mentioned in the previous lecture consisting of miniweb of 11 pages and 17 hyperlinks. Colours, variable sizes and PageRanks have been removed and the bottom five vertices have been labelled G to K, following the given labelling of the top six vertices. The layout is similar to that in the original.


## Example 5



Using the steady state method we have been discussing, we will derive the PageRanks given on the Wikipedia diagram.

At left is the Wikipedia example we mentioned in the previous lecture consisting of miniweb of 11 pages and 17 hyperlinks. Colours, variable sizes and PageRanks have been removed and the bottom five vertices have been labelled G to K, following the given labelling of the top six vertices. The layout is similar to that in the original.

## Example 5



Using the steady state method we have been discussing, we will derive the PageRanks given on the Wikipedia diagram.

**Step 1:** Construct  $G^+$  from  $G$

At left is the Wikipedia example we mentioned in the previous lecture consisting of miniweb of 11 pages and 17 hyperlinks. Colours, variable sizes and PageRanks have been removed and the bottom five vertices have been labelled G to K, following the given labelling of the top six vertices. The layout is similar to that in the original.

## Example 5 (cont.)

**Step 2:** Compile the basic transition matrix  $T$ .

## Example 5 (cont.)

**Step 2:** Compile the basic transition matrix  $T$ .

Let  $n_i$  be the number vertices to which  $i$  is adjacent in  $G^+$ , that is

$$n_i = |\{j : (i, j) \in E(G^+)\}|$$

The  $p_{ij}$  of a transition from vertex  $j$  to  $i$  is given by

$$p_{ij} = \begin{cases} 1/n_i & \text{if } n_i \neq 0 \text{ and } (j, i) \in E(G^+) \\ 0 & \text{otherwise} \end{cases}$$



## Example 5 (cont.)

**Step 2:** Compile the basic transition matrix  $T$ .

Let  $n_i$  be the number vertices to which  $i$  is adjacent in  $G^+$ , that is

$$n_i = |\{j : (i, j) \in E(G^+)\}|$$

The  $p_{ij}$  of a transition from vertex  $j$  to  $i$  is given by

$$p_{ij} = \begin{cases} 1/n_i & \text{if } n_i \neq 0 \text{ and } (j, i) \in E(G^+) \\ 0 & \text{otherwise} \end{cases}$$

## Example 5 (cont.)

**Step 2:** Compile the basic transition matrix  $T$ .

Let  $n_i$  be the number vertices to which  $i$  is adjacent in  $G^+$ , that is

$$n_i = |\{j : (i, j) \in E(G^+)\}|$$

The  $p_{ij}$  of a transition from vertex  $j$  to  $i$  is given by

$$p_{ij} = \begin{cases} 1/n_i & \text{if } n_i \neq 0 \text{ and } (j, i) \in E(G^+) \\ 0 & \text{otherwise} \end{cases}$$

For our Wikipedia example we get

$$T = \begin{bmatrix} 0 & 0 & 0 & 1/2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1/10 & 0 & 1 & 1/2 & 1/3 & 1/2 & 1/2 & 1/2 & 1/2 & 0 & 0 \\ 1/10 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1/10 & 0 & 0 & 0 & 1/3 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1/10 & 0 & 0 & 0 & 0 & 1/2 & 1/2 & 1/2 & 1/2 & 1 & 1 \\ 1/10 & 0 & 0 & 0 & 1/3 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1/10 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1/10 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1/10 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1/10 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1/10 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

## Example 5 (cont.)

**Step 3:** Compile the matrix  $M = (\alpha/n)U + (1 - \alpha)T$

## Example 5 (cont.)

**Step 3:** Compile the matrix  $M = (\alpha/n)U + (1 - \alpha)T$

That is multiply each entry of  $T$  by  $(1 - \alpha)$  and add  $\alpha/n$  to each entry. Recall that Google uses an 85% damping factor, so we set  $(1 - \alpha) = 0.85$ .

## Example 5 (cont.)

**Step 3:** Compile the matrix  $M = (\alpha/n)U + (1 - \alpha)T$

That is multiply each entry of  $T$  by  $(1 - \alpha)$  and add  $\alpha/n$  to each entry. Recall that Google uses an 85% damping factor, so we set  $(1 - \alpha) = 0.85$ .

$$M = (\alpha/n)U + (1 - \alpha)T =$$

$$\begin{bmatrix} 3/220 & 3/220 & 3/220 & 193/440 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 \\ 217/2200 & 3/220 & 19/22 & 193/440 & 49/165 & 193/440 & 193/440 & 193/440 & 193/440 & 3/220 & 3/220 \\ 217/2200 & 19/22 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 \\ 217/2200 & 3/220 & 3/220 & 3/220 & 49/165 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 \\ 217/2200 & 3/220 & 3/220 & 3/220 & 3/220 & 193/440 & 193/440 & 193/440 & 193/440 & 19/22 & 19/22 \\ 217/2200 & 3/220 & 3/220 & 3/220 & 49/165 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 \\ 217/2200 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 \\ 217/2200 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 \\ 217/2200 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 \\ 217/2200 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 \\ 217/2200 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 & 3/220 \end{bmatrix}$$

## Example 5 (cont.)

**Step 4:** Use a computer to solve  $M\mathcal{R} = \mathcal{R}$  using our 'Shortcut method'. For example 'Gauss-Jordan Elimination' in the *Matrix Reshish* online matrix calculator, gives

## Example 5 (cont.)

**Step 4:** Use a computer to solve  $M\mathbb{R} = \mathbb{R}$  using our 'Shortcut method'. For example 'Gauss-Jordan Elimination' in the *Matrix Reshish* online matrix calculator, gives

$$\mathbb{R} = \begin{bmatrix} 5135730/169545563 \\ 3195978661779342/8292828011094295 \\ 165932254119056/482651365725065 \\ 6644106/169545563 \\ 5317726843699199/65574887090606070 \\ 6644106/169545563 \\ 2748522/169545563 \\ 2748522/169545563 \\ 2748522/169545563 \\ 2748522/169545563 \\ 2748522/169545563 \end{bmatrix} \approx \begin{bmatrix} 0.0303 \\ 0.3854 \\ 0.3438 \\ 0.0392 \\ 0.0811 \\ 0.0392 \\ 0.0162 \\ 0.0162 \\ 0.0162 \\ 0.0162 \\ 0.0162 \end{bmatrix} \quad (4d.p.)$$

# Iteration method

---



# Iterative Approximation method

When  $n$  is huge, as it is with the WWW, solving the the  $n \times n$  linear system  $(M - I)\mathbf{PR} = 0$  becomes computationally infeasible.

# Iterative Approximation method

When  $n$  is huge, as it is with the WWW, solving the the  $n \times n$  linear system  $(M - I)\overline{PR} = 0$  becomes computationally infeasible.

A computationally simpler method starts from the defining equation attempts to find  $\overline{PR}$  by iteratively calculating the chain of probability vectors  $P_0, P_1, \dots$  where  $P_0$  is arbitrary and  $P_k = MP_{k-1}$  for  $k \geq 1$  (so  $P_k = M^k P_0$ ).

A steady state is reached when  $P_k \approx P_{k-1}$ . Then declare that  $\overline{PR} \approx P_k$ .

# Iterative Approximation method

When  $n$  is huge, as it is with the WWW, solving the the  $n \times n$  linear system  $(M - I)\overline{PR} = 0$  becomes computationally infeasible.

A computationally simpler method starts from the defining equation attempts to find  $\overline{PR}$  by iteratively calculating the chain of probability vectors  $P_0, P_1, \dots$  where  $P_0$  is arbitrary and  $P_k = MP_{k-1}$  for  $k \geq 1$  (so  $P_k = M^k P_0$ ).

A steady state is reached when  $P_k \approx P_{k-1}$ . Then declare that  $\overline{PR} \approx P_k$ .

$$\begin{aligned}\text{Now } P_k = MP_{k-1} &= [(\alpha/n)U + (1 - \alpha)T]P_{k-1} \\ &= (\alpha/n)\mathbf{1} + (1 - \alpha)TP_{k-1}\end{aligned}$$

where  $\mathbf{1}$  is column of 1's.

# Iterative Approximation method

When  $n$  is huge, as it is with the WWW, solving the the  $n \times n$  linear system  $(M - I)\overline{PR} = 0$  becomes computationally infeasible.

A computationally simpler method starts from the defining equation attempts to find  $\overline{PR}$  by iteratively calculating the chain of probability vectors  $P_0, P_1, \dots$  where  $P_0$  is arbitrary and  $P_k = MP_{k-1}$  for  $k \geq 1$  (so  $P_k = M^k P_0$ ).

A steady state is reached when  $P_k \approx P_{k-1}$ . Then declare that  $\overline{PR} \approx P_k$ .

$$\begin{aligned}\text{Now } P_k = MP_{k-1} &= [(\alpha/n)U + (1 - \alpha)T]P_{k-1} \\ &= (\alpha/n)\mathbf{1} + (1 - \alpha)TP_{k-1}\end{aligned}$$

where  $\mathbf{1}$  is column of 1's. It is natural to start with all ranks equal.

# Iterative Approximation method

When  $n$  is huge, as it is with the WWW, solving the the  $n \times n$  linear system  $(M - I)\overline{PR} = 0$  becomes computationally infeasible.

A computationally simpler method starts from the defining equation attempts to find  $\overline{PR}$  by iteratively calculating the chain of probability vectors  $P_0, P_1, \dots$  where  $P_0$  is arbitrary and  $P_k = MP_{k-1}$  for  $k \geq 1$  (so  $P_k = M^k P_0$ ).

A steady state is reached when  $P_k \approx P_{k-1}$ . Then declare that  $\overline{PR} \approx P_k$ .

$$\begin{aligned}\text{Now } P_k = MP_{k-1} &= [(\alpha/n)U + (1 - \alpha)T]P_{k-1} \\ &= (\alpha/n)\mathbf{1} + (1 - \alpha)TP_{k-1}\end{aligned}$$

where  $\mathbf{1}$  is column of 1's. It is natural to start with all ranks equal. So the iterative scheme is

$$P_0 = (1/n)\mathbf{1}; \quad P_k = \alpha P_0 + (1 - \alpha)TP_{k-1}, \quad k \geq 1.$$

# Iterative Approximation method

When  $n$  is huge, as it is with the WWW, solving the the  $n \times n$  linear system  $(M - I)\overline{PR} = 0$  becomes computationally infeasible.

A computationally simpler method starts from the defining equation attempts to find  $\overline{PR}$  by iteratively calculating the chain of probability vectors  $P_0, P_1, \dots$  where  $P_0$  is arbitrary and  $P_k = MP_{k-1}$  for  $k \geq 1$  (so  $P_k = M^k P_0$ ).

A steady state is reached when  $P_k \approx P_{k-1}$ . Then declare that  $\overline{PR} \approx P_k$ .

$$\begin{aligned}\text{Now } P_k = MP_{k-1} &= [(\alpha/n)U + (1 - \alpha)T]P_{k-1} \\ &= (\alpha/n)\mathbf{1} + (1 - \alpha)TP_{k-1}\end{aligned}$$

where  $\mathbf{1}$  is column of 1's. It is natural to start with all ranks equal. So the iterative scheme is

$$P_0 = (1/n)\mathbf{1}; \quad P_k = \alpha P_0 + (1 - \alpha)TP_{k-1}, \quad k \geq 1.$$

Each iteration takes a weighted average of teleporting and hyperlinking.

## Example 4C

In Example 4B we used the equation-solving method to find  $\mathcal{PR}$ .

$$\text{For } T = \begin{bmatrix} 0 & 0 & 0 & 1/3 \\ 1/2 & 0 & 1/2 & 1/3 \\ 0 & 1 & 0 & 1/3 \\ 1/2 & 0 & 1/2 & 0 \end{bmatrix} \text{ and } \alpha = 0.1 \text{ we found } \mathcal{PR} = \begin{bmatrix} .10 \\ .30 \\ .37 \\ .23 \end{bmatrix} \text{ to 2d.p.}$$

## Example 4C

In Example 4B we used the equation-solving method to find  $\mathcal{PR}$ .

$$\text{For } T = \begin{bmatrix} 0 & 0 & 0 & 1/3 \\ 1/2 & 0 & 1/2 & 1/3 \\ 0 & 1 & 0 & 1/3 \\ 1/2 & 0 & 1/2 & 0 \end{bmatrix} \text{ and } \alpha = 0.1 \text{ we found } \mathcal{PR} = \begin{bmatrix} .10 \\ .30 \\ .37 \\ .23 \end{bmatrix} \text{ to 2d.p.}$$

Let's try the same problem using iterative approximation:

$$P_0 = \begin{bmatrix} .25 \\ .25 \\ .25 \\ .25 \end{bmatrix} \quad P_k = \begin{bmatrix} .025 \\ .025 \\ .025 \\ .025 \end{bmatrix} + \begin{bmatrix} 0 & 0 & 0 & .3 \\ .45 & 0 & .45 & .3 \\ 0 & .9 & 0 & .3 \\ .45 & 0 & .45 & 0 \end{bmatrix} P_{k-1}$$



## Example 4C

In Example 4B we used the equation-solving method to find  $\overline{PR}$ .

$$\text{For } T = \begin{bmatrix} 0 & 0 & 0 & 1/3 \\ 1/2 & 0 & 1/2 & 1/3 \\ 0 & 1 & 0 & 1/3 \\ 1/2 & 0 & 1/2 & 0 \end{bmatrix} \text{ and } \alpha = 0.1 \text{ we found } \overline{PR} = \begin{bmatrix} .10 \\ .30 \\ .37 \\ .23 \end{bmatrix} \text{ to 2d.p.}$$

Let's try the same problem using iterative approximation:

$$P_0 = \begin{bmatrix} .25 \\ .25 \\ .25 \\ .25 \end{bmatrix} \quad P_k = \begin{bmatrix} .025 \\ .025 \\ .025 \\ .025 \end{bmatrix} + \begin{bmatrix} 0 & 0 & 0 & .3 \\ .45 & 0 & .45 & .3 \\ 0 & .9 & 0 & .3 \\ .45 & 0 & .45 & 0 \end{bmatrix} P_{k-1}$$

Results of the first ten iterations, rounded to 2d.p. Calcs used 15d.p.

$k$	1	2	3	4	5	6	7	8	9	10
$P_k$	$\begin{bmatrix} .10 \\ .33 \\ .33 \\ .25 \end{bmatrix}$	$\begin{bmatrix} .10 \\ .29 \\ .39 \\ .22 \end{bmatrix}$	$\begin{bmatrix} .09 \\ .31 \\ .35 \\ .25 \end{bmatrix}$	$\begin{bmatrix} .10 \\ .30 \\ .38 \\ .22 \end{bmatrix}$	$\begin{bmatrix} .09 \\ .31 \\ .36 \\ .24 \end{bmatrix}$	$\begin{bmatrix} .10 \\ .30 \\ .37 \\ .23 \end{bmatrix}$	$\begin{bmatrix} .09 \\ .31 \\ .36 \\ .24 \end{bmatrix}$	$\begin{bmatrix} .10 \\ .30 \\ .37 \\ .23 \end{bmatrix}$	$\begin{bmatrix} .09 \\ .30 \\ .37 \\ .24 \end{bmatrix}$	$\begin{bmatrix} .10 \\ .30 \\ .37 \\ .23 \end{bmatrix}$

## Example 4C

In Example 4B we used the equation-solving method to find  $PR$ .

$$\text{For } T = \begin{bmatrix} 0 & 0 & 0 & 1/3 \\ 1/2 & 0 & 1/2 & 1/3 \\ 0 & 1 & 0 & 1/3 \\ 1/2 & 0 & 1/2 & 0 \end{bmatrix} \text{ and } \alpha = 0.1 \text{ we found } PR = \begin{bmatrix} .10 \\ .30 \\ .37 \\ .23 \end{bmatrix} \text{ to 2d.p.}$$

Let's try the same problem using iterative approximation:

$$P_0 = \begin{bmatrix} .25 \\ .25 \\ .25 \\ .25 \end{bmatrix} \quad P_k = \begin{bmatrix} .025 \\ .025 \\ .025 \\ .025 \end{bmatrix} + \begin{bmatrix} 0 & 0 & 0 & .3 \\ .45 & 0 & .45 & .3 \\ 0 & .9 & 0 & .3 \\ .45 & 0 & .45 & 0 \end{bmatrix} P_{k-1}$$

Results of the first ten iterations, rounded to 2d.p. Calcs used 15d.p.

$k$	1	2	3	4	5	6	7	8	9	10
$P_k$	$\begin{bmatrix} .10 \\ .33 \\ .33 \\ .25 \end{bmatrix}$	$\begin{bmatrix} .10 \\ .29 \\ .39 \\ .22 \end{bmatrix}$	$\begin{bmatrix} .09 \\ .31 \\ .35 \\ .25 \end{bmatrix}$	$\begin{bmatrix} .10 \\ .30 \\ .38 \\ .22 \end{bmatrix}$	$\begin{bmatrix} .09 \\ .31 \\ .36 \\ .24 \end{bmatrix}$	$\begin{bmatrix} .10 \\ .30 \\ .37 \\ .23 \end{bmatrix}$	$\begin{bmatrix} .09 \\ .31 \\ .36 \\ .24 \end{bmatrix}$	$\begin{bmatrix} .10 \\ .30 \\ .37 \\ .23 \end{bmatrix}$	$\begin{bmatrix} .09 \\ .30 \\ .37 \\ .24 \end{bmatrix}$	$\begin{bmatrix} .10 \\ .30 \\ .37 \\ .23 \end{bmatrix}$

Pretty good after just 2 iterations! Within 1%-point after 4 iterations.

## Example 4C

In Example 4B we used the equation-solving method to find  $PR$ .

$$\text{For } T = \begin{bmatrix} 0 & 0 & 0 & 1/3 \\ 1/2 & 0 & 1/2 & 1/3 \\ 0 & 1 & 0 & 1/3 \\ 1/2 & 0 & 1/2 & 0 \end{bmatrix} \text{ and } \alpha = 0.1 \text{ we found } PR = \begin{bmatrix} .10 \\ .30 \\ .37 \\ .23 \end{bmatrix} \text{ to 2d.p.}$$

Let's try the same problem using iterative approximation:

$$P_0 = \begin{bmatrix} .25 \\ .25 \\ .25 \\ .25 \end{bmatrix} \quad P_k = \begin{bmatrix} .025 \\ .025 \\ .025 \\ .025 \end{bmatrix} + \begin{bmatrix} 0 & 0 & 0 & .3 \\ .45 & 0 & .45 & .3 \\ 0 & .9 & 0 & .3 \\ .45 & 0 & .45 & 0 \end{bmatrix} P_{k-1}$$

Results of the first ten iterations, rounded to 2d.p. Calcs used 15d.p.

$k$	1	2	3	4	5	6	7	8	9	10
$P_k$	$\begin{bmatrix} .10 \\ .33 \\ .33 \\ .25 \end{bmatrix}$	$\begin{bmatrix} .10 \\ .29 \\ .39 \\ .22 \end{bmatrix}$	$\begin{bmatrix} .09 \\ .31 \\ .35 \\ .25 \end{bmatrix}$	$\begin{bmatrix} .10 \\ .30 \\ .38 \\ .22 \end{bmatrix}$	$\begin{bmatrix} .09 \\ .31 \\ .36 \\ .24 \end{bmatrix}$	$\begin{bmatrix} .10 \\ .30 \\ .37 \\ .23 \end{bmatrix}$	$\begin{bmatrix} .09 \\ .31 \\ .36 \\ .24 \end{bmatrix}$	$\begin{bmatrix} .10 \\ .30 \\ .37 \\ .23 \end{bmatrix}$	$\begin{bmatrix} .09 \\ .30 \\ .37 \\ .24 \end{bmatrix}$	$\begin{bmatrix} .10 \\ .30 \\ .37 \\ .23 \end{bmatrix}$

Pretty good after just 2 iterations! Within 1%-point after 4 iterations.

**END OF SECTION D3**

End of Course Notes :')

# My research in graph theory

---

## Set up!

Let  $G$  be a simple graph with some upper bound on vertex degree. That means no loops, parallel edges and not infinitely many edges at one vertex!

## Set up!

Let  $G$  be a simple graph with some upper bound on vertex degree. That means no loops, parallel edges and not infinitely many edges at one vertex!

We assign weights of 1 to each edge in  $G$ .

## Set up!

Let  $G$  be a simple graph with some upper bound on vertex degree. That means no loops, parallel edges and not infinitely many edges at one vertex!

We assign weights of 1 to each edge in  $G$ .

A shortest path from  $u$  to  $v$  is called a **geodesic** in  $G$ .



## Set up!

Let  $G$  be a simple graph with some upper bound on vertex degree. That means no loops, parallel edges and not infinitely many edges at one vertex!

We assign weights of 1 to each edge in  $G$ .

A shortest path from  $u$  to  $v$  is called a **geodesic** in  $G$ .



A graph  $G$  is called **geodetic** if for all vertices  $u, v \in V(G)$ , there exists a unique geodesic from  $u$  to  $v$

# Geodetic graphs

Examples: Trees, complete graphs, odd cycle graphs etc.



# Geodetic graphs

Examples: Trees, complete graphs, odd cycle graphs etc.



# Geodetic graphs


Examples: Trees, complete graphs, odd cycle graphs etc.




Question: Characterise all geodetic graphs (Ore in The Theory of Graphs (1962)).

The classic example of a non-geodetic graph is an even cycle graph.

The classic example of a non-geodetic graph is an even cycle graph.



The classic example of a non-geodetic graph is an even cycle graph.



Given a graph there are minimal spanning trees of  $G$  consisting of geodesics from  $v$  to every other vertex in  $G$ . However, in a geodetic graph  $G$ , such a minimal spanning tree is unique  $v \in V(G)$ .

## Combining geodetic graphs with bridges and cut vertices




Given two geodetic graphs  $G_1$  and  $G_2$  we can join them at a single vertex or via a single edge and produce a new geodetic graph. Also, if a graph  $G$  is geodetic then one can subdivide the edges and obtain a new geodetic graph.



# Combining geodetic graphs with bridges and cut vertices

Given two geodetic graphs  $G_1$  and  $G_2$  we can join them at a single vertex or via a single edge and produce a new geodetic graph. Also, if a graph  $G$  is geodetic then one can subdivide the edges and obtain a new geodetic graph.

Recently my previous supervisor's colleagues (Weißband his PhD student Stober) used the NAUTY program to generate geodetic graphs (with no cut vertices) of up to 25 vertices (<https://arxiv.org/pdf/2308.08970>). There are 12969219563 geodetic graphs with 19 vertices, only 19 of them have no cut vertices. Of those 19, only 2 are not subdivisions of a smaller geodetic graph, one is  $K_{19}$  and other is  $F_3$ .




## Geodesics in a geodetic graph cannot stay close forever

Suppose we have two long geodesics in a geodetic graph  $G$  travelling next to each other (never touching) such that they are at most  $M > 0$  away from each other. Then by the generalised pigeonhole principle there are many times they are some they are distance  $0 < m \leq M$  arbitrarily many times.


## Geodesics in a geodetic graph cannot stay close forever

Suppose we have two long geodesics in a geodetic graph  $G$  travelling next to each other (never touching) such that they are at most  $M > 0$  away from each other. Then by the generalised pigeonhole principle there are many times they are some they are distance  $0 < m \leq M$  arbitrarily many times.



# Geodesics in a geodetic graph cannot stay close forever

Suppose we have two long geodesics in a geodetic graph  $G$  travelling next to each other (never touching) such that they are at most  $M > 0$  away from each other. Then by the generalised pigeonhole principle there are many times they are some they are distance  $0 < m \leq M$  arbitrarily many times.



We get a contradiction!

## Vertex transitive graphs

Let  $\text{Aut}(G)$  be the collection of all graph isomorphisms  $G \rightarrow G$ .

# Vertex transitive graphs

Let  $\text{Aut}(G)$  be the collection of all graph isomorphisms  $G \rightarrow G$ .

A graph is called **vertex transitive** if for every  $u, v \in V(G)$  there exists  $f \in \text{Aut}(G)$  such that  $f(u) = v$ .

# Vertex transitive graphs

Let  $\text{Aut}(G)$  be the collection of all graph isomorphisms  $G \rightarrow G$ .

A graph is called **vertex transitive** if for every  $u, v \in V(G)$  there exists  $f \in \text{Aut}(G)$  such that  $f(u) = v$ .


Examples: Cycle graphs, complete graphs and more interesting infinite examples like this one below:

# Vertex transitive graphs

Let  $\text{Aut}(G)$  be the collection of all graph isomorphisms  $G \rightarrow G$ .

A graph is called **vertex transitive** if for every  $u, v \in V(G)$  there exists  $f \in \text{Aut}(G)$  such that  $f(u) = v$ .

Examples: Cycle graphs, complete graphs and more interesting infinite examples like this one below:





## Theorem and conjecture

Recall that a **simple circuit** is a circuit with no repeated vertices.

# Theorem and conjecture

Recall that a **simple circuit** is a circuit with no repeated vertices.

We call a simple circuit in a graph  $G$  **isometric** if the geodesic between any two vertices in the simple circuit lie on the circuit.

# Theorem and conjecture

Recall that a **simple circuit** is a circuit with no repeated vertices.

We call a simple circuit in a graph  $G$  **isometric** if the geodesic between any two vertices in the simple circuit lie on the circuit.

## **Theorem (Elder-Gardam-Spriano-Piggott-T)**

*Let  $G$  be a vertex transitive geodetic graph. Then  $G$  has bounded size isometric simple circuits.*

# Theorem and conjecture

Recall that a **simple circuit** is a circuit with no repeated vertices.

We call a simple circuit in a graph  $G$  **isometric** if the geodesic between any two vertices in the simple circuit lie on the circuit.

## **Theorem (Elder-Gardam-Spriano-Piggott-T)**

*Let  $G$  be a vertex transitive geodetic graph. Then  $G$  has bounded size isometric simple circuits.*

This means that vertex transitive geodetic graphs are ‘quasi-transitive’ to a tree (like a tree up to bounded tinkering).

# Theorem and conjecture

Recall that a **simple circuit** is a circuit with no repeated vertices.

We call a simple circuit in a graph  $G$  **isometric** if the geodesic between any two vertices in the simple circuit lie on the circuit.

## **Theorem (Elder-Gardam-Spriano-Piggott-T)**

*Let  $G$  be a vertex transitive geodetic graph. Then  $G$  has bounded size isometric simple circuits.*

This means that vertex transitive geodetic graphs are ‘quasi-transitive’ to a tree (like a tree up to bounded tinkering).

**Conjecture:** Let  $G$  be a transitive geodetic graph. Then every vertex is a cut vertex? Then there is a bound on simple circuits in  $G$ ?

## Proof of theorem outline

The proof requires understanding of topology outside the scope of this course.

# Proof of theorem outline

The proof requires understanding of topology outside the scope of this course.

**Step 1:** Define something called the geodesic boundary in an infinite graph by setting up an equivalence relation on the collection of geodesic rays in the graph. Motivated by prior work on the Gromov boundary of hyperbolic metric spaces.




# Proof of theorem outline

The proof requires understanding of topology outside the scope of this course.

**Step 1:** Define something called the geodesic boundary in an infinite graph by setting up an equivalence relation on the collection of geodesic rays in the graph. Motivated by prior work on the Gromov boundary of hyperbolic metric spaces.




**Step 2:** Define a topology/metric on the geodesic boundary of a geodesic graph which is dependent on a basepoint vertex.






## Proof of theorem outline (cont.)

**Step 3:** Prove that geodetic graphs cannot contain an ‘infinite onion’ (proof by contradiction using the topological properties of the geodesic boundary of a geodetic graph).



## Proof of theorem outline (cont.)

**Step 3:** Prove that geodetic graphs cannot contain an ‘infinite onion’ (proof by contradiction using the topological properties of the geodesic boundary of a geodetic graph).



**Step 4:** Prove that a transitive geodetic graph cannot have unbounded size of isometric simple circuits as this implies that you can construct an infinite onion.

## References to preprints

<https://arxiv.org/pdf/2311.03730>

<https://arxiv.org/pdf/2211.13397>