

Discrete Mathematical Models

Lecture 11

Kane Townsend

Semester 2, 2024

Section B: Digital Information (cont.)

Section B2: Sequences, Induction, Sorting

Sequences

Text Reference (Epp)

3ed: Sections 4.1-4, 8.1-3 (Sequences and induction),
9.3,5 (Sorting)

4ed: Sections 5.1-4,6-8, (Sequences and induction),
11.3,5 (Sorting)

5ed: Sections 5.1-4,6-7, (Sequences and induction),
11.3,5 (Sorting)

Sequences

Let S be a set and $I \subseteq \mathbb{Z}_{\geq 0} = \mathbb{N}^*$. A function $a: I \rightarrow S$ is called a **sequence in S** .

Sequences

Let S be a set and $I \subseteq \mathbb{Z}_{\geq 0} = \mathbb{N}^*$. A function $a: I \rightarrow S$ is called a **sequence in S** . Special sequence notation is often used:

Function notation	Sequence notation
$a: I \rightarrow S, n \mapsto a(n)$	$(a_n)_{n \in I} \subseteq S$

Sequences

Let S be a set and $I \subseteq \mathbb{Z}_{\geq 0} = \mathbb{N}^*$. A function $a: I \rightarrow S$ is called a **sequence in S** . Special sequence notation is often used:

Function notation	Sequence notation
$a: I \rightarrow S, n \mapsto a(n)$	$(a_n)_{n \in I} \subseteq S$

The notation $(a_n)_{n \in I}$ indicates that the function can be represented as an *ordered -tuple* (possibly infinite -tuple).

Sequences

Let S be a set and $I \subseteq \mathbb{Z}_{\geq 0} = \mathbb{N}^*$. A function $a: I \rightarrow S$ is called a **sequence in S** . Special sequence notation is often used:

Function notation	Sequence notation
$a: I \rightarrow S, n \mapsto a(n)$	$(a_n)_{n \in I} \subseteq S$

The notation $(a_n)_{n \in I}$ indicates that the function can be represented as an *ordered -tuple* (possibly infinite -tuple).

Examples:

- $I = \{1, 2, 3\} : (a_n)_{n \in I} = (a_1, a_2, a_3).$

Sequences

Let S be a set and $I \subseteq \mathbb{Z}_{\geq 0} = \mathbb{N}^*$. A function $a: I \rightarrow S$ is called a **sequence in S** . Special sequence notation is often used:

Function notation	Sequence notation
$a: I \rightarrow S, n \mapsto a(n)$	$(a_n)_{n \in I} \subseteq S$

The notation $(a_n)_{n \in I}$ indicates that the function can be represented as an *ordered -tuple* (possibly infinite -tuple).

Examples:

- $I = \{1, 2, 3\} : (a_n)_{n \in I} = (a_1, a_2, a_3).$
- $I = \mathbb{N}^* : (a_n)_{n \in I} = (a_0, a_1, a_2, \dots).$

Sequences

Let S be a set and $I \subseteq \mathbb{Z}_{\geq 0} = \mathbb{N}^*$. A function $a: I \rightarrow S$ is called a **sequence in S** . Special sequence notation is often used:

Function notation	Sequence notation
$a: I \rightarrow S, n \mapsto a(n)$	$(a_n)_{n \in I} \subseteq S$

The notation $(a_n)_{n \in I}$ indicates that the function can be represented as an *ordered -tuple* (possibly infinite -tuple).

Examples:

- $I = \{1, 2, 3\} : (a_n)_{n \in I} = (a_1, a_2, a_3).$
- $I = \mathbb{N}^* : (a_n)_{n \in I} = (a_0, a_1, a_2, \dots).$

In practice we sometimes leave out the parentheses and speak of “the sequence a_1, a_2, a_3 ” or “the sequence a_0, a_1, a_2, \dots .”

Examples of Sequences

The “ $\subseteq S$ ” part of the sequence notation $(a_n)_{n \in I} \subseteq S$ indicates that the sequence members belong to S ;

Examples of Sequences

The “ $\subseteq S$ ” part of the sequence notation $(a_n)_{n \in I} \subseteq S$ indicates that the sequence members belong to S ; *i.e.* that the range of the sequence function $a : I \rightarrow S$ is a subset of its codomain S .

Examples of Sequences

The “ $\subseteq S$ ” part of the sequence notation $(a_n)_{n \in I} \subseteq S$ indicates that the sequence members belong to S ; *i.e.* that the range of the sequence function $a : I \rightarrow S$ is a subset of its codomain S .

The sequence *itself* is **not** a subset of S , since it is not a *set*.

Examples of Sequences

The “ $\subseteq S$ ” part of the sequence notation $(a_n)_{n \in I} \subseteq S$ indicates that the sequence members belong to S ; *i.e.* that the range of the sequence function $a : I \rightarrow S$ is a subset of its codomain S .

The sequence *itself* is **not** a subset of S , since it is not a *set*.

Examples:

1. $(a_n)_{n \in \mathbb{N}} \subseteq \mathbb{Q}$ sequence of interests rates. a_n is interest rate at time n .

Examples of Sequences

The “ $\subseteq S$ ” part of the sequence notation $(a_n)_{n \in I} \subseteq S$ indicates that the sequence members belong to S ; *i.e.* that the range of the sequence function $a : I \rightarrow S$ is a subset of its codomain S .

The sequence *itself* is **not** a subset of S , since it is not a *set*.

Examples:

1. $(a_n)_{n \in \mathbb{N}} \subseteq \mathbb{Q}$ sequence of interests rates. a_n is interest rate at time n .
2. $(p_n)_{n \in \mathbb{N}} \subseteq \mathbb{N}^* \times \mathbb{N}^* \times \mathbb{N}^*$ sequence of states of an ecosystem.
 $p_n = (r_n, s_n, t_n)$: population of species r, s, t at time n .

Examples of Sequences

The “ $\subseteq S$ ” part of the sequence notation $(a_n)_{n \in I} \subseteq S$ indicates that the sequence members belong to S ; i.e. that the range of the sequence function $a : I \rightarrow S$ is a subset of its codomain S .

The sequence *itself* is **not** a subset of S , since it is not a *set*.

Examples:

1. $(a_n)_{n \in \mathbb{N}} \subseteq \mathbb{Q}$ sequence of interests rates. a_n is interest rate at time n .
2. $(p_n)_{n \in \mathbb{N}} \subseteq \mathbb{N}^* \times \mathbb{N}^* \times \mathbb{N}^*$ sequence of states of an ecosystem.
 $p_n = (r_n, s_n, t_n)$: population of species r, s, t at time n .
3. U set of users. $(u_n)_{n \in \{1,2,3,4,5\}} \subseteq U$: a list of 5 users.

Describing sequences

An **explicit definition** of a sequence is a formula for a_n .

Describing sequences

An **explicit definition** of a sequence is a formula for a_n .

Examples:

1. $\forall n \in \mathbb{N} \quad a_n = 2^n.$

Describing sequences

An **explicit definition** of a sequence is a formula for a_n .

Examples:

1. $\forall n \in \mathbb{N} \ a_n = 2^n. \ (a_n)_{n \in \mathbb{N}} = 2, 4, 8, 16, \dots,$

Describing sequences

An **explicit definition** of a sequence is a formula for a_n .

Examples:

1. $\forall n \in \mathbb{N} \ a_n = 2^n$. $(a_n)_{n \in \mathbb{N}} = 2, 4, 8, 16, \dots$,
2. $a_1 = \text{Pierre}$, $a_2 = \text{Julie}$, $a_3 = \text{Paul}$.
 $(a_n)_{n \in \{1,2,3\}} = (\text{Pierre}, \text{Julie}, \text{Paul})$.

Describing sequences

An **explicit definition** of a sequence is a formula for a_n .

Examples:

1. $\forall n \in \mathbb{N} \ a_n = 2^n$. $(a_n)_{n \in \mathbb{N}} = 2, 4, 8, 16, \dots$,
2. $a_1 = \text{Pierre}$, $a_2 = \text{Julie}$, $a_3 = \text{Paul}$.
 $(a_n)_{n \in \{1,2,3\}} = (\text{Pierre}, \text{Julie}, \text{Paul})$.

An **implicit definition** of a sequence comprises starting value(s) and a relationship between the a_n 's.

Describing sequences

An **explicit definition** of a sequence is a formula for a_n .

Examples:

1. $\forall n \in \mathbb{N} \quad a_n = 2^n. \quad (a_n)_{n \in \mathbb{N}} = 2, 4, 8, 16, \dots,$
2. $a_1 = \text{Pierre}, a_2 = \text{Julie}, a_3 = \text{Paul}.$
 $(a_n)_{n \in \{1,2,3\}} = (\text{Pierre}, \text{Julie}, \text{Paul}).$

An **implicit definition** of a sequence comprises starting value(s) and a relationship between the a_n 's.

Examples:

1.
$$\begin{cases} a_{n+1} = 2a_n, \\ a_1 = 2. \end{cases}$$

Describing sequences

An **explicit definition** of a sequence is a formula for a_n .

Examples:

1. $\forall n \in \mathbb{N} \quad a_n = 2^n. \quad (a_n)_{n \in \mathbb{N}} = 2, 4, 8, 16, \dots,$
2. $a_1 = \text{Pierre}, a_2 = \text{Julie}, a_3 = \text{Paul}.$
 $(a_n)_{n \in \{1,2,3\}} = (\text{Pierre}, \text{Julie}, \text{Paul}).$

An **implicit definition** of a sequence comprises starting value(s) and a relationship between the a_n 's.

Examples:

1.
$$\begin{array}{ll} & a_1=2 \\ \left\{ \begin{array}{l} a_{n+1} = 2a_n, \\ a_1 = 2. \end{array} \right. & \begin{array}{l} a_2=2(2) = 4 \\ a_3=2(4) = 8 \\ \dots\dots \end{array} \end{array}$$

Describing sequences

An **explicit definition** of a sequence is a formula for a_n .

Examples:

1. $\forall n \in \mathbb{N} \quad a_n = 2^n. \quad (a_n)_{n \in \mathbb{N}} = 2, 4, 8, 16, \dots,$
2. $a_1 = \text{Pierre}, a_2 = \text{Julie}, a_3 = \text{Paul}.$
 $(a_n)_{n \in \{1,2,3\}} = (\text{Pierre}, \text{Julie}, \text{Paul}).$

An **implicit definition** of a sequence comprises starting value(s) and a relationship between the a_n 's.

Examples:

1.
$$\begin{array}{lll} & a_1=2 & \\ \left\{ \begin{array}{l} a_{n+1} = 2a_n, \\ a_1 = 2. \end{array} \right. & \begin{array}{l} a_2=2(2) = 4 \\ a_3=2(4) = 8 \\ \dots\dots \end{array} & (a_n)_{n \in \mathbb{N}} = 2, 4, 8, 16, \dots, \end{array}$$

Describing sequences

An **explicit definition** of a sequence is a formula for a_n .

Examples:

1. $\forall n \in \mathbb{N} \quad a_n = 2^n. \quad (a_n)_{n \in \mathbb{N}} = 2, 4, 8, 16, \dots,$

2. $a_1 = \text{Pierre}, a_2 = \text{Julie}, a_3 = \text{Paul}.$

$$(a_n)_{n \in \{1,2,3\}} = (\text{Pierre}, \text{Julie}, \text{Paul}).$$

An **implicit definition** of a sequence comprises starting value(s) and a relationship between the a_n 's.

Examples:

1.
$$\begin{cases} a_{n+1} = 2a_n, & a_1 = 2 \\ a_1 = 2. \end{cases} \quad \begin{matrix} a_2 = 2(2) = 4 \\ a_3 = 2(4) = 8 \end{matrix} \quad (a_n)_{n \in \mathbb{N}} = 2, 4, 8, 16, \dots,$$

.....

2.
$$\begin{cases} a_{n+1} = -a_n + a_{n-1}, \\ a_2 = 1, \\ a_1 = 0. \end{cases}$$

Describing sequences

An **explicit definition** of a sequence is a formula for a_n .

Examples:

1. $\forall n \in \mathbb{N} \quad a_n = 2^n. \quad (a_n)_{n \in \mathbb{N}} = 2, 4, 8, 16, \dots,$

2. $a_1 = \text{Pierre}, a_2 = \text{Julie}, a_3 = \text{Paul}.$

$$(a_n)_{n \in \{1,2,3\}} = (\text{Pierre}, \text{Julie}, \text{Paul}).$$

An **implicit definition** of a sequence comprises starting value(s) and a relationship between the a_n 's.

Examples:

1.
$$\begin{cases} a_{n+1} = 2a_n, & a_1 = 2 \\ a_1 = 2. \end{cases} \quad \begin{array}{l} a_2 = 2(2) = 4 \\ a_3 = 2(4) = 8 \\ \dots\dots \end{array} \quad (a_n)_{n \in \mathbb{N}} = 2, 4, 8, 16, \dots,$$

2.
$$\begin{cases} a_{n+1} = -a_n + a_{n-1}, & a_1 = 0 \\ a_2 = 1, & a_2 = 1 \\ a_1 = 0. \end{cases} \quad \begin{array}{l} a_3 = -1 + 0 = -1 \\ a_4 = -(-1) + 1 = 2 \\ a_5 = -2 + (-1) = -3 \\ \dots\dots\dots \end{array}$$

Proofs about sequences

Mathematical induction

Let $P(n)$ be a predicate with variable $n \in \mathbb{N}$.

Mathematical induction

Let $P(n)$ be a predicate with variable $n \in \mathbb{N}$.

How to prove that $\forall n \in \mathbb{N} \ P(n)$?

Mathematical induction

Let $P(n)$ be a predicate with variable $n \in \mathbb{N}$.

How to prove that $\forall n \in \mathbb{N} \ P(n)$?

Proof of $\forall n \in \mathbb{N} \ P(n)$ by mathematical induction:

- Prove $P(1)$.

Mathematical induction

Let $P(n)$ be a predicate with variable $n \in \mathbb{N}$.

How to prove that $\forall n \in \mathbb{N} \ P(n)$?

Proof of $\forall n \in \mathbb{N} \ P(n)$ by mathematical induction:

- Prove $P(1)$.
- Let $n \in \mathbb{N}$. Treat n as temporarily fixed and then:

Mathematical induction

Let $P(n)$ be a predicate with variable $n \in \mathbb{N}$.

How to prove that $\forall n \in \mathbb{N} \ P(n)$?

Proof of $\forall n \in \mathbb{N} \ P(n)$ by mathematical induction:

- Prove $P(1)$.
- Let $n \in \mathbb{N}$. Treat n as temporarily fixed and then:
Assume $P(1) \wedge P(2) \wedge P(3) \wedge \dots \wedge P(n)$.

Mathematical induction

Let $P(n)$ be a predicate with variable $n \in \mathbb{N}$.

How to prove that $\forall n \in \mathbb{N} \ P(n)$?

Proof of $\forall n \in \mathbb{N} \ P(n)$ by mathematical induction:

- Prove $P(1)$.
- Let $n \in \mathbb{N}$. Treat n as temporarily fixed and then:
Assume $P(1) \wedge P(2) \wedge P(3) \wedge \dots \wedge P(n)$.
- Prove $P(n+1)$.

Mathematical induction

Let $P(n)$ be a predicate with variable $n \in \mathbb{N}$.

How to prove that $\forall n \in \mathbb{N} \ P(n)$?

Proof of $\forall n \in \mathbb{N} \ P(n)$ by mathematical induction:

- Prove $P(1)$.
- Let $n \in \mathbb{N}$. Treat n as temporarily fixed and then:
Assume $P(1) \wedge P(2) \wedge P(3) \wedge \dots \wedge P(n)$.
- Prove $P(n+1)$.

This is also known as **strong** mathematical induction.

Mathematical induction

Let $P(n)$ be a predicate with variable $n \in \mathbb{N}$.

How to prove that $\forall n \in \mathbb{N} \ P(n)$?

Proof of $\forall n \in \mathbb{N} \ P(n)$ by mathematical induction:

- Prove $P(1)$.
- Let $n \in \mathbb{N}$. Treat n as temporarily fixed and then:
Assume $P(1) \wedge P(2) \wedge P(3) \wedge \dots \wedge P(n)$.
- Prove $P(n+1)$.

This is also known as **strong** mathematical induction.

This works because:

$P(1)$ is true.

Mathematical induction

Let $P(n)$ be a predicate with variable $n \in \mathbb{N}$.

How to prove that $\forall n \in \mathbb{N} \ P(n)$?

Proof of $\forall n \in \mathbb{N} \ P(n)$ by mathematical induction:

- Prove $P(1)$.
- Let $n \in \mathbb{N}$. Treat n as temporarily fixed and then:
Assume $P(1) \wedge P(2) \wedge P(3) \wedge \dots \wedge P(n)$.
- Prove $P(n+1)$.

This is also known as **strong** mathematical induction.

This works because:

$P(1)$ is true.

$P(1) \implies P(2)$, so $P(2)$ is true.

Mathematical induction

Let $P(n)$ be a predicate with variable $n \in \mathbb{N}$.

How to prove that $\forall n \in \mathbb{N} \ P(n)$?

Proof of $\forall n \in \mathbb{N} \ P(n)$ by mathematical induction:

- Prove $P(1)$.
- Let $n \in \mathbb{N}$. Treat n as temporarily fixed and then:
Assume $P(1) \wedge P(2) \wedge P(3) \wedge \dots \wedge P(n)$.
- Prove $P(n+1)$.

This is also known as **strong** mathematical induction.

This works because:

$P(1)$ is true.

$P(1) \implies P(2)$, so $P(2)$ is true.

$P(1) \wedge P(2) \implies P(3)$, so $P(3)$ is true.

Mathematical induction

Let $P(n)$ be a predicate with variable $n \in \mathbb{N}$.

How to prove that $\forall n \in \mathbb{N} \ P(n)$?

Proof of $\forall n \in \mathbb{N} \ P(n)$ by mathematical induction:

- Prove $P(1)$.
- Let $n \in \mathbb{N}$. Treat n as temporarily fixed and then:
Assume $P(1) \wedge P(2) \wedge P(3) \wedge \dots \wedge P(n)$.
- Prove $P(n+1)$.

This is also known as **strong** mathematical induction.

This works because:

$P(1)$ is true.

$P(1) \implies P(2)$, so $P(2)$ is true.

$P(1) \wedge P(2) \implies P(3)$, so $P(3)$ is true.

$P(1) \wedge P(2) \wedge P(3) \implies P(4)$, so $P(4)$ is true too.

Mathematical induction

Let $P(n)$ be a predicate with variable $n \in \mathbb{N}$.

How to prove that $\forall n \in \mathbb{N} \ P(n)$?

Proof of $\forall n \in \mathbb{N} \ P(n)$ by mathematical induction:

- Prove $P(1)$.
- Let $n \in \mathbb{N}$. Treat n as temporarily fixed and then:
Assume $P(1) \wedge P(2) \wedge P(3) \wedge \dots \wedge P(n)$.
- Prove $P(n+1)$.

This is also known as **strong** mathematical induction.

This works because:

$P(1)$ is true.

$P(1) \implies P(2)$, so $P(2)$ is true.

$P(1) \wedge P(2) \implies P(3)$, so $P(3)$ is true.

$P(1) \wedge P(2) \wedge P(3) \implies P(4)$, so $P(4)$ is true too.

Continuing to argue in this manner gives $P(n)$ for all $n \in \mathbb{N}$.

From implicit to explicit definitions; Example 1

A sequence is defined implicitly by
$$\begin{cases} a_{n+1} = 3a_n & \forall n \in \mathbb{N}, \\ a_1 = 3. \end{cases}$$

From implicit to explicit definitions; Example 1

A sequence is defined implicitly by
$$\begin{cases} a_{n+1} = 3a_n & \forall n \in \mathbb{N}, \\ a_1 = 3. \end{cases}$$

Can we get an explicit definition?

From implicit to explicit definitions; Example 1

A sequence is defined implicitly by
$$\begin{cases} a_{n+1} = 3a_n & \forall n \in \mathbb{N}, \\ a_1 = 3. \end{cases}$$

Can we get an explicit definition? First generate some values:

$$a_1 = 3,$$

From implicit to explicit definitions; Example 1

A sequence is defined implicitly by
$$\begin{cases} a_{n+1} = 3a_n & \forall n \in \mathbb{N}, \\ a_1 = 3. \end{cases}$$

Can we get an explicit definition? First generate some values:

$$a_1 = 3, a_2 = 9,$$

From implicit to explicit definitions; Example 1

A sequence is defined implicitly by
$$\begin{cases} a_{n+1} = 3a_n & \forall n \in \mathbb{N}, \\ a_1 = 3. \end{cases}$$

Can we get an explicit definition? First generate some values:

$$a_1 = 3, a_2 = 9, a_3 = 27,$$

From implicit to explicit definitions; Example 1

A sequence is defined implicitly by
$$\begin{cases} a_{n+1} = 3a_n & \forall n \in \mathbb{N}, \\ a_1 = 3. \end{cases}$$

Can we get an explicit definition? First generate some values:

$a_1=3$, $a_2=9$, $a_3=27$, $a_4=81$,...

From implicit to explicit definitions; Example 1

A sequence is defined implicitly by
$$\begin{cases} a_{n+1} = 3a_n & \forall n \in \mathbb{N}, \\ a_1 = 3. \end{cases}$$

Can we get an explicit definition? First generate some values:

$a_1=3, a_2=9, a_3=27, a_4=81, \dots$

Claim: $\forall n \in \mathbb{N} \ a_n = 3^n$.

From implicit to explicit definitions; Example 1

A sequence is defined implicitly by
$$\begin{cases} a_{n+1} = 3a_n & \forall n \in \mathbb{N}, \\ a_1 = 3. \end{cases}$$

Can we get an explicit definition? First generate some values:

$a_1=3, a_2=9, a_3=27, a_4=81, \dots$

Claim: $\forall n \in \mathbb{N} \ a_n = 3^n$.

Proof by mathematical induction that the claim is correct:

From implicit to explicit definitions; Example 1

A sequence is defined implicitly by
$$\begin{cases} a_{n+1} = 3a_n & \forall n \in \mathbb{N}, \\ a_1 = 3. \end{cases}$$

Can we get an explicit definition? First generate some values:

$a_1=3, a_2=9, a_3=27, a_4=81, \dots$

Claim: $\forall n \in \mathbb{N} \ a_n = 3^n$.

Proof by mathematical induction that the claim is correct:

Basis step: For $n=1$, formula gives $a_1 = 3^1 = 3$, agreeing with the implicit definition.

From implicit to explicit definitions; Example 1

A sequence is defined implicitly by
$$\begin{cases} a_{n+1} = 3a_n & \forall n \in \mathbb{N}, \\ a_1 = 3. \end{cases}$$

Can we get an explicit definition? First generate some values:

$a_1=3, a_2=9, a_3=27, a_4=81, \dots$

Claim: $\forall n \in \mathbb{N} \ a_n = 3^n$.

Proof by mathematical induction that the claim is correct:

Basis step: For $n=1$, formula gives $a_1 = 3^1 = 3$, agreeing with the implicit definition.

Inductive step: Assume the formula is correct for up to and including some fixed n .

From implicit to explicit definitions; Example 1

A sequence is defined implicitly by
$$\begin{cases} a_{n+1} = 3a_n & \forall n \in \mathbb{N}, \\ a_1 = 3. \end{cases}$$

Can we get an explicit definition? First generate some values:

$$a_1=3, a_2=9, a_3=27, a_4=81, \dots$$

$$\textbf{Claim: } \forall n \in \mathbb{N} \ a_n = 3^n.$$

Proof by mathematical induction that the claim is correct:

Basis step: For $n=1$, formula gives $a_1 = 3^1 = 3$, agreeing with the implicit definition.

Inductive step: Assume the formula is correct for up to and including some fixed n . Then

$$a_{n+1} = 3a_n \quad (\text{from the implicit definition})$$

From implicit to explicit definitions; Example 1

A sequence is defined implicitly by
$$\begin{cases} a_{n+1} = 3a_n & \forall n \in \mathbb{N}, \\ a_1 = 3. \end{cases}$$

Can we get an explicit definition? First generate some values:

$$a_1=3, a_2=9, a_3=27, a_4=81, \dots$$

Claim: $\forall n \in \mathbb{N} \ a_n = 3^n$.

Proof by mathematical induction that the claim is correct:

Basis step: For $n=1$, formula gives $a_1 = 3^1 = 3$, agreeing with the implicit definition.

Inductive step: Assume the formula is correct for up to and including some fixed n . Then

$$\begin{aligned} a_{n+1} &= 3a_n \quad (\text{from the implicit definition}) \\ &= 3(3^n) \quad (\text{by the inductive assumption}) \end{aligned}$$

From implicit to explicit definitions; Example 1

A sequence is defined implicitly by
$$\begin{cases} a_{n+1} = 3a_n & \forall n \in \mathbb{N}, \\ a_1 = 3. \end{cases}$$

Can we get an explicit definition? First generate some values:

$$a_1=3, a_2=9, a_3=27, a_4=81, \dots$$

$$\textbf{Claim: } \forall n \in \mathbb{N} \ a_n = 3^n.$$

Proof by mathematical induction that the claim is correct:

Basis step: For $n=1$, formula gives $a_1=3^1=3$, agreeing with the implicit definition.

Inductive step: Assume the formula is correct for up to and including some fixed n . Then

$$\begin{aligned} a_{n+1} &= 3a_n && \text{(from the implicit definition)} \\ &= 3(3^n) && \text{(by the inductive assumption)} \\ &= 3^{n+1} \end{aligned}$$

From implicit to explicit definitions; Example 1

A sequence is defined implicitly by
$$\begin{cases} a_{n+1} = 3a_n & \forall n \in \mathbb{N}, \\ a_1 = 3. \end{cases}$$

Can we get an explicit definition? First generate some values:

$$a_1=3, a_2=9, a_3=27, a_4=81, \dots$$

$$\textbf{Claim: } \forall n \in \mathbb{N} \ a_n = 3^n.$$

Proof by mathematical induction that the claim is correct:

Basis step: For $n=1$, formula gives $a_1=3^1=3$, agreeing with the implicit definition.

Inductive step: Assume the formula is correct for up to and including some fixed n . Then

$$\begin{aligned} a_{n+1} &= 3a_n \quad (\text{from the implicit definition}) \\ &= 3(3^n) \quad (\text{by the inductive assumption}) \\ &= 3^{n+1} \end{aligned}$$

and so the formula is also correct for $n+1$.

Sum and products of terms

Terms of a sequence can be summed: $a_1 + a_2 + a_3 + \dots$

Sum and products of terms

Terms of a sequence can be summed: $a_1 + a_2 + a_3 + \dots$ or multiplied:
 $a_1 \times a_2 \times a_3 \times \dots$

Sum and products of terms

Terms of a sequence can be summed: $a_1 + a_2 + a_3 + \dots$ or multiplied: $a_1 \times a_2 \times a_3 \times \dots$. We use the special notation

$$\sum_{n=1}^k a_n = a_1 + a_2 + a_3 + \dots + a_k,$$

Sum and products of terms

Terms of a sequence can be summed: $a_1 + a_2 + a_3 + \dots$ or multiplied: $a_1 \times a_2 \times a_3 \times \dots$. We use the special notation

$$\sum_{n=1}^k a_n = a_1 + a_2 + a_3 + \dots + a_k, \quad \prod_{n=1}^k a_n = a_1 \times a_2 \times a_3 \times \dots \times a_k.$$

Sum and products of terms

Terms of a sequence can be summed: $a_1 + a_2 + a_3 + \dots$ or multiplied: $a_1 \times a_2 \times a_3 \times \dots$. We use the special notation

$$\sum_{n=1}^k a_n = a_1 + a_2 + a_3 + \dots + a_k, \quad \prod_{n=1}^k a_n = a_1 \times a_2 \times a_3 \times \dots \times a_k.$$

Examples: (1) $\sum_{n=1}^{10} n = 1 + 2 + 3 + 4 + \dots + 9 + 10 = 55.$

Sum and products of terms

Terms of a sequence can be summed: $a_1 + a_2 + a_3 + \dots$ or multiplied: $a_1 \times a_2 \times a_3 \times \dots$. We use the special notation

$$\sum_{n=1}^k a_n = a_1 + a_2 + a_3 + \dots + a_k, \quad \prod_{n=1}^k a_n = a_1 \times a_2 \times a_3 \times \dots \times a_k.$$

Examples: (1) $\sum_{n=1}^{10} n = 1 + 2 + 3 + 4 + \dots + 9 + 10 = 55.$

(2) $\sum_{n=0}^7 2^n = 1 + 2 + 4 + 8 + \dots + 128 = 255.$

Sum and products of terms

Terms of a sequence can be summed: $a_1 + a_2 + a_3 + \dots$ or multiplied: $a_1 \times a_2 \times a_3 \times \dots$. We use the special notation

$$\sum_{n=1}^k a_n = a_1 + a_2 + a_3 + \dots + a_k, \quad \prod_{n=1}^k a_n = a_1 \times a_2 \times a_3 \times \dots \times a_k.$$

Examples:

- (1) $\sum_{n=1}^{10} n = 1 + 2 + 3 + 4 + \dots + 9 + 10 = 55.$
- (2) $\sum_{n=0}^7 2^n = 1 + 2 + 4 + 8 + \dots + 128 = 255.$
- (3) $\prod_{n=1}^5 n = 1 \times 2 \times 3 \times 4 \times 5 = 5! = 120.$

Sum and products of terms

Terms of a sequence can be summed: $a_1 + a_2 + a_3 + \dots$ or multiplied: $a_1 \times a_2 \times a_3 \times \dots$. We use the special notation

$$\sum_{n=1}^k a_n = a_1 + a_2 + a_3 + \dots + a_k, \quad \prod_{n=1}^k a_n = a_1 \times a_2 \times a_3 \times \dots \times a_k.$$

Examples: (1) $\sum_{n=1}^{10} n = 1 + 2 + 3 + 4 + \dots + 9 + 10 = 55.$

(2) $\sum_{n=0}^7 2^n = 1 + 2 + 4 + 8 + \dots + 128 = 255.$

(3) $\prod_{n=1}^5 n = 1 \times 2 \times 3 \times 4 \times 5 = 5! = 120.$

(4) $\prod_{n=1}^8 n^2 = 4 \times 9 \times 16 \times \dots \times 64 = 1\,625\,702\,400.$

Given a sequence $(a_n)_{n \in I} \subseteq \mathbb{Q}$, we define a **series** as $\sum_{n \in I} a_n$.

Given a sequence we can also consider **partial sums** of the series, which is just a series associated to some 'subsequence'.

We will now consider a few important examples of sequences and series:

- Geometric Sequence and Series
- Arithmetic Sequence and Series
- Mixed Geometric-Arithmetic Sequence

how we can calculate their terms and their associated series.

Geometric Sequences and Series

Slide 6 is a special case of a

Geometric Sequence	
Implicit Definition	Explicit Definition
$a_k = a$ (a is the first term) $a_{n+1} = ra_n, \forall n \geq k$ (r is the common ratio)	$\forall n \geq k$ $a_n = ar^{n-k}$

Geometric Sequences and Series

Slide 6 is a special case of a

Geometric Sequence	
Implicit Definition	Explicit Definition
$a_k = a$ (a is the first term) $a_{n+1} = ra_n, \forall n \geq k$ (r is the common ratio)	$\forall n \geq k$ $a_n = ar^{n-k}$

When terms of a sequence are summed, we get a **series**. In particular

Geometric Series	
Series of N terms	Sum of N terms
$\sum_{n=k}^{k+(N-1)} ar^{n-k} = a + ar + \dots + ar^{N-1}$ [Usually $k=0$ or $k=1$.]	$\begin{cases} \frac{a(1-r^N)}{(1-r)} & \text{if } r \neq 1 \\ Na & \text{if } r = 1 \end{cases}$

Geometric Sequences and Series

Slide 6 is a special case of a

Geometric Sequence	
Implicit Definition	Explicit Definition
$a_k = a$ (a is the first term) $a_{n+1} = ra_n, \forall n \geq k$ (r is the common ratio)	$\forall n \geq k$ $a_n = ar^{n-k}$

When terms of a sequence are summed, we get a **series**. In particular

Geometric Series	
Series of N terms	Sum of N terms
$\sum_{n=k}^{k+(N-1)} ar^{n-k} = a + ar + \dots + ar^{N-1}$ [Usually $k=0$ or $k=1$.]	$\begin{cases} \frac{a(1-r^N)}{(1-r)} & \text{if } r \neq 1 \\ Na & \text{if } r = 1 \end{cases}$

Example: $k = 0, a = 6, r = \frac{1}{2}, N = 5$:

Geometric Sequences and Series

Slide 6 is a special case of a

Geometric Sequence	
Implicit Definition	Explicit Definition
$a_k = a$ (a is the first term) $a_{n+1} = ra_n, \forall n \geq k$ (r is the common ratio)	$\forall n \geq k$ $a_n = ar^{n-k}$

When terms of a sequence are summed, we get a **series**. In particular

Geometric Series	
Series of N terms	Sum of N terms
$\sum_{n=k}^{k+(N-1)} ar^{n-k} = a + ar + \dots + ar^{N-1}$ [Usually $k=0$ or $k=1$.]	$\begin{cases} \frac{a(1-r^N)}{(1-r)} & \text{if } r \neq 1 \\ Na & \text{if } r = 1 \end{cases}$

Example: $k = 0, a = 6, r = \frac{1}{2}, N = 5$:

$$6 + 3 + \frac{3}{2} + \frac{3}{4} + \frac{3}{8} = \sum_{n=0}^4 6\left(\frac{1}{2}\right)^n$$

Geometric Sequences and Series

Slide 6 is a special case of a

Geometric Sequence	
Implicit Definition	Explicit Definition
$a_k = a$ (a is the first term) $a_{n+1} = ra_n, \forall n \geq k$ (r is the common ratio)	$\forall n \geq k$ $a_n = ar^{n-k}$

When terms of a sequence are summed, we get a **series**. In particular

Geometric Series	
Series of N terms	Sum of N terms
$\sum_{n=k}^{k+(N-1)} ar^{n-k} = a + ar + \dots + ar^{N-1}$ [Usually $k=0$ or $k=1$.]	$\begin{cases} \frac{a(1-r^N)}{(1-r)} & \text{if } r \neq 1 \\ Na & \text{if } r = 1 \end{cases}$

Example: $k = 0, a = 6, r = \frac{1}{2}, N = 5$:

$$6 + 3 + \frac{3}{2} + \frac{3}{4} + \frac{3}{8} = \sum_{n=0}^4 6\left(\frac{1}{2}\right)^n = \frac{6(1-2^{-5})}{1-2^{-1}}$$

Geometric Sequences and Series

Slide 6 is a special case of a

Geometric Sequence	
Implicit Definition	Explicit Definition
$a_k = a$ (a is the first term) $a_{n+1} = ra_n, \forall n \geq k$ (r is the common ratio)	$\forall n \geq k$ $a_n = ar^{n-k}$

When terms of a sequence are summed, we get a **series**. In particular

Geometric Series	
Series of N terms	Sum of N terms
$\sum_{n=k}^{k+(N-1)} ar^{n-k} = a + ar + \dots + ar^{N-1}$ [Usually $k=0$ or $k=1$.]	$\begin{cases} \frac{a(1-r^N)}{(1-r)} & \text{if } r \neq 1 \\ Na & \text{if } r = 1 \end{cases}$

Example: $k = 0, a = 6, r = \frac{1}{2}, N = 5$:

$$6 + 3 + \frac{3}{2} + \frac{3}{4} + \frac{3}{8} = \sum_{n=0}^4 6\left(\frac{1}{2}\right)^n = \frac{6(1-2^{-5})}{1-2^{-1}} = \frac{6(32-1)}{32} \cdot \frac{2}{1} = \frac{93}{8} = 11\frac{5}{8}$$

From implicit to explicit definitions; Example 2

A sequence is defined implicitly by
$$\begin{cases} b_{n+1} = b_n + 5 & \forall n \in \mathbb{N}, \\ b_1 = 0. \end{cases}$$

From implicit to explicit definitions; Example 2

A sequence is defined implicitly by
$$\begin{cases} b_{n+1} = b_n + 5 & \forall n \in \mathbb{N}, \\ b_1 = 0. \end{cases}$$

Can we get an explicit definition?

From implicit to explicit definitions; Example 2

A sequence is defined implicitly by
$$\begin{cases} b_{n+1} = b_n + 5 & \forall n \in \mathbb{N}, \\ b_1 = 0. \end{cases}$$

Can we get an explicit definition? First generate some values:

$b_1=0$, $b_2=5$, $b_3=10$, $b_4=15,..$

From implicit to explicit definitions; Example 2

A sequence is defined implicitly by
$$\begin{cases} b_{n+1} = b_n + 5 & \forall n \in \mathbb{N}, \\ b_1 = 0. \end{cases}$$

Can we get an explicit definition? First generate some values:

$b_1=0, b_2=5, b_3=10, b_4=15, \dots$

Claim: $\forall n \in \mathbb{N} \ b_n = 5(n-1).$

From implicit to explicit definitions; Example 2

A sequence is defined implicitly by
$$\begin{cases} b_{n+1} = b_n + 5 & \forall n \in \mathbb{N}, \\ b_1 = 0. \end{cases}$$

Can we get an explicit definition? First generate some values:

$b_1=0, b_2=5, b_3=10, b_4=15, \dots$

Claim: $\forall n \in \mathbb{N} \ b_n = 5(n-1).$

Proof by mathematical induction that the claim is correct:

From implicit to explicit definitions; Example 2

A sequence is defined implicitly by
$$\begin{cases} b_{n+1} = b_n + 5 & \forall n \in \mathbb{N}, \\ b_1 = 0. \end{cases}$$

Can we get an explicit definition? First generate some values:

$b_1=0, b_2=5, b_3=10, b_4=15, \dots$

Claim: $\forall n \in \mathbb{N} \ b_n = 5(n-1).$

Proof by mathematical induction that the claim is correct:

Basis step: For $n=1$, formula gives $b_1=0$, agreeing with the definition.

From implicit to explicit definitions; Example 2

A sequence is defined implicitly by
$$\begin{cases} b_{n+1} = b_n + 5 & \forall n \in \mathbb{N}, \\ b_1 = 0. \end{cases}$$

Can we get an explicit definition? First generate some values:

$b_1=0, b_2=5, b_3=10, b_4=15, \dots$

Claim: $\forall n \in \mathbb{N} \ b_n = 5(n-1).$

Proof by mathematical induction that the claim is correct:

Basis step: For $n=1$, formula gives $b_1=0$, agreeing with the definition.

Inductive step: Assume the formula is correct for up to and including some fixed n .

From implicit to explicit definitions; Example 2

A sequence is defined implicitly by
$$\begin{cases} b_{n+1} = b_n + 5 & \forall n \in \mathbb{N}, \\ b_1 = 0. \end{cases}$$

Can we get an explicit definition? First generate some values:

$b_1=0, b_2=5, b_3=10, b_4=15, \dots$

Claim: $\forall n \in \mathbb{N} \ b_n = 5(n-1).$

Proof by mathematical induction that the claim is correct:

Basis step: For $n=1$, formula gives $b_1=0$, agreeing with the definition.

Inductive step: Assume the formula is correct for up to and including some fixed n . Then

$$b_{n+1} = b_n + 5 \quad (\text{from the implicit definition})$$

From implicit to explicit definitions; Example 2

A sequence is defined implicitly by
$$\begin{cases} b_{n+1} = b_n + 5 & \forall n \in \mathbb{N}, \\ b_1 = 0. \end{cases}$$

Can we get an explicit definition? First generate some values:

$$b_1=0, b_2=5, b_3=10, b_4=15, \dots$$

$$\textbf{Claim: } \forall n \in \mathbb{N} \ b_n = 5(n-1).$$

Proof by mathematical induction that the claim is correct:

Basis step: For $n=1$, formula gives $b_1=0$, agreeing with the definition.

Inductive step: Assume the formula is correct for up to and including some fixed n . Then

$$\begin{aligned} b_{n+1} &= b_n + 5 && \text{(from the implicit definition)} \\ &= 5(n-1) + 5 && \text{(by the inductive assumption)} \end{aligned}$$

From implicit to explicit definitions; Example 2

A sequence is defined implicitly by
$$\begin{cases} b_{n+1} = b_n + 5 & \forall n \in \mathbb{N}, \\ b_1 = 0. \end{cases}$$

Can we get an explicit definition? First generate some values:

$b_1=0, b_2=5, b_3=10, b_4=15, \dots$

Claim: $\forall n \in \mathbb{N} \ b_n = 5(n-1).$

Proof by mathematical induction that the claim is correct:

Basis step: For $n=1$, formula gives $b_1=0$, agreeing with the definition.

Inductive step: Assume the formula is correct for up to and including some fixed n . Then

$$\begin{aligned} b_{n+1} &= b_n + 5 && \text{(from the implicit definition)} \\ &= 5(n-1) + 5 && \text{(by the inductive assumption)} \\ &= 5n \end{aligned}$$

From implicit to explicit definitions; Example 2

A sequence is defined implicitly by
$$\begin{cases} b_{n+1} = b_n + 5 & \forall n \in \mathbb{N}, \\ b_1 = 0. \end{cases}$$

Can we get an explicit definition? First generate some values:

$b_1=0, b_2=5, b_3=10, b_4=15, \dots$

Claim: $\forall n \in \mathbb{N} \ b_n = 5(n-1).$

Proof by mathematical induction that the claim is correct:

Basis step: For $n=1$, formula gives $b_1=0$, agreeing with the definition.

Inductive step: Assume the formula is correct for up to and including some fixed n . Then

$$\begin{aligned} b_{n+1} &= b_n + 5 && \text{(from the implicit definition)} \\ &= 5(n-1) + 5 && \text{(by the inductive assumption)} \\ &= 5n = 5((n+1)-1) \end{aligned}$$

From implicit to explicit definitions; Example 2

A sequence is defined implicitly by
$$\begin{cases} b_{n+1} = b_n + 5 & \forall n \in \mathbb{N}, \\ b_1 = 0. \end{cases}$$

Can we get an explicit definition? First generate some values:

$b_1=0, b_2=5, b_3=10, b_4=15, \dots$

Claim: $\forall n \in \mathbb{N} \ b_n = 5(n-1).$

Proof by mathematical induction that the claim is correct:

Basis step: For $n=1$, formula gives $b_1=0$, agreeing with the definition.

Inductive step: Assume the formula is correct for up to and including some fixed n . Then

$$\begin{aligned} b_{n+1} &= b_n + 5 && \text{(from the implicit definition)} \\ &= 5(n-1) + 5 && \text{(by the inductive assumption)} \\ &= 5n = 5((n+1)-1) \end{aligned}$$

and so the formula is also correct for $n+1$.

Arithmetic Sequences and Series

Example 2 is a special case of

Arithmetic Sequence	
Implicit Definition	Explicit Definition
$a_k = a$ (a is the first term) $a_{n+1} = a_n + d, \forall n \geq k$ (d is the common difference)	$\forall n \geq k$ $a_n = a + (n - k)d$

Arithmetic Sequences and Series

Example 2 is a special case of

Arithmetic Sequence	
Implicit Definition	Explicit Definition
$a_k = a$ (a is the first term) $a_{n+1} = a_n + d, \forall n \geq k$ (d is the common difference)	$\forall n \geq k$ $a_n = a + (n - k)d$

When the terms of an arithmetic sequence are summed, we get:

Arithmetic Series	
Series of N terms	Sum of N terms
$\sum_{n=k}^{k+(N-1)} [a + (n-k)d] = a + (a+d) + \cdots + (a + (N-1)d)$ [Usually $k=0$ or $k=1$.]	$\frac{N}{2} [2a + (N-1)d]$ $=$ $N \left[\frac{a + (a + (N-1)d)}{2} \right]$

Arithmetic Sequences and Series

Example 2 is a special case of

Arithmetic Sequence	
Implicit Definition	Explicit Definition
$a_k = a$ (a is the first term) $a_{n+1} = a_n + d, \forall n \geq k$ (d is the common difference)	$\forall n \geq k$ $a_n = a + (n - k)d$

When the terms or an arithmetic sequence are summed, we get:

Arithmetic Series	
Series of N terms	Sum of N terms
$\sum_{n=k}^{k+(N-1)} [a + (n-k)d] = a + (a+d) + \dots + (a + (N-1)d)$ [Usually $k=0$ or $k=1$.]	$\frac{N}{2} [2a + (N-1)d]$ $=$ $N \left[\frac{a + (a + (N-1)d)}{2} \right]$

sum = number of terms times average of first and last

Arithmetic Sequences and Series

Example 2 is a special case of

Arithmetic Sequence	
Implicit Definition	Explicit Definition
$a_k = a$ (a is the first term) $a_{n+1} = a_n + d, \forall n \geq k$ (d is the common difference)	$\forall n \geq k$ $a_n = a + (n - k)d$

When the terms or an arithmetic sequence are summed, we get:

Arithmetic Series	
Series of N terms	Sum of N terms
$\sum_{n=k}^{k+(N-1)} [a + (n-k)d] = a + (a+d) + \dots + (a + (N-1)d)$ [Usually $k=0$ or $k=1$.]	$\frac{N}{2} [2a + (N-1)d]$ $=$ $N \left[\frac{a + (a + (N-1)d)}{2} \right]$

sum = number of terms times average of first and last

Example: $1 + 3 + 5 + 7 + \dots + 99 = 50 \left(\frac{1 + 99}{2} \right) = 2500$.

In this example $a = 1, d = 2, k = 0, N = 50$, check yourself!

Simplest Malthusian model in population dynamics

Exponential Growth:

Let p_n be number of individuals in a population after n years.

Simplest Malthusian model in population dynamics

Exponential Growth:

Let p_n be number of individuals in a population after n years.

Assume the population doubles every year,

Simplest Malthusian model in population dynamics

Exponential Growth:

Let p_n be number of individuals in a population after n years.

Assume the population doubles every year, *i.e.* $\forall n \in \mathbb{N}^* \quad p_{n+1} = 2p_n$.

Simplest Malthusian model in population dynamics

Exponential Growth:

Let p_n be number of individuals in a population after n years.

Assume the population doubles every year, *i.e.* $\forall n \in \mathbb{N}^* \quad p_{n+1} = 2p_n$.

Assume the current population is five million.

Simplest Malthusian model in population dynamics

Exponential Growth:

Let p_n be number of individuals in a population after n years.

Assume the population doubles every year, *i.e.* $\forall n \in \mathbb{N}^* \quad p_{n+1} = 2p_n$.

Assume the current population is five million. *i.e.* $p_0 = 5 \times 10^6$.

Simplest Malthusian model in population dynamics

Exponential Growth:

Let p_n be number of individuals in a population after n years.

Assume the population doubles every year, *i.e.* $\forall n \in \mathbb{N}^* \quad p_{n+1} = 2p_n$.

Assume the current population is five million. *i.e.* $p_0 = 5 \times 10^6$.

Questions: 1. What will the population size be in 10 years time?

Simplest Malthusian model in population dynamics

Exponential Growth:

Let p_n be number of individuals in a population after n years.

Assume the population doubles every year, *i.e.* $\forall n \in \mathbb{N}^* \quad p_{n+1} = 2p_n$.

Assume the current population is five million. *i.e.* $p_0 = 5 \times 10^6$.

- Questions:
1. What will the population size be in 10 years time?
 2. When will the population size reach 10^9 ?

Simplest Malthusian model in population dynamics

Exponential Growth:

Let p_n be number of individuals in a population after n years.

Assume the population doubles every year, i.e. $\forall n \in \mathbb{N}^* \quad p_{n+1} = 2p_n$.

Assume the current population is five million. i.e. $p_0 = 5 \times 10^6$.

- Questions:
1. What will the population size be in 10 years time?
 2. When will the population size reach 10^9 ?

We have a geometric sequence with $k = 0$, $a = 5 \times 10^6$, $r = 2$.

Simplest Malthusian model in population dynamics

Exponential Growth:

Let p_n be number of individuals in a population after n years.

Assume the population doubles every year, i.e. $\forall n \in \mathbb{N}^* \quad p_{n+1} = 2p_n$.

Assume the current population is five million. i.e. $p_0 = 5 \times 10^6$.

- Questions:
1. What will the population size be in 10 years time?
 2. When will the population size reach 10^9 ?

We have a geometric sequence with $k = 0$, $a = 5 \times 10^6$, $r = 2$. Hence:

Answers: 1. $a_{10} = 5 \times 10^6 \times 2^{10} = 5.12 \times 10^9$

Simplest Malthusian model in population dynamics

Exponential Growth:

Let p_n be number of individuals in a population after n years.

Assume the population doubles every year, i.e. $\forall n \in \mathbb{N}^* \quad p_{n+1} = 2p_n$.

Assume the current population is five million. i.e. $p_0 = 5 \times 10^6$.

- Questions:
1. What will the population size be in 10 years time?
 2. When will the population size reach 10^9 ?

We have a geometric sequence with $k = 0$, $a = 5 \times 10^6$, $r = 2$. Hence:

- Answers:
1. $a_{10} = 5 \times 10^6 \times 2^{10} = 5.12 \times 10^9$
 2. We need N so that $a_N \geq 10^9 > a_{N-1}$.

Simplest Malthusian model in population dynamics

Exponential Growth:

Let p_n be number of individuals in a population after n years.

Assume the population doubles every year, i.e. $\forall n \in \mathbb{N}^* \quad p_{n+1} = 2p_n$.

Assume the current population is five million. i.e. $p_0 = 5 \times 10^6$.

- Questions:
1. What will the population size be in 10 years time?
 2. When will the population size reach 10^9 ?

We have a geometric sequence with $k = 0$, $a = 5 \times 10^6$, $r = 2$. Hence:

- Answers:
1. $a_{10} = 5 \times 10^6 \times 2^{10} = 5.12 \times 10^9$
 2. We need N so that $a_N \geq 10^9 > a_{N-1}$.
Note that $a_8 = \frac{1}{4}(5.12 \times 10^9) = 1.28 \times 10^9$

Simplest Malthusian model in population dynamics

Exponential Growth:

Let p_n be number of individuals in a population after n years.

Assume the population doubles every year, i.e. $\forall n \in \mathbb{N}^* \quad p_{n+1} = 2p_n$.

Assume the current population is five million. i.e. $p_0 = 5 \times 10^6$.

- Questions:
1. What will the population size be in 10 years time?
 2. When will the population size reach 10^9 ?

We have a geometric sequence with $k = 0$, $a = 5 \times 10^6$, $r = 2$. Hence:

Answers: 1. $a_{10} = 5 \times 10^6 \times 2^{10} = 5.12 \times 10^9$

2. We need N so that $a_N \geq 10^9 > a_{N-1}$.

Note that $a_8 = \frac{1}{4}(5.12 \times 10^9) = 1.28 \times 10^9$

and $a_7 = \frac{1}{8}(5.12 \times 10^9) = 0.64 \times 10^9$.

Simplest Malthusian model in population dynamics

Exponential Growth:

Let p_n be number of individuals in a population after n years.

Assume the population doubles every year, i.e. $\forall n \in \mathbb{N}^* \quad p_{n+1} = 2p_n$.

Assume the current population is five million. i.e. $p_0 = 5 \times 10^6$.

- Questions:
1. What will the population size be in 10 years time?
 2. When will the population size reach 10^9 ?

We have a geometric sequence with $k = 0$, $a = 5 \times 10^6$, $r = 2$. Hence:

Answers: 1. $a_{10} = 5 \times 10^6 \times 2^{10} = 5.12 \times 10^9$

2. We need N so that $a_N \geq 10^9 > a_{N-1}$.

Note that $a_8 = \frac{1}{4}(5.12 \times 10^9) = 1.28 \times 10^9$

and $a_7 = \frac{1}{8}(5.12 \times 10^9) = 0.64 \times 10^9$.

So $N = 8$ i.e. eight years.

Compound Interest

Let c_n be the capital (in \$) after n years.

Compound Interest

Let c_n be the capital (in \$) after n years.

Assume the capital yields 3% interest every year,

Compound Interest

Let c_n be the capital (in \$) after n years.

Assume the capital yields 3% interest every year, *i.e.*

$$\forall n \in \mathbb{N}^* \quad c_{n+1} = c_n + 0.03c_n = 1.03c_n.$$

Compound Interest

Let c_n be the capital (in \$) after n years.

Assume the capital yields 3% interest every year, *i.e.*

$$\forall n \in \mathbb{N}^* \quad c_{n+1} = c_n + 0.03c_n = 1.03c_n.$$

Assume the current capital is \$20K, *i.e.* $c_0 = 2 \times 10^4$.

Compound Interest

Let c_n be the capital (in \$) after n years.

Assume the capital yields 3% interest every year, *i.e.*

$$\forall n \in \mathbb{N}^* \quad c_{n+1} = c_n + 0.03c_n = 1.03c_n.$$

Assume the current capital is \$20K, *i.e.* $c_0 = 2 \times 10^4$.

Question: What will the capital be in 10 years?

Compound Interest

Let c_n be the capital (in \$) after n years.

Assume the capital yields 3% interest every year, *i.e.*

$$\forall n \in \mathbb{N}^* \quad c_{n+1} = c_n + 0.03c_n = 1.03c_n.$$

Assume the current capital is \$20K, *i.e.* $c_0 = 2 \times 10^4$.

Question: What will the capital be in 10 years?

$$\text{Answer: } \forall n \in \mathbb{N}^* \quad c_n = 2 \times 10^4 \times 1.03^n.$$

Compound Interest

Let c_n be the capital (in \$) after n years.

Assume the capital yields 3% interest every year, *i.e.*

$$\forall n \in \mathbb{N}^* \quad c_{n+1} = c_n + 0.03c_n = 1.03c_n.$$

Assume the current capital is \$20K, *i.e.* $c_0 = 2 \times 10^4$.

Question: What will the capital be in 10 years?

$$\text{Answer: } \forall n \in \mathbb{N}^* \quad c_n = 2 \times 10^4 \times 1.03^n.$$

$$c_{10} = 2 \times 10^4 \times 1.03^{10} = \$26\,87833.$$

Compound Interest

Let c_n be the capital (in \$) after n years.

Assume the capital yields 3% interest every year, *i.e.*

$$\forall n \in \mathbb{N}^* \quad c_{n+1} = c_n + 0.03c_n = 1.03c_n.$$

Assume the current capital is \$20K, *i.e.* $c_0 = 2 \times 10^4$.

Question: What will the capital be in 10 years?

Answer: $\forall n \in \mathbb{N}^* \quad c_n = 2 \times 10^4 \times 1.03^n$.

$$c_{10} = 2 \times 10^4 \times 1.03^{10} = \$26\,87833.$$

Monthly compounding:

Now suppose the capital yields $\frac{3}{12} = 0.25\%$ interest per *month*.

Compound Interest

Let c_n be the capital (in \$) after n years.

Assume the capital yields 3% interest every year, *i.e.*

$$\forall n \in \mathbb{N}^* \quad c_{n+1} = c_n + 0.03c_n = 1.03c_n.$$

Assume the current capital is \$20K, *i.e.* $c_0 = 2 \times 10^4$.

Question: What will the capital be in 10 years?

Answer: $\forall n \in \mathbb{N}^* \quad c_n = 2 \times 10^4 \times 1.03^n$.

$$c_{10} = 2 \times 10^4 \times 1.03^{10} = \$26\,87833.$$

Monthly compounding:

Now suppose the capital yields $\frac{3}{12} = 0.25\%$ interest per *month*.

Question: Is it the same as 3% per year?

Compound Interest

Let c_n be the capital (in \$) after n years.

Assume the capital yields 3% interest every year, *i.e.*

$$\forall n \in \mathbb{N}^* \quad c_{n+1} = c_n + 0.03c_n = 1.03c_n.$$

Assume the current capital is \$20K, *i.e.* $c_0 = 2 \times 10^4$.

Question: What will the capital be in 10 years?

$$\text{Answer: } \forall n \in \mathbb{N}^* \quad c_n = 2 \times 10^4 \times 1.03^n.$$

$$c_{10} = 2 \times 10^4 \times 1.03^{10} = \$26\,87833.$$

Monthly compounding:

Now suppose the capital yields $\frac{3}{12} = 0.25\%$ interest per *month*.

Question: Is it the same as 3% per year?

Answer: Now n is time in months and d_n capital at time n .

Compound Interest

Let c_n be the capital (in \$) after n years.

Assume the capital yields 3% interest every year, *i.e.*

$$\forall n \in \mathbb{N}^* \quad c_{n+1} = c_n + 0.03c_n = 1.03c_n.$$

Assume the current capital is \$20K, *i.e.* $c_0 = 2 \times 10^4$.

Question: What will the capital be in 10 years?

$$\text{Answer: } \forall n \in \mathbb{N}^* \quad c_n = 2 \times 10^4 \times 1.03^n.$$

$$c_{10} = 2 \times 10^4 \times 1.03^{10} = \$26\,87833.$$

Monthly compounding:

Now suppose the capital yields $\frac{3}{12} = 0.25\%$ interest per *month*.

Question: Is it the same as 3% per year?

Answer: Now n is time in months and d_n capital at time n .

$$d_0 = 2 \times 10^4, \quad \forall n \in \mathbb{N} \quad d_{n+1} = d_n + 0.0025d_n = 1.0025d_n.$$

Compound Interest

Let c_n be the capital (in \$) after n years.

Assume the capital yields 3% interest every year, *i.e.*

$$\forall n \in \mathbb{N}^* \quad c_{n+1} = c_n + 0.03c_n = 1.03c_n.$$

Assume the current capital is \$20K, *i.e.* $c_0 = 2 \times 10^4$.

Question: What will the capital be in 10 years?

$$\text{Answer: } \forall n \in \mathbb{N}^* \quad c_n = 2 \times 10^4 \times 1.03^n.$$

$$c_{10} = 2 \times 10^4 \times 1.03^{10} = \$26\,87833.$$

Monthly compounding:

Now suppose the capital yields $\frac{3}{12} = 0.25\%$ interest per *month*.

Question: Is it the same as 3% per year?

Answer: Now n is time in months and d_n capital at time n .

$$d_0 = 2 \times 10^4, \quad \forall n \in \mathbb{N} \quad d_{n+1} = d_n + 0.0025d_n = 1.0025d_n.$$

$$d_{120} = 2 \times 10^4 \times 1.0025^{120} = \$26\,98771.$$

Compound Interest

Let c_n be the capital (in \$) after n years.

Assume the capital yields 3% interest every year, *i.e.*

$$\forall n \in \mathbb{N}^* \quad c_{n+1} = c_n + 0.03c_n = 1.03c_n.$$

Assume the current capital is \$20K, *i.e.* $c_0 = 2 \times 10^4$.

Question: What will the capital be in 10 years?

$$\text{Answer: } \forall n \in \mathbb{N}^* \quad c_n = 2 \times 10^4 \times 1.03^n.$$

$$c_{10} = 2 \times 10^4 \times 1.03^{10} = \$26\,87833.$$

Monthly compounding:

Now suppose the capital yields $\frac{3}{12} = 0.25\%$ interest per *month*.

Question: Is it the same as 3% per year?

Answer: Now n is time in months and d_n capital at time n .

$$d_0 = 2 \times 10^4, \quad \forall n \in \mathbb{N} \quad d_{n+1} = d_n + 0.0025d_n = 1.0025d_n.$$

$$d_{120} = 2 \times 10^4 \times 1.0025^{120} = \$26\,98771. \quad \text{Slightly better!}$$

More applications of sequences

Compound Interest: If the bank is charging a fee of \$10 per year, the compound interest model becomes

$$\begin{cases} c_{n+1} = 1.03c_n - 10 & \forall n \in \mathbb{N}^*, \\ c_0 = 2 \times 10^4. \end{cases}$$

Compound Interest: If the bank is charging a fee of \$10 per year, the compound interest model becomes

$$\begin{cases} c_{n+1} = 1.03c_n - 10 & \forall n \in \mathbb{N}^*, \\ c_0 = 2 \times 10^4. \end{cases}$$

Population growth: If there is some immigration, bringing 10^3 new individuals to the population each year, the population dynamics model becomes

$$\begin{cases} p_{n+1} = 2p_n + 10^3 & \forall n \in \mathbb{N}^*, \\ p_0 = 5 \times 10^6. \end{cases}$$

Compound Interest: If the bank is charging a fee of \$10 per year, the compound interest model becomes

$$\begin{cases} c_{n+1} = 1.03c_n - 10 & \forall n \in \mathbb{N}^*, \\ c_0 = 2 \times 10^4. \end{cases}$$

Population growth: If there is some immigration, bringing 10^3 new individuals to the population each year, the population dynamics model becomes

$$\begin{cases} p_{n+1} = 2p_n + 10^3 & \forall n \in \mathbb{N}^*, \\ p_0 = 5 \times 10^6. \end{cases}$$

The sequences defined by these models are neither geometric nor arithmetic, but are a generalisation of both.

Compound Interest: If the bank is charging a fee of \$10 per year, the compound interest model becomes

$$\begin{cases} c_{n+1} = 1.03c_n - 10 & \forall n \in \mathbb{N}^*, \\ c_0 = 2 \times 10^4. \end{cases}$$

Population growth: If there is some immigration, bringing 10^3 new individuals to the population each year, the population dynamics model becomes

$$\begin{cases} p_{n+1} = 2p_n + 10^3 & \forall n \in \mathbb{N}^*, \\ p_0 = 5 \times 10^6. \end{cases}$$

The sequences defined by these models are neither geometric nor arithmetic, but are a generalisation of both.

We need to start again from scratch.

From implicit to explicit definitions; Example 3

We seek an explicit formula for the population given by the implicit formula at right, where d and p are shorthand for 10^3 and 5×10^6 .

$$\begin{cases} p_{n+1} = 2p_n + d \quad \forall n \in \mathbb{N}^*, \\ p_0 = p. \end{cases}$$

From implicit to explicit definitions; Example 3

We seek an explicit formula for the population given by the implicit formula at right, where d and p are shorthand for 10^3 and 5×10^6 .

$$\begin{cases} p_{n+1} = 2p_n + d \quad \forall n \in \mathbb{N}^*, \\ p_0 = p. \end{cases}$$

We start by generating the first few terms of the sequence:

$$p_0 = p,$$

From implicit to explicit definitions; Example 3

We seek an explicit formula for the population given by the implicit formula at right, where d and p are shorthand for 10^3 and 5×10^6 .

$$\begin{cases} p_{n+1} = 2p_n + d \quad \forall n \in \mathbb{N}^*, \\ p_0 = p. \end{cases}$$

We start by generating the first few terms of the sequence:

$$p_0 = p, \quad p_1 = 2p + d,$$

From implicit to explicit definitions; Example 3

We seek an explicit formula for the population given by the implicit formula at right, where d and p are shorthand for 10^3 and 5×10^6 .

$$\begin{cases} p_{n+1} = 2p_n + d \quad \forall n \in \mathbb{N}^*, \\ p_0 = p. \end{cases}$$

We start by generating the first few terms of the sequence:

$$p_0 = p, \quad p_1 = 2p + d, \quad p_2 = 2(2p + d) + d =$$

From implicit to explicit definitions; Example 3

We seek an explicit formula for the population given by the implicit formula at right, where d and p are shorthand for 10^3 and 5×10^6 .

$$\begin{cases} p_{n+1} = 2p_n + d \quad \forall n \in \mathbb{N}^*, \\ p_0 = p. \end{cases}$$

We start by generating the first few terms of the sequence:

$$p_0 = p, \quad p_1 = 2p + d, \quad p_2 = 2(2p + d) + d = 4p + 3d,$$

From implicit to explicit definitions; Example 3

We seek an explicit formula for the population given by the implicit formula at right, where d and p are shorthand for 10^3 and 5×10^6 .

$$\begin{cases} p_{n+1} = 2p_n + d \quad \forall n \in \mathbb{N}^*, \\ p_0 = p. \end{cases}$$

We start by generating the first few terms of the sequence:

$$p_0 = p, \quad p_1 = 2p + d, \quad p_2 = 2(2p + d) + d = 4p + 3d,$$

$$p_3 =$$

From implicit to explicit definitions; Example 3

We seek an explicit formula for the population given by the implicit formula at right, where d and p are shorthand for 10^3 and 5×10^6 .

$$\begin{cases} p_{n+1} = 2p_n + d \quad \forall n \in \mathbb{N}^*, \\ p_0 = p. \end{cases}$$

We start by generating the first few terms of the sequence:

$$\begin{aligned} p_0 &= p, & p_1 &= 2p + d, & p_2 &= 2(2p + d) + d = 4p + 3d, \\ p_3 &= 8p + 7d, \end{aligned}$$

From implicit to explicit definitions; Example 3

We seek an explicit formula for the population given by the implicit formula at right, where d and p are shorthand for 10^3 and 5×10^6 .

$$\begin{cases} p_{n+1} = 2p_n + d \quad \forall n \in \mathbb{N}^*, \\ p_0 = p. \end{cases}$$

We start by generating the first few terms of the sequence:

$$p_0 = p, \quad p_1 = 2p + d, \quad p_2 = 2(2p + d) + d = 4p + 3d,$$

$$p_3 = 8p + 7d,$$

$$\textbf{Claim: } \forall n \in \mathbb{N}^* \quad p_n = 2^n p + (2^n - 1)d.$$

From implicit to explicit definitions; Example 3

We seek an explicit formula for the population given by the implicit formula at right, where d and p are shorthand for 10^3 and 5×10^6 .

$$\begin{cases} p_{n+1} = 2p_n + d \quad \forall n \in \mathbb{N}^*, \\ p_0 = p. \end{cases}$$

We start by generating the first few terms of the sequence:

$$p_0 = p, \quad p_1 = 2p + d, \quad p_2 = 2(2p + d) + d = 4p + 3d,$$

$$p_3 = 8p + 7d,$$

$$\textbf{Claim: } \forall n \in \mathbb{N}^* \quad p_n = 2^n p + (2^n - 1)d.$$

Proof by mathematical induction that the claim is correct:

From implicit to explicit definitions; Example 3

We seek an explicit formula for the population given by the implicit formula at right, where d and p are shorthand for 10^3 and 5×10^6 .

$$\begin{cases} p_{n+1} = 2p_n + d \quad \forall n \in \mathbb{N}^*, \\ p_0 = p. \end{cases}$$

We start by generating the first few terms of the sequence:

$$p_0 = p, \quad p_1 = 2p + d, \quad p_2 = 2(2p + d) + d = 4p + 3d,$$

$$p_3 = 8p + 7d,$$

$$\textbf{Claim: } \forall n \in \mathbb{N}^* \quad p_n = 2^n p + (2^n - 1)d.$$

Proof by mathematical induction that the claim is correct:

Basis step: For $n=0$, formula gives $p_0 = 2^0 p + (2^0 - 1)d = p$, agreeing with the implicit definition.

From implicit to explicit definitions; Example 3

We seek an explicit formula for the population given by the implicit formula at right, where d and p are shorthand for 10^3 and 5×10^6 .

$$\begin{cases} p_{n+1} = 2p_n + d \quad \forall n \in \mathbb{N}^*, \\ p_0 = p. \end{cases}$$

We start by generating the first few terms of the sequence:

$$p_0 = p, \quad p_1 = 2p + d, \quad p_2 = 2(2p + d) + d = 4p + 3d,$$

$$p_3 = 8p + 7d,$$

$$\textbf{Claim: } \forall n \in \mathbb{N}^* \quad p_n = 2^n p + (2^n - 1)d.$$

Proof by mathematical induction that the claim is correct:

Basis step: For $n=0$, formula gives $p_0 = 2^0 p + (2^0 - 1)d = p$, agreeing with the implicit definition.

Inductive step: Assume the formula is correct for up to and including some fixed n .

From implicit to explicit definitions; Example 3

We seek an explicit formula for the population given by the implicit formula at right, where d and p are shorthand for 10^3 and 5×10^6 .

$$\begin{cases} p_{n+1} = 2p_n + d \quad \forall n \in \mathbb{N}^*, \\ p_0 = p. \end{cases}$$

We start by generating the first few terms of the sequence:

$$p_0 = p, \quad p_1 = 2p + d, \quad p_2 = 2(2p + d) + d = 4p + 3d,$$

$$p_3 = 8p + 7d,$$

$$\textbf{Claim: } \forall n \in \mathbb{N}^* \quad p_n = 2^n p + (2^n - 1)d.$$

Proof by mathematical induction that the claim is correct:

Basis step: For $n=0$, formula gives $p_0 = 2^0 p + (2^0 - 1)d = p$, agreeing with the implicit definition.

Inductive step: Assume the formula is correct for up to and including some fixed n . Then

$$p_{n+1} = 2p_n + d \quad (\text{from the implicit definition})$$

From implicit to explicit definitions; Example 3

We seek an explicit formula for the population given by the implicit formula at right, where d and p are shorthand for 10^3 and 5×10^6 .

$$\begin{cases} p_{n+1} = 2p_n + d \quad \forall n \in \mathbb{N}^*, \\ p_0 = p. \end{cases}$$

We start by generating the first few terms of the sequence:

$$p_0 = p, \quad p_1 = 2p + d, \quad p_2 = 2(2p + d) + d = 4p + 3d,$$

$$p_3 = 8p + 7d,$$

$$\textbf{Claim: } \forall n \in \mathbb{N}^* \quad p_n = 2^n p + (2^n - 1)d.$$

Proof by mathematical induction that the claim is correct:

Basis step: For $n=0$, formula gives $p_0 = 2^0 p + (2^0 - 1)d = p$, agreeing with the implicit definition.

Inductive step: Assume the formula is correct for up to and including some fixed n . Then

$$\begin{aligned} p_{n+1} &= 2p_n + d \quad (\text{from the implicit definition}) \\ &= 2(2^n p + (2^n - 1)d) + d \quad (\text{by the inductive assumption}) \end{aligned}$$

From implicit to explicit definitions; Example 3

We seek an explicit formula for the population given by the implicit formula at right, where d and p are shorthand for 10^3 and 5×10^6 .

$$\begin{cases} p_{n+1} = 2p_n + d \quad \forall n \in \mathbb{N}^*, \\ p_0 = p. \end{cases}$$

We start by generating the first few terms of the sequence:

$$p_0 = p, \quad p_1 = 2p + d, \quad p_2 = 2(2p + d) + d = 4p + 3d,$$

$$p_3 = 8p + 7d,$$

$$\textbf{Claim: } \forall n \in \mathbb{N}^* \quad p_n = 2^n p + (2^n - 1)d.$$

Proof by mathematical induction that the claim is correct:

Basis step: For $n=0$, formula gives $p_0 = 2^0 p + (2^0 - 1)d = p$, agreeing with the implicit definition.

Inductive step: Assume the formula is correct for up to and including some fixed n . Then

$$\begin{aligned} p_{n+1} &= 2p_n + d \quad (\text{from the implicit definition}) \\ &= 2(2^n p + (2^n - 1)d) + d \quad (\text{by the inductive assumption}) \\ &= 2^{n+1} p + (2^{n+1} - 1)d \quad (\text{by algebraic simplification}) \end{aligned}$$

From implicit to explicit definitions; Example 3

We seek an explicit formula for the population given by the implicit formula at right, where d and p are shorthand for 10^3 and 5×10^6 .

$$\begin{cases} p_{n+1} = 2p_n + d \quad \forall n \in \mathbb{N}^*, \\ p_0 = p. \end{cases}$$

We start by generating the first few terms of the sequence:

$$p_0 = p, \quad p_1 = 2p + d, \quad p_2 = 2(2p + d) + d = 4p + 3d,$$

$$p_3 = 8p + 7d, \quad \textbf{Claim: } \forall n \in \mathbb{N}^* \quad p_n = 2^n p + (2^n - 1)d.$$

Proof by mathematical induction that the claim is correct:

Basis step: For $n=0$, formula gives $p_0 = 2^0 p + (2^0 - 1)d = p$, agreeing with the implicit definition.

Inductive step: Assume the formula is correct for up to and including some fixed n . Then

$$\begin{aligned} p_{n+1} &= 2p_n + d \quad (\text{from the implicit definition}) \\ &= 2(2^n p + (2^n - 1)d) + d \quad (\text{by the inductive assumption}) \\ &= 2^{n+1} p + (2^{n+1} - 1)d \quad (\text{by algebraic simplification}) \end{aligned}$$

and so the formula is also correct for $n+1$.