

Discrete Mathematical Models

Lecture 18

Kane Townsend

Semester 2, 2024

The Product Rule (with some clarity)

Suppose you run n experiments with sample spaces S_1, S_2, \dots, S_n . For each $1 \leq i \leq n$ let $E_i \subseteq S_i$ be an event. If the events E_1, \dots, E_n are 'independent' of each other; then the probability of composite event ' E_1 and E_2 and ... and E_n ' is

$$\mathbb{P}_{S_1 \times S_2 \times \dots \times S_n}(E_1 \times E_2 \times \dots \times E_n) = \mathbb{P}_{S_1}(E_1) \times \mathbb{P}_{S_2}(E_2) \times \dots \times \mathbb{P}_{S_n}(E_n).$$

The Product Rule (with some clarity)

Suppose you run n experiments with sample spaces S_1, S_2, \dots, S_n . For each $1 \leq i \leq n$ let $E_i \subseteq S_i$ be an event. If the events E_1, \dots, E_n are 'independent' of each other; then the probability of composite event ' E_1 and E_2 and ... and E_n ' is

$$\mathbb{P}_{S_1 \times S_2 \times \dots \times S_n}(E_1 \times E_2 \times \dots \times E_n) = \mathbb{P}_{S_1}(E_1) \times \mathbb{P}_{S_2}(E_2) \times \dots \times \mathbb{P}_{S_n}(E_n).$$

- $\mathbb{P}(E_1) = \frac{|E_1|}{|S_1|}$

The Product Rule (with some clarity)

Suppose you run n experiments with sample spaces S_1, S_2, \dots, S_n . For each $1 \leq i \leq n$ let $E_i \subseteq S_i$ be an event. If the events E_1, \dots, E_n are 'independent' of each other; then the probability of composite event ' E_1 and E_2 and ... and E_n ' is

$$\mathbb{P}_{S_1 \times S_2 \times \dots \times S_n}(E_1 \times E_2 \times \dots \times E_n) = \mathbb{P}_{S_1}(E_1) \times \mathbb{P}_{S_2}(E_2) \times \dots \times \mathbb{P}_{S_n}(E_n).$$

- $\mathbb{P}(E_1) = \frac{|E_1|}{|S_1|}$
- $\mathbb{P}(E_2) = \frac{|E_2|}{|S_2|}$
- ...
- $\mathbb{P}(E_n) = \frac{|E_n|}{|S_n|}$

The Product Rule (with some clarity)

Suppose you run n experiments with sample spaces S_1, S_2, \dots, S_n . For each $1 \leq i \leq n$ let $E_i \subseteq S_i$ be an event. If the events E_1, \dots, E_n are 'independent' of each other; then the probability of composite event ' E_1 and E_2 and ... and E_n ' is


$$\mathbb{P}_{S_1 \times S_2 \times \dots \times S_n}(E_1 \times E_2 \times \dots \times E_n) = \mathbb{P}_{S_1}(E_1) \times \mathbb{P}_{S_2}(E_2) \times \dots \times \mathbb{P}_{S_n}(E_n).$$

- $\mathbb{P}(E_1) = \frac{|E_1|}{|S_1|}$
- $\mathbb{P}(E_2) = \frac{|E_2|}{|S_2|}$
- ...
- $\mathbb{P}(E_n) = \frac{|E_n|}{|S_n|}$
- $\mathbb{P}(E_n) = \frac{|E_1 \times E_2 \times \dots \times E_n|}{|S_1 \times S_2 \times \dots \times S_n|}$

Density Functions


Frequency Histograms

- One way to visualize all possible outcomes of an experiment together is to draw a **frequency histogram**.
- We have already seen some simple examples, like tossing a die with equally likely possible outcomes: 1, 2, 3, 4, 5, 6:



Probability Density Functions


- The **Probability Density Function** (or just **Density**) is obtained from a Frequency Histogram by **normalizing**. We divide the vertical axis by the total number of outcomes.
- Continuing the die-tossing example, we have



What is the area under the curve? Why?

Cumulative Probability Distribution Functions


- The Cumulative Probability Distribution Function (or Distribution) is obtained from the Density Function by graphing cumulative totals.
- Continuing the die-tossing example, we have




- We will only use of cumulative distributions when looking up probability values in tables or online.

Uniform Distribution

- When every event has the same probability the resulting densities and distributions are called 'uniform'. Examples:
- Uniform density:



- Uniform distribution:



Tossing two coins:

- Some more interesting densities and distributions are obtained by considering events which combine several outcomes.
- For example, tossing two coins. A neat way to list all possible outcomes is to expand

$$\begin{aligned} & (T + H)(T + H) \\ = & TT + TH + HT + HH \end{aligned}$$

- *What is the sample space?*

$$\{TT, TH, HT, HH\}$$

Tossing two coins:


- Now consider events:
 - E_0 : 'No heads'
 - E_1 : 'exactly 1 Head'
 - E_2 : 'exactly 2 Heads'
- *Which subsets of the sample space correspond to these events?*

Tossing two coins:

- Now consider events:
 - E_0 : 'No heads'
 - E_1 : 'exactly 1 Head'
 - E_2 : 'exactly 2 Heads'
- *Which subsets of the sample space correspond to these events?*

$$E_0 = \{TT\}, \quad E_1 = \{TH, HT\}, \quad E_2 = \{HH\}$$

Frequency Histogram:




Two fair coins: Density and Distribution Functions

- Assuming a fair coin (equally likely outcomes), divide out by the size of the sample space to get density function.

Two fair coins: Density and Distribution Functions

- Assuming a fair coin (equally likely outcomes), divide out by the size of the sample space to get density function. Then take the cumulative sum of the density to get the distribution:



Class Example - Three Fair Coins

Q: *For tossing three fair coins, what is the sample space?*

Class Example - Three Fair Coins

Q: *For tossing three fair coins, what is the sample space?*

A: All possible outcomes are given by $(T + H)(T + H)(T + H)$
 $= TTT + TTH + THT + THH + HTT + HTH + HHT + HHH.$
S.Space = $\{TTT, TTH, THT, THH, HTT, HTH, HHT, HHH\}.$

Class Example - Three Fair Coins

Q: *For tossing three fair coins, what is the sample space?*

A: All possible outcomes are given by $(T + H)(T + H)(T + H)$
 $= TTT + TTH + THT + THH + HTT + HTH + HHT + HHH$.
S.Space = $\{TTT, TTH, THT, THH, HTT, HTH, HHT, HHH\}$.

Q: *What subsets of the sample space correspond to the events E_0 : '0 heads', ' E_1 : 'exactly 1 Head', E_2 : 'exactly 2 Heads' and E_3 : 'exactly 3 Heads' ?*

Class Example - Three Fair Coins

Q: *For tossing three fair coins, what is the sample space?*

A: All possible outcomes are given by $(T + H)(T + H)(T + H)$
 $= TTT + TTH + THT + THH + HTT + HTH + HHT + HHH$.
S.Space = $\{TTT, TTH, THT, THH, HTT, HTH, HHT, HHH\}$.

Q: *What subsets of the sample space correspond to the events E_0 : '0 heads', ' E_1 : 'exactly 1 Head', E_2 : 'exactly 2 Heads' and E_3 : 'exactly 3 Heads' ?*

A: $E_0 = \{TTT\}$; $E_1 = \{TTH, THT, HTT\}$; $E_2 = \{THH, HTH, HHT\}$;
 $E_3 = \{HHH\}$.

Class Example - Three Fair Coins

Q: *For tossing three fair coins, what is the sample space?*

A: All possible outcomes are given by $(T + H)(T + H)(T + H)$
 $= TTT + TTH + THT + THH + HTT + HTH + HHT + HHH$.
S.Space = $\{TTT, TTH, THT, THH, HTT, HTH, HHT, HHH\}$.

Q: *What subsets of the sample space correspond to the events E_0 : '0 heads', ' E_1 : 'exactly 1 Head', ' E_2 : 'exactly 2 Heads' and ' E_3 : 'exactly 3 Heads' ?*

A: $E_0 = \{TTT\}$; $E_1 = \{TTH, THT, HTT\}$; $E_2 = \{THH, HTH, HHT\}$;
 $E_3 = \{HHH\}$.

Q: *What is the size of the sample space?*

Class Example - Three Fair Coins

Q: *For tossing three fair coins, what is the sample space?*

A: All possible outcomes are given by $(T + H)(T + H)(T + H)$
 $= TTT + TTH + THT + THH + HTT + HTH + HHT + HHH$.
S.Space = $\{TTT, TTH, THT, THH, HTT, HTH, HHT, HHH\}$.

Q: *What subsets of the sample space correspond to the events E_0 : '0 heads', ' E_1 : 'exactly 1 Head', ' E_2 : 'exactly 2 Heads' and ' E_3 : 'exactly 3 Heads' ?*

A: $E_0 = \{TTT\}$; $E_1 = \{TTH, THT, HTT\}$; $E_2 = \{THH, HTH, HHT\}$;
 $E_3 = \{HHH\}$.

Q: *What is the size of the sample space?* A: 8

Class Example - Three Fair Coins

Q: *For tossing three fair coins, what is the sample space?*

A: All possible outcomes are given by $(T + H)(T + H)(T + H)$
 $= TTT + TTH + THT + THH + HTT + HTH + HHT + HHH$.
S.Space = $\{TTT, TTH, THT, THH, HTT, HTH, HHT, HHH\}$.

Q: *What subsets of the sample space correspond to the events E_0 : '0 heads', ' E_1 : 'exactly 1 Head', ' E_2 : 'exactly 2 Heads' and ' E_3 : 'exactly 3 Heads' ?*

A: $E_0 = \{TTT\}$; $E_1 = \{TTH, THT, HTT\}$; $E_2 = \{THH, HTH, HHT\}$;
 $E_3 = \{HHH\}$.

Q: *What is the size of the sample space?* A: 8

Q: *What do we divide frequencies by (on the Frequency Histogram) to get the Density Function?*

Class Example - Three Fair Coins

Q: *For tossing three fair coins, what is the sample space?*

A: All possible outcomes are given by $(T + H)(T + H)(T + H)$
 $= TTT + TTH + THT + THH + HTT + HTH + HHT + HHH$.
S.Space = $\{TTT, TTH, THT, THH, HTT, HTH, HHT, HHH\}$.

Q: *What subsets of the sample space correspond to the events E_0 : '0 heads', E_1 : 'exactly 1 Head', E_2 : 'exactly 2 Heads' and E_3 : 'exactly 3 Heads' ?*

A: $E_0 = \{TTT\}$; $E_1 = \{TTH, THT, HTT\}$; $E_2 = \{THH, HTH, HHT\}$;
 $E_3 = \{HHH\}$.

Q: *What is the size of the sample space?* A: 8

Q: *What do we divide frequencies by (on the Frequency Histogram) to get the Density Function?* A: 8

Class Example - Three Fair Coins

Q: *For tossing three fair coins, what is the sample space?*

A: All possible outcomes are given by $(T + H)(T + H)(T + H)$
 $= TTT + TTH + THT + THH + HTT + HTH + HHT + HHH$.
S.Space = $\{TTT, TTH, THT, THH, HTT, HTH, HHT, HHH\}$.

Q: *What subsets of the sample space correspond to the events E_0 : '0 heads', E_1 : 'exactly 1 Head', E_2 : 'exactly 2 Heads' and E_3 : 'exactly 3 Heads' ?*

A: $E_0 = \{TTT\}$; $E_1 = \{TTH, THT, HTT\}$; $E_2 = \{THH, HTH, HHT\}$;
 $E_3 = \{HHH\}$.

Q: *What is the size of the sample space?* A: 8

Q: *What do we divide frequencies by (on the Frequency Histogram) to get the Density Function?* A: 8

Q: *What are the probabilities of the four events E_0, E_1, E_2, E_3 ?*

Class Example - Three Fair Coins

Q: *For tossing three fair coins, what is the sample space?*

A: All possible outcomes are given by $(T + H)(T + H)(T + H)$
 $= TTT + TTH + THT + THH + HTT + HTH + HHT + HHH$.
S.Space = $\{TTT, TTH, THT, THH, HTT, HTH, HHT, HHH\}$.

Q: *What subsets of the sample space correspond to the events E_0 : '0 heads', ' E_1 : 'exactly 1 Head', E_2 : 'exactly 2 Heads' and E_3 : 'exactly 3 Heads' ?*

A: $E_0 = \{TTT\}$; $E_1 = \{TTH, THT, HTT\}$; $E_2 = \{THH, HTH, HHT\}$;
 $E_3 = \{HHH\}$.


Q: *What is the size of the sample space?* A: 8

Q: *What do we divide frequencies by (on the Frequency Histogram) to get the Density Function?* A: 8

Q: *What are the probabilities of the four events E_0, E_1, E_2, E_3 ?*


A: $\mathbb{P}(E_0) = \frac{1}{8}$; $\mathbb{P}(E_1) = \frac{3}{8}$; $\mathbb{P}(E_2) = \frac{3}{8}$; $\mathbb{P}(E_3) = \frac{1}{8}$.

Three Fair Coins: Density and Distribution Functions




Binomial Probability Distributions

The family of functions that come from coin-tossing are all examples of **binomial** densities/distributions:



Bell-like curves for large n


As n gets larger and larger these **binomial probability density functions** get closer and closer to the famous Bell Curve:



which is the so-called **'Normal' Probability Density Function**. This convergence to the normal probability function is a result of the **Central Limit Theorem**.


Binomial Theorem

Pascal's Triangle



Pascal's Triangle

- Frequencies in Coin-Tossing are numbers in Pascal's Triangle




- Each row is generated by expanding a binomial, eg:

$$(y + x)^4 = y^4 + 4y^3x + 6y^2x^2 + 4yx^3 + x^4.$$

Pascal's Triangle

- We've seen these numbers before in 'combinations': $\binom{n}{k}$:



The Binomial Theorem

- The **Binomial Theorem** states that

$$(y + x)^n = \binom{n}{0} y^n x^0 + \binom{n}{1} y^{n-1} x^1 + \cdots + \binom{n}{n} y^0 x^n$$

and gives the rows of Pascal's Triangle in its coefficients.

Idea of Proof of Binomial Theorem:

$$(y + x)(y + x)(y + x)$$

$$= yyy + yyx + yxy + yxx + xyy + xyx + xxy + xxx$$

$$= \underbrace{yyy}_{\binom{3}{0} x's} + \underbrace{yyx + yxy + xyy}_{\binom{3}{1} x's} + \underbrace{yxx + xyx + xxy}_{\binom{3}{2} x's} + \underbrace{xxx}_{\binom{3}{3} x's}$$



Tossing n Coins

- Let's toss a (fair) coin n times ($n \in \mathbb{N}$.)

Tossing n Coins

- Let's toss a (fair) coin n times ($n \in \mathbb{N}$.)
- As earlier, let E_k denote the event of obtaining k heads.

Tossing n Coins

- Let's toss a (fair) coin n times ($n \in \mathbb{N}$.)
- As earlier, let E_k denote the event of obtaining k heads.
- Then
 - E_0 can occur in $\binom{n}{0} = 1$ way, i.e. $|E_0| = \binom{n}{0}$

Tossing n Coins

- Let's toss a (fair) coin n times ($n \in \mathbb{N}$.)
- As earlier, let E_k denote the event of obtaining k heads.
- Then
 - E_0 can occur in $\binom{n}{0} = 1$ way, i.e. $|E_0| = \binom{n}{0}$
 - E_1 can occur in $\binom{n}{1} = n$ ways, i.e. $|E_1| = \binom{n}{1}$

Tossing n Coins

- Let's toss a (fair) coin n times ($n \in \mathbb{N}$.)
- As earlier, let E_k denote the event of obtaining k heads.
- Then
 - E_0 can occur in $\binom{n}{0} = 1$ way, i.e. $|E_0| = \binom{n}{0}$
 - E_1 can occur in $\binom{n}{1} = n$ ways, i.e. $|E_1| = \binom{n}{1}$
 - E_2 can occur in $\binom{n}{2}$ ways, i.e. $|E_2| = \binom{n}{2}$
 - \vdots
 - E_n can occur in $\binom{n}{n}$ ways, i.e. $|E_n| = \binom{n}{n}$

Tossing n Coins

- Let's toss a (fair) coin n times ($n \in \mathbb{N}$.)
- As earlier, let E_k denote the event of obtaining k heads.
- Then
 - E_0 can occur in $\binom{n}{0} = 1$ way, i.e. $|E_0| = \binom{n}{0}$
 - E_1 can occur in $\binom{n}{1} = n$ ways, i.e. $|E_1| = \binom{n}{1}$
 - E_2 can occur in $\binom{n}{2}$ ways, i.e. $|E_2| = \binom{n}{2}$
 - \vdots
 - E_n can occur in $\binom{n}{n}$ ways, i.e. $|E_n| = \binom{n}{n}$

What is the total size of the sample space?

- I.e. what is

$$\binom{n}{0} + \binom{n}{1} + \binom{n}{2} + \dots + \binom{n}{n}?$$

- The binomial theorem gives a neat way to find the sum.
- Set $x = y = 1$ in $(x + y)^n$. Then

$$\binom{n}{0}1^n1^0 + \binom{n}{1}1^{n-1}1^1 + \cdots + \binom{n}{n}1^01^n = (1 + 1)^n$$

so that

$$\boxed{\binom{n}{0} + \binom{n}{1} + \cdots + \binom{n}{n} = 2^n.}$$

- The binomial theorem gives a neat way to find the sum.
- Set $x = y = 1$ in $(x + y)^n$. Then

$$\binom{n}{0}1^n1^0 + \binom{n}{1}1^{n-1}1^1 + \cdots + \binom{n}{n}1^01^n = (1 + 1)^n$$

so that

$$\boxed{\binom{n}{0} + \binom{n}{1} + \cdots + \binom{n}{n} = 2^n.}$$

- We can also get 2^n by observing that there are two possible outcomes for each toss, and so $2 \times 2 \times \cdots \times 2 = 2^n$ possible outcomes for n tosses.

- The binomial theorem gives a neat way to find the sum.
- Set $x = y = 1$ in $(x + y)^n$. Then

$$\binom{n}{0}1^n1^0 + \binom{n}{1}1^{n-1}1^1 + \cdots + \binom{n}{n}1^01^n = (1 + 1)^n$$

so that

$$\boxed{\binom{n}{0} + \binom{n}{1} + \cdots + \binom{n}{n} = 2^n.}$$

- We can also get 2^n by observing that there are two possible outcomes for each toss, and so $2 \times 2 \times \cdots \times 2 = 2^n$ possible outcomes for n tosses.
- So, for example, the probability of obtaining exactly three heads from six tosses of a fair coin is

$$\frac{\binom{6}{3}}{2^6} = \frac{6 \cdot 5 \cdot 4}{3 \cdot 2 \cdot 1} = \frac{20}{64} = \frac{5}{16}.$$

- The binomial theorem gives a neat way to find the sum.
- Set $x = y = 1$ in $(x + y)^n$. Then

$$\binom{n}{0}1^n1^0 + \binom{n}{1}1^{n-1}1^1 + \cdots + \binom{n}{n}1^01^n = (1 + 1)^n$$

so that

$$\boxed{\binom{n}{0} + \binom{n}{1} + \cdots + \binom{n}{n} = 2^n.}$$

- We can also get 2^n by observing that there are two possible outcomes for each toss, and so $2 \times 2 \times \cdots \times 2 = 2^n$ possible outcomes for n tosses.
- So, for example, the probability of obtaining exactly three heads from six tosses of a fair coin is


$$\frac{\binom{6}{3}}{2^6} = \frac{6 \cdot 5 \cdot 4}{3 \cdot 2 \cdot 1} = \frac{20}{64} = \frac{5}{16}.$$

- Examples will be found in worksheet questions.

Binomial Distribution


A tree representation of Coin-tossing

- Another way to list all the outcomes of an event is to draw a
Tree Diagram of the Possibilities



Three tosses of a fair coin

- This allows us to deal with fair coins, as before:



Outcome:

$$\Pr(TTT) = 1/8$$

$$\Pr(TTH) = 1/8$$

$$\Pr(THT) = 1/8$$

$$\Pr(THH) = 1/8$$

$$\Pr(HTT) = 1/8$$

$$\Pr(HTH) = 1/8$$

$$\Pr(HHT) = 1/8$$

$$\Pr(HHH) = 1/8$$

Three tosses of a fair coin

Collecting possibilities from the tree and using the sum rule gives

$$\mathbb{P}(0\text{heads}) = \frac{1}{8}, \quad \mathbb{P}(1\text{head}) = \frac{3}{8}, \quad \mathbb{P}(2\text{heads}) = \frac{3}{8}, \quad \mathbb{P}(3\text{heads}) = \frac{1}{8}$$

Three tosses of a fair coin

Collecting possibilities from the tree and using the sum rule gives

$$\mathbb{P}(0\text{heads}) = \frac{1}{8}, \quad \mathbb{P}(1\text{head}) = \frac{3}{8}, \quad \mathbb{P}(2\text{heads}) = \frac{3}{8}, \quad \mathbb{P}(3\text{heads}) = \frac{1}{8}$$


– the same density function as before; $n = 3$ and $p = \frac{1}{2}$ (fair coin):

Three tosses of a fair coin

Collecting possibilities from the tree and using the sum rule gives

$$\mathbb{P}(0\text{heads}) = \frac{1}{8}, \quad \mathbb{P}(1\text{head}) = \frac{3}{8}, \quad \mathbb{P}(2\text{heads}) = \frac{3}{8}, \quad \mathbb{P}(3\text{heads}) = \frac{1}{8}$$

– the same density function as before; $n = 3$ and $p = \frac{1}{2}$ (fair coin):




Three tosses of an unfair coin

- But we can also deal with an **unfair coin** – not equal likelihood:

Three tosses of an unfair coin

- But we can also deal with an **unfair coin** – not equal likelihood:



Three tosses of an unfair coin

Collecting possibilities from the tree and using the sum rule gives

$$\mathbb{P}(0\text{heads}) = \frac{125}{512}, \mathbb{P}(1\text{head}) = \frac{225}{512}, \mathbb{P}(2\text{heads}) = \frac{135}{512}, \mathbb{P}(3\text{heads}) = \frac{27}{512}$$

Three tosses of an unfair coin

Collecting possibilities from the tree and using the sum rule gives

$$\mathbb{P}(0\text{heads}) = \frac{125}{512}, \mathbb{P}(1\text{head}) = \frac{225}{512}, \mathbb{P}(2\text{heads}) = \frac{135}{512}, \mathbb{P}(3\text{heads}) = \frac{27}{512}$$


The unfair coin with $n = 3$ tosses and probability $p = 3/8$ of heads on a single toss, gives a non-symmetric binomial density function:

Three tosses of an unfair coin

Collecting possibilities from the tree and using the sum rule gives

$$\mathbb{P}(0\text{heads}) = \frac{125}{512}, \mathbb{P}(1\text{head}) = \frac{225}{512}, \mathbb{P}(2\text{heads}) = \frac{135}{512}, \mathbb{P}(3\text{heads}) = \frac{27}{512}$$

The unfair coin with $n = 3$ tosses and probability $p = 3/8$ of heads on a single toss, gives a non-symmetric binomial density function:




Three tosses of an unfair coin

Collecting possibilities from the tree and using the sum rule gives

$$\mathbb{P}(0\text{heads}) = \frac{125}{512}, \mathbb{P}(1\text{head}) = \frac{225}{512}, \mathbb{P}(2\text{heads}) = \frac{135}{512}, \mathbb{P}(3\text{heads}) = \frac{27}{512}$$

The unfair coin with $n = 3$ tosses and probability $p = 3/8$ of heads on a single toss, gives a non-symmetric binomial density function:



The general binomial density function for n trials (e.g. tosses) with probability p of a success (e.g. head) on each trial is given by

$$\mathbb{P}(k \text{ successes}) = \binom{n}{k} p^k (1-p)^{n-k}.$$

“Fun” Problems

Another general probability example

In a group of 10 students, 5 are studying computer science, 2 are studying art history, and 3 are studying mathematics. We pick a student from this group and ask what her/his major is.

Another general probability example

In a group of 10 students, 5 are studying computer science, 2 are studying art history, and 3 are studying mathematics. We pick a student from this group and ask what her/his major is.

The sample space is $S = \{M, A, C\}$, where M stand for maths, A for art history, and C for computer science.

Another general probability example

In a group of 10 students, 5 are studying computer science, 2 are studying art history, and 3 are studying mathematics. We pick a student from this group and ask what her/his major is.

The sample space is $S = \{M, A, C\}$, where M stand for maths, A for art history, and C for computer science.

The probability density function $\mathbb{P} : S \rightarrow \mathbb{Q}_+$ is given by

$$\mathbb{P}(M) = \frac{3}{10}, \quad \mathbb{P}(A) = \frac{2}{10}, \quad \mathbb{P}(C) = \frac{5}{10}.$$

Another general probability example

In a group of 10 students, 5 are studying computer science, 2 are studying art history, and 3 are studying mathematics. We pick a student from this group and ask what her/his major is.

The sample space is $S = \{M, A, C\}$, where M stand for maths, A for art history, and C for computer science.

The probability density function $\mathbb{P} : S \rightarrow \mathbb{Q}_+$ is given by

$$\mathbb{P}(M) = \frac{3}{10}, \quad \mathbb{P}(A) = \frac{2}{10}, \quad \mathbb{P}(C) = \frac{5}{10}.$$

The associated event probabilities are

$$\begin{aligned} \mathbb{P}(\emptyset) &= 0, & \mathbb{P}(\{M\}) &= \frac{3}{10}, & \mathbb{P}(\{A\}) &= \frac{2}{10}, & \mathbb{P}(\{C\}) &= \frac{5}{10}, \\ \mathbb{P}(\{M, A\}) &= \frac{5}{10}, & \mathbb{P}(\{A, C\}) &= \frac{7}{10}, & \mathbb{P}(\{M, C\}) &= \frac{8}{10}, & \mathbb{P}(S) &= 1. \end{aligned}$$

The Monty Hall problem

A game: Three doors, with a prize behind one of them.

The contestant chooses one door.

The host, who knows where the prize is, opens one of the other two doors, revealing that the prize is not there.

The Monty Hall problem

A game: Three doors, with a prize behind one of them.

The contestant chooses one door.

The host, who knows where the prize is, opens one of the other two doors, revealing that the prize is not there.

The contestant can then change her/his choice, or not.

The Monty Hall problem

A game: Three doors, with a prize behind one of them.

The contestant chooses one door.

The host, who knows where the prize is, opens one of the other two doors, revealing that the prize is not there.

The contestant can then change her/his choice, or not.

Should she/he change doors?

The Monty Hall problem

A game: Three doors, with a prize behind one of them.

The contestant chooses one door.

The host, who knows where the prize is, opens one of the other two doors, revealing that the prize is not there.

The contestant can then change her/his choice, or not.

Should she/he change doors?

Before the host opens the door:

Sample space is $S = \{d_1, d_2, d_3\}$ with $\mathbb{P}(d_1) = \mathbb{P}(d_2) = \mathbb{P}(d_3) = \frac{1}{3}$.

The Monty Hall problem

A game: Three doors, with a prize behind one of them.

The contestant chooses one door.

The host, who knows where the prize is, opens one of the other two doors, revealing that the prize is not there.

The contestant can then change her/his choice, or not.

Should she/he change doors?

Before the host opens the door:

Sample space is $S = \{d_1, d_2, d_3\}$ with $\mathbb{P}(d_1) = \mathbb{P}(d_2) = \mathbb{P}(d_3) = \frac{1}{3}$.

The contestant chooses d_1 and has a $\frac{1}{3}$ chance of winning.

The Monty Hall problem

A game: Three doors, with a prize behind one of them.

The contestant chooses one door.

The host, who knows where the prize is, opens one of the other two doors, revealing that the prize is not there.

The contestant can then change her/his choice, or not.

Should she/he change doors?

Before the host opens the door:

Sample space is $S = \{d_1, d_2, d_3\}$ with $\mathbb{P}(d_1) = \mathbb{P}(d_2) = \mathbb{P}(d_3) = \frac{1}{3}$.

The contestant chooses d_1 and has a $\frac{1}{3}$ chance of winning.

The host then opens door 2:

The sample space is now $S' = \{d_1, d_3\}$, but $\mathbb{P}(d_1)$ remains at $\frac{1}{3}$ (since the prize hasn't moved)

The Monty Hall problem

A game: Three doors, with a prize behind one of them.

The contestant chooses one door.

The host, who knows where the prize is, opens one of the other two doors, revealing that the prize is not there.

The contestant can then change her/his choice, or not.

Should she/he change doors?

Before the host opens the door:

Sample space is $S = \{d_1, d_2, d_3\}$ with $\mathbb{P}(d_1) = \mathbb{P}(d_2) = \mathbb{P}(d_3) = \frac{1}{3}$.

The contestant chooses d_1 and has a $\frac{1}{3}$ chance of winning.

The host then opens door 2:

The sample space is now $S' = \{d_1, d_3\}$, but $\mathbb{P}(d_1)$ remains at $\frac{1}{3}$ (since the prize hasn't moved) so $\mathbb{P}(d_3) = \frac{2}{3}$ since the sum of the probabilities must be 1.

The Monty Hall problem

A game: Three doors, with a prize behind one of them.

The contestant chooses one door.

The host, who knows where the prize is, opens one of the other two doors, revealing that the prize is not there.

The contestant can then change her/his choice, or not.

Should she/he change doors?

Before the host opens the door:

Sample space is $S = \{d_1, d_2, d_3\}$ with $\mathbb{P}(d_1) = \mathbb{P}(d_2) = \mathbb{P}(d_3) = \frac{1}{3}$.

The contestant chooses d_1 and has a $\frac{1}{3}$ chance of winning.

The host then opens door 2:

The sample space is now $S' = \{d_1, d_3\}$, but $\mathbb{P}(d_1)$ remains at $\frac{1}{3}$ (since the prize hasn't moved) so $\mathbb{P}(d_3) = \frac{2}{3}$ since the sum of the probabilities must be 1.

Thus the prize is twice as likely to be behind door 3 as behind door 1.

The Monty Hall problem

A game: Three doors, with a prize behind one of them.

The contestant chooses one door.

The host, who knows where the prize is, opens one of the other two doors, revealing that the prize is not there.

The contestant can then change her/his choice, or not.

Should she/he change doors?

Before the host opens the door:

Sample space is $S = \{d_1, d_2, d_3\}$ with $\mathbb{P}(d_1) = \mathbb{P}(d_2) = \mathbb{P}(d_3) = \frac{1}{3}$.

The contestant chooses d_1 and has a $\frac{1}{3}$ chance of winning.

The host then opens door 2:

The sample space is now $S' = \{d_1, d_3\}$, but $\mathbb{P}(d_1)$ remains at $\frac{1}{3}$ (since the prize hasn't moved) so $\mathbb{P}(d_3) = \frac{2}{3}$ since the sum of the probabilities must be 1.

Thus the prize is twice as likely to be behind door 3 as behind door 1.

So the contestant should change doors.

The birthday problem

In a group of 50 people, what is the probability that at least two people have the same birthday (assuming all birthdays are equally likely)?

The birthday problem

In a group of 50 people, what is the probability that at least two people have the same birthday (assuming all birthdays are equally likely)?

Let $D = \{1, \dots, 365\}$ represent the days of the year.

The birthday problem

In a group of 50 people, what is the probability that at least two people have the same birthday (assuming all birthdays are equally likely)?

Let $D = \{1, \dots, 365\}$ represent the days of the year.

Then the sample space is $S = D^{50} = \{(a_n)_{n=1, \dots, 50} \subseteq D\}$, representing all sequences of 50 birthdays.

The birthday problem

In a group of 50 people, what is the probability that at least two people have the same birthday (assuming all birthdays are equally likely)?

Let $D = \{1, \dots, 365\}$ represent the days of the year.

Then the sample space is $S = D^{50} = \{(a_n)_{n=1, \dots, 50} \subseteq D\}$, representing all sequences of 50 birthdays.

We have $|S| = |D^{50}| = |D|^{50} = 365^{50}$.

The birthday problem

In a group of 50 people, what is the probability that at least two people have the same birthday (assuming all birthdays are equally likely)?

Let $D = \{1, \dots, 365\}$ represent the days of the year.

Then the sample space is $S = D^{50} = \{(a_n)_{n=1, \dots, 50} \subseteq D\}$, representing all sequences of 50 birthdays.

We have $|S| = |D^{50}| = |D|^{50} = 365^{50}$.

The event E of 'at least two persons have the same birthday' is

$E = \{(a_n)_{n=1, \dots, 50} \subseteq D ; \exists j \neq k \in \{1, \dots, 50\} a_j = a_k\}$.

The birthday problem

In a group of 50 people, what is the probability that at least two people have the same birthday (assuming all birthdays are equally likely)?

Let $D = \{1, \dots, 365\}$ represent the days of the year.

Then the sample space is $S = D^{50} = \{(a_n)_{n=1, \dots, 50} \subseteq D\}$, representing all sequences of 50 birthdays.

We have $|S| = |D^{50}| = |D|^{50} = 365^{50}$.

The event E of 'at least two persons have the same birthday' is

$$E = \{(a_n)_{n=1, \dots, 50} \subseteq D ; \exists j \neq k \in \{1, \dots, 50\} a_j = a_k\}.$$

The complementary event 'everybody has a different birthday' is

$$E^c = \{(a_n)_{n=1, \dots, 50} \subseteq D ; \forall j, k \in \{1, \dots, 50\} j \neq k \implies a_j \neq a_k\}.$$

The birthday problem

In a group of 50 people, what is the probability that at least two people have the same birthday (assuming all birthdays are equally likely)?

Let $D = \{1, \dots, 365\}$ represent the days of the year.

Then the sample space is $S = D^{50} = \{(a_n)_{n=1, \dots, 50} \subseteq D\}$, representing all sequences of 50 birthdays.

We have $|S| = |D^{50}| = |D|^{50} = 365^{50}$.

The event E of 'at least two persons have the same birthday' is

$$E = \{(a_n)_{n=1, \dots, 50} \subseteq D ; \exists j \neq k \in \{1, \dots, 50\} a_j = a_k\}.$$

The complementary event 'everybody has a different birthday' is

$$E^c = \{(a_n)_{n=1, \dots, 50} \subseteq D ; \forall j, k \in \{1, \dots, 50\} j \neq k \implies a_j \neq a_k\}.$$

$$\mathbb{P}(E^c) = \frac{|E^c|}{|S|} = \frac{P(365, 50)}{365^{50}} = \frac{365!}{365^{50} \times 315!} \sim 0.03.$$

The birthday problem

In a group of 50 people, what is the probability that at least two people have the same birthday (assuming all birthdays are equally likely)?

Let $D = \{1, \dots, 365\}$ represent the days of the year.

Then the sample space is $S = D^{50} = \{(a_n)_{n=1, \dots, 50} \subseteq D\}$, representing all sequences of 50 birthdays.

We have $|S| = |D^{50}| = |D|^{50} = 365^{50}$.

The event E of 'at least two persons have the same birthday' is

$$E = \{(a_n)_{n=1, \dots, 50} \subseteq D ; \exists j \neq k \in \{1, \dots, 50\} a_j = a_k\}.$$

The complementary event 'everybody has a different birthday' is

$$E^c = \{(a_n)_{n=1, \dots, 50} \subseteq D ; \forall j, k \in \{1, \dots, 50\} j \neq k \implies a_j \neq a_k\}.$$

$$\mathbb{P}(E^c) = \frac{|E^c|}{|S|} = \frac{P(365, 50)}{365^{50}} = \frac{365!}{365^{50} \times 315!} \sim 0.03.$$

Thus

$$\mathbb{P}(E) = 1 - \mathbb{P}(E^c) \sim 0.97.$$

There is a 97% chance that two people will have the same birthday.