

Gene name: **C3AR1**

**External Ids for C3AR1 Gene :** HGNC: [131](#) NCBI Gene: [719](#) Ensembl: [ENSG00000171860](#) OMIM®: [605246](#) UniProtKB/Swiss-Prot: [Q16581](#)

**NCBI Gene Summary:** C3a is an anaphylatoxin released during activation of the **complement system**. The protein encoded by this gene is an orphan G protein-coupled receptor for C3a. Binding of C3a by the encoded receptor activates chemotaxis, granule enzyme release, superoxide anion production, and bacterial opsonization.

**GeneCards Summary:** C3AR1 (Complement C3a Receptor 1) is a Protein Coding gene. Diseases associated with C3AR1 include [Atypical Hemolytic-Uremic Syndrome](#) and [Prostate Cancer](#). Among its related pathways are [Class A/1 \(Rhodopsin-like receptors\)](#) and [Complement cascade](#). Gene Ontology (GO) annotations related to this gene include *G protein-coupled receptor activity* and *complement component C3a receptor activity*. An important paralog of this gene is [C5AR1](#).

**UniProtKB/Swiss-Prot Summary:** Receptor for the chemotactic and inflammatory peptide anaphylatoxin C3a. This receptor stimulates chemotaxis, granule enzyme release and superoxide anion production. ( [C3AR\\_HUMAN,Q16581](#) )

**Cellular localization:** Cell membrane; Multi-pass membrane protein is the most important.

**Full name:** Complement component 3a receptor 1 (C3AR1).

**Receptor type:** G protein-coupled receptor (GPCR).

**Ligand:** C3a, an anaphylatoxin generated during activation of the **complement system**.

**Expression:** Highly expressed on myeloid cells — particularly neutrophils, monocytes/macrophages, dendritic cells, and some endothelial and epithelial cells.

## **Biological Function of C3AR1**

- Binds **C3a**, a cleavage product of complement protein C3
- Activates **pro-inflammatory signaling cascades**, leading to:
  - **Chemotaxis** of neutrophils and macrophages
  - **Release of reactive oxygen species (ROS)**
  - **Cytokine and chemokine secretion**
  - **Increased vascular permeability**
- Plays a key role in **innate immunity**, particularly in **early response to infection**.

## Role in Sepsis

Sepsis is a dysregulated immune response to infection, and C3AR1 is central to the hyperinflammatory phase:

### Early-phase sepsis (hyperinflammatory):

- C3AR1 is upregulated in sepsis
- Drives excessive neutrophil activation and cytokine storms
- Associated with endothelial damage, vascular leakage, and multi-organ dysfunction

### Later-phase (immunosuppression):

- Some studies show compensatory downregulation or desensitization of C3AR1 as a mechanism to prevent tissue damage.

## Evidence in Sepsis Studies

### Diagnostic Value

- C3AR1 gene expression is significantly upregulated in whole blood transcriptome studies of sepsis patients compared to healthy controls.
- Detected early in sepsis onset, including in neonates and adults.

### Prognostic Value

- High C3AR1 expression is associated with:
  - Increased mortality risk.
  - Organ failure severity (SOFA scores).
- May help predict poor outcomes, especially in patients with Gram-negative bacteremia or systemic inflammation.

## Supporting Literature


DOI: 10.1097/MD.00000000000037519

DOI: 10.1038/s41598-024-59400-0

DOI: 10.3389/fimmu.2019.00543

DOI: 10.1111/jth.12956

KEGG :



Reactome results:

