



A

Shared primary warm-start-region selection



B

Selected-family joint-objective eligibility



C

Context-specific top-50 endpoint robustness

		Selected warm-start family	
		C01	C02
in vivo	Both ploidy regimes present at 0% O2	yes 100%	yes 100%
	Both ploidy regimes present at 1% O2	yes 100%	no 100%
	Both ploidy regimes present at 5% O2	no 100%	yes 100%
	Unmodified trajectory: O2 response direction	decrease 100%	decrease 100%
	Surface O2 response at low p_misseg	approximately flat 100%	increase 100%
	Surface O2 response at intermediate p_misseg	decrease 100%	approximately flat 100%
	Surface O2 response at high p_misseg	decrease 100%	decrease 100%
	At least half of the prespecified diagnostic union has spectral gap ≥ 0.005	no 100%	no 100%
in vitro	Both ploidy regimes present at 0% O2	yes 100%	yes 100%
	Both ploidy regimes present at 1% O2	yes 100%	yes 100%
	Both ploidy regimes present at 5% O2	yes 100%	yes 100%
	Unmodified trajectory: O2 response direction	increase 100%	increase 100%
	Surface O2 response at low p_misseg	approximately flat 100%	approximately flat 100%
	Surface O2 response at intermediate p_misseg	increase 100%	increase 100%
	Surface O2 response at high p_misseg	decrease 100%	decrease 100%
	At least half of the prespecified diagnostic union has spectral gap ≥ 0.005	yes 100%	yes 100%