Xiaoxingkai Transfer Result Report

Run directory: S:\STU-Papers\My_Papers\EXP1-Pre-Training-Standalone\outputs\runs\xiaoxingkai_transfer_daily12\xiaoxingkai_transfer_daily12_20260415_112733
Experiment: xiaoxingkai_transfer_daily12
Source file: S:\STU-Papers\My_Papers\Pre-Training-Data\LakeIce_ERA5_less60_XiaoxingkaiDaily12.xlsx
Target test lake: [Li] Lake Xiaoxingkai

Front Summary

Split	Loss	RMSE	MAE	R2	Bias	Negative Preds
val	40.9290	2.8430	1.3937	-280.0793	1.3642	0
test	7.6504	2.7659	1.4554	-1585.1862	1.0229	0

Per-Lake Numeric Test Summary

Lake	Count	RMSE	MAE	R2	Bias
[Li] Lake Xiaoxingkai	66	2.7659	1.4554	-1585.1862	1.0229

Key Diagnostic Notes

Best epoch: 1

Observed test mean/std: 0.7052 / 0.0694

Predicted test mean/std: 1.7281 / 2.5887

Validation windows: 36 | Test windows: 66

Interpretation: if predicted std is much smaller than observed std, the model is behaving too conservatively and is under-expressing peak-to-trough variation.

0. Training Inputs And Hyperparameters

The model consumes one irregular path per valid anchor row. Each path contains relative time plus meteorological and seasonal channels; the target total_ice_m is predicted, not fed back as an input feature.

Input Channels

1. relative_time
2. doy_sin
3. doy_cos
4. Ten_Meter_Elevation_Wind_Speed_meterPerSecond
5. Air_Temperature_celsius
6. Relative_Humidity_percent
7. Shortwave_Radiation_Downwelling_wattPerMeterSquared
8. Longwave_Radiation_Downwelling_wattPerMeterSquared
9. Sea_Level_Barometric_Pressure_pascal
10. Surface_Level_Barometric_Pressure_pascal
11. Precipitation_millimeterPerDay
12. Snowfall_millimeterPerDay

Core Training Configuration

Field	Value
window.window_days	15
coeffs.interpolation	hermite
model.method	midpoint
model.hidden_channels	48
model.hidden_hidden_channels	96
model.num_hidden_layers	2

Field	Value
train.batch_size	64
train.learning_rate	0.0002
train.weight_decay	0.0005
train.loss	huber
train.huber_delta	0.1
train.max_epochs	120
train.gradient_clip_norm	0.5
custom_split.val_fraction	0.1
custom_split.min_train_rows_per_lake	24
custom_split.min_val_rows_per_lake	8
custom_split.max_train_windows_per_lake	160
custom_split.max_val_windows_per_lake	20

1. Source-Lake Selection, Windowing, And Balance Effects

Each source lake is first split by time into train/val. The held-out target lake [Li] Lake Xiaoxingkai is kept entirely for test. A row only becomes a valid window if there are at least two observations inside the 15-day lookback. After that, train and val windows are capped per lake for balance.

Row Assignment Counts From The Source File

Lake	Rows In Source	Train Rows	Val Rows	Test Rows
[Li] Lake Xiaoxingkai	68	0	0	68
[Huo] Lake Hnagzhang	101	90	11	0
[Huo] Lake Ulansu	163	146	17	0
[IISD Experimental] Lake 239	160	144	16	0
[IISD Experimental] Lake 302	101	90	11	0
[North Temperate Lakes LTER] Allequash Lake	118	106	12	0
[North Temperate Lakes LTER] Big Muskellunge Lake	121	108	13	0
[North Temperate Lakes LTER] Bog 12-15 (Trout Bog)	124	111	13	0
[North Temperate Lakes LTER] Bog 27-2 (Crystal Bog)	121	108	13	0
[North Temperate Lakes LTER] Crystal Lake (NTL)	120	108	12	0
[North Temperate Lakes LTER] Sparkling Lake	121	108	13	0
[North Temperate Lakes LTER] Trout Lake	120	108	12	0

Window Eligibility And Balance Effects

Lake	Train Valid	Train Kept	Val Valid	Val Kept	Test Windows	Median Points
[Li] Lake Xiaoxingkai	0	0	0	0	66	16.0
[Huo] Lake Hnagzhang	89	89	11	11	0	16.0

Lake	Train Valid	Train Kept	Val Valid	Val Kept	Test Windows	Median Points
[Huo] Lake Ulansu	144	144	17	17	0	16.0
[IISD Experimental] Lake 239	32	32	3	3	0	2.0
[IISD Experimental] Lake 302	9	9	3	3	0	2.0
[North Temperate Lakes LTER] Allequash Lake	1	1	0	0	0	2.0
[North Temperate Lakes LTER] Big Muskellunge Lake	1	1	0	0	0	2.0
[North Temperate Lakes LTER] Bog 12-15 (Trout Bog)	0	0	1	1	0	2.0
[North Temperate Lakes LTER] Bog 27-2 (Crystal Bog)	0	0	0	0	0	0.0
[North Temperate Lakes LTER] Crystal Lake (NTL)	2	2	1	1	0	2.0
[North Temperate Lakes LTER] Sparkling Lake	0	0	0	0	0	0.0
[North Temperate Lakes LTER] Trout Lake	0	0	0	0	0	0.0

Total Counts

Total included source rows: 1438

Total train rows / val rows / test rows: 1227 / 143 / 68

Total train windows before/after balance: 278 / 278

Total val windows before/after balance: 36 / 36

Total test windows: 66

2. Core Result Figures

Training loss curve



Validation metric curves



Test predicted vs observed



Held-out Xiaoxingkai time series



Prediction vs observation distribution


Prediction Distribution Vs Observation Distribution
Observed
Predicted
Count
0
5
10
15
20
0
2
4
6
8
10
12
Ice thickness (m)