

# Modelling with Differential Equations 1

Semester 1, 2023

*Prof Matthew Simpson*

Tarang Janawalkar

This work is licensed under a Creative Commons  
“Attribution-NonCommercial-ShareAlike 4.0 International” license.



# Contents

<b>Contents</b>	<b>1</b>
<b>1 Differential Equations</b>	<b>2</b>
1.1 Ordinary and Partial Differential Equations . . . . .	2
1.2 Order . . . . .	2
1.3 Linearity . . . . .	2
1.4 Homogeneity . . . . .	3
1.5 Useful Theorems . . . . .	3
<b>2 Constant Coefficient ODEs</b>	<b>4</b>
2.1 Homogeneous Solution . . . . .	4
2.1.1 Real Distinct Roots . . . . .	5
2.1.2 Complex Conjugate Roots . . . . .	5
2.1.3 Real Repeated Roots . . . . .	5
2.2 Reduction of Order . . . . .	5
2.2.1 Obtaining the General Solution in a Repeated Roots Problem . . . . .	5
2.3 Fundamental Set of Solutions . . . . .	6
2.3.1 Test for Linear Independence . . . . .	7
2.3.2 The Wronskian . . . . .	7
2.4 Particular Solution . . . . .	8
2.4.1 Choosing a Particular Solution . . . . .	8
2.4.2 Undetermined Coefficients . . . . .	9
2.4.3 The Resonance Case . . . . .	9
2.5 Initial Value Problems . . . . .	9
<b>3 Nonconstant Coefficient ODEs</b>	<b>10</b>
3.1 Variation of Parameters . . . . .	10
<b>4 Power Series Solutions</b>	<b>12</b>
4.1 Classification of Points . . . . .	12
4.2 Frobenius Method . . . . .	13
4.2.1 Cases of Indicial Roots . . . . .	13
<b>5 Laplace Transforms</b>	<b>15</b>
5.1 Partial Fraction Decomposition . . . . .	15
5.2 Heaviside Step Function . . . . .	16
5.3 Properties of Laplace Transforms . . . . .	16
<b>6 Linear Systems</b>	<b>17</b>
6.1 Stability . . . . .	17
6.2 Linearisation of nonlinear systems . . . . .	17
6.3 Nullclines . . . . .	18
<b>A Second-Order Variation of Parameters</b>	<b>19</b>

# 1 Differential Equations

The study of differential equations (DE) finds techniques to solve equations relating unknown functions with their derivatives. These equations allow us to model and predict the behaviour of systems evolving over time or any other dimension.

## 1.1 Ordinary and Partial Differential Equations

Differential equations fall into one of two categories:

- *Ordinary differential equations* (ODE) — derivatives are taken with respect to only one variable.
- *Partial differential equations* (PDE) — derivatives are taken with respect to several variables.

Some examples of ODEs are shown below:

$$\begin{aligned}\frac{dy}{dt} &= ky && \text{(Exponential growth)} \\ \frac{dy}{dt} &= k(A - y) && \text{(Newton's law of cooling)} \\ m \frac{d^2x}{dt^2} + c \frac{dx}{dt} + kx &= f(t) && \text{(Mechanical vibrations)}\end{aligned}$$

where  $x$ ,  $y$ , and  $f$  are functions of time -  $t$ . Below are some examples of PDEs:

$$\begin{aligned}\frac{\partial u}{\partial t} + c \frac{\partial u}{\partial x} &= 0 && \text{(Transport equation)} \\ \frac{\partial u}{\partial t} &= k \frac{\partial^2 u}{\partial x^2} && \text{(Heat equation)} \\ \Delta u &= 0 && \text{(Laplace's equation)}\end{aligned}$$

where  $u$  is typically a function of space ( $x$ ,  $y$ ,  $z$ ) or space-time ( $x$ ,  $y$ ,  $z$ ,  $t$ ).

## 1.2 Order

The order of a differential equation refers to the highest derivative term that appears in the equation.

## 1.3 Linearity

A linear ODE is defined by a linear polynomial in the unknown function and its derivatives:

$$a_n(t)y^{(n)} + \dots + a_2(t)y'' + a_1(t)y' + a_0(t)y = f(t).$$

Here the notation  $y^{(n)}$  denotes the  $n$ th derivative of  $y$ , and we use this in-place of the apostrophe (') for compactness. Note that the function  $y$  itself, can be thought of as the "0"th derivative of  $y$ . Any ODE that cannot be expressed in the above form is nonlinear.

We can also classify PDEs in a similar manner. We will only consider a second-order linear PDE for brevity:

$$A(x, y)u_{xx} + B(x, y)u_{xy} + C(x, y)u_{yy} + D(x, y)u_x + E(x, y)u_y + F(x, y)u = f(x, y)$$

where we have a linear combination of second-order partial derivatives of  $u$ . Due to the complexity of PDEs, other classifications also exist, but these will not be discussed.

## 1.4 Homogeneity

Finally, a differential equation is homogeneous when it is an equation consisting only of the unknown function and its derivatives. In other words, there are no constant terms (or functions).

Both ODEs and PDEs are classified by the same rules, below is an example of homogeneous DEs:

$$ay'' + by' + cy = 0$$

$$\frac{\partial^2 u}{\partial x^2} + k \frac{\partial^2 u}{\partial t^2} = 0.$$

As we will see in further sections, differential equations are often simpler to solve when they are homogeneous.

## 1.5 Useful Theorems

**Theorem 1.5.1** (Existence and Uniqueness Theorem<sup>1</sup>). *Let the functions  $f$  and  $\frac{\partial f}{\partial y}$  be Lipschitz continuous<sup>2</sup> on the closed rectangle  $D \subseteq \mathbb{R}$  with  $(t_0, y_0) \in D$ . Then there exists some interval  $I \subseteq D$ , centred on  $t_0$  such that there is a unique solution to the initial value problem:*

$$\begin{cases} y'(t) = f(t, y) \\ y(t_0) = y_0 \end{cases}$$

**Corollary 1.5.1.1** (Existence and Uniqueness Theorem for Higher-Order ODEs). *Let the functions  $\frac{\partial^i f}{\partial y^i}$  for all  $i = 0, 1, \dots, n-1$ , be Lipschitz continuous on the closed  $(n+1)$ -dimensional hyperrectangle  $D \subseteq \mathbb{R} \times \mathbb{R}^n$  with  $(t_0, y_0, y'_0, y''_0, \dots, y_0^{(n-1)}) \in D$ . Then there exists some interval  $I \subseteq D$  containing  $t_0$ , such that there is a unique solution to the initial value problem:*

$$\begin{cases} y^{(n)}(t) = f(t, y, y', y'', \dots, y^{(n-1)}) \\ y^{(i)}(t_0) = y_0^{(i)} \end{cases}$$

for all  $i = 0, 1, \dots, n-1$ . Note that  $y_0^{(i)}$  is the constant value associated with the initial value of the  $i$ th derivative of  $y$  at  $t_0$ .

**Theorem 1.5.2** (Principle of Superposition). *If  $y_1$  is a solution to the equation*

$$ay'' + by' + cy = f_1(t),$$

and  $y_2$  are solutions to

$$ay'' + by' + cy = f_2(t),$$

then for any constants  $c_1$  and  $c_2$ , the linear combination  $y = c_1 y_1 + c_2 y_2$  is a solution to the differential equation

$$ay'' + by' + cy = c_1 f_1(t) + c_2 f_2(t).$$

<sup>1</sup>Also known as the Picard-Lindelöf Theorem

<sup>2</sup>When  $f$  and  $\frac{\partial f}{\partial y}$  are continuous, then a solution exists with no information about its uniqueness. See the Peano Existence Theorem for more information.

*Proof.* We can prove the above result through substitution,

$$\begin{aligned} ay'' + by' + cy &= a(c_1y_1 + c_2y_2)'' + b(c_1y_1 + c_2y_2)' + c(c_1y_1 + c_2y_2) \\ &= ac_1y_1'' + ac_2y_2'' + bc_1y_1' + bc_2y_2' + cc_1y_1 + cc_2y_2 \\ &= c_1(ay_1'' + by_1' + cy_1) + c_2(ay_2'' + by_2' + cy_2) \\ &= c_1f_1(t) + c_2f_2(t) \end{aligned}$$

A similar proof demonstrates that this principle also applies for higher-order linear ODEs.  $\square$

## 2 Constant Coefficient ODEs

A constant coefficient ODE is a linear ODE with coefficients that do not depend on the independent variable:

$$a_n y^{(n)} + \dots + a_2 y'' + a_1 y' + a_0 y = f(t).$$

Using the superposition principle, we can express solutions as

$$y = c_1 y_h + c_2 y_p.$$

where:

- $y_h$  is a solution to the *homogeneous* ODE, that is, when  $f(t) = 0$ , and is known as the “homogeneous solution”.
- $y_p$  is a solution to the *nonhomogeneous* ODE, and is known as the “particular solution”.

### 2.1 Homogeneous Solution

As the homogeneous ODE is linear, we will assume<sup>3</sup> that the solution  $y_h$  has the form

$$y_h(t) = e^{rt}.$$

Let us then substitute this into the homogeneous ODE:

$$\begin{aligned} a_n y_h^{(n)} + \dots + a_2 y_h'' + a_1 y_h' + a_0 y_h &= 0 \\ a_n r^n e^{rt} + \dots + a_2 r^2 e^{rt} + a_1 r e^{rt} + a_0 e^{rt} &= 0 \\ (a_n r^n + \dots + a_2 r^2 + a_1 r + a_0) e^{rt} &= 0 \\ a_n r^n + \dots + a_2 r^2 + a_1 r + a_0 &= 0 \end{aligned}$$

This result is known as the **characteristic equation** (also **complementary equation** or **auxiliary equation**) of our ODE. Finding the *roots* of this algebraic equation allows us to find  $y_h$ . As this is a polynomial expression, we expect three kinds of solutions:

- Real distinct roots:  $r_1, r_2, \dots, r_k \in \mathbb{R}$
- Complex conjugate roots:  $r, r^* \in \mathbb{C}$
- Real repeated roots:  $r \in \mathbb{R}$

For higher order ODEs that may have real, repeated, and complex roots simultaneously, we can take the linear combination of the solutions discussed below.

<sup>3</sup>A common term for such assumptions is also *ansatz*.

### 2.1.1 Real Distinct Roots

Given real distinct roots  $r_1, r_2, \dots, r_k$ , the homogeneous solution is given by

$$y_h = c_1 e^{r_1 t} + c_2 e^{r_2 t} \dots + c_k e^{r_k t}$$

for  $k \leq n$ .

### 2.1.2 Complex Conjugate Roots

Given complex conjugate roots  $r = \alpha + \beta i$  and  $r^* = \alpha - \beta i$ , the homogeneous solution is given by

$$y_h = k_1 e^{(\alpha + \beta i)t} + k_2 e^{(\alpha - \beta i)t}.$$

We can then simplify this solution using Euler's identity.

$$\begin{aligned} y_h &= e^{\alpha t} (k_1 e^{\beta i t} + k_2 e^{-\beta i t}) \\ &= e^{\alpha t} [k_1 (\cos(\beta t) + i \sin(\beta t)) + k_2 (\cos(\beta t) - i \sin(\beta t))] \\ &= e^{\alpha t} [k_1 \cos(\beta t) + i k_1 \sin(\beta t) + k_2 \cos(\beta t) - i k_2 \sin(\beta t)] \\ &= e^{\alpha t} [(k_1 + k_2) \cos(\beta t) + i (k_1 - k_2) \sin(\beta t)] \\ &= e^{\alpha t} [c_1 \cos(\beta t) + c_2 \sin(\beta t)] \end{aligned}$$

which demonstrates that complex conjugate roots correspond to oscillatory solutions.

### 2.1.3 Real Repeated Roots

When  $r = -\frac{a_1}{2a_2}$  is a repeated root with multiplicity  $k$ , a solution is given by

$$y_{h,1} = e^{rt}.$$

However, as we will see later, this is not sufficient.

## 2.2 Reduction of Order

The reduction of order method is a method for converting any linear ODE to another linear ODE of lower order. If the solution  $y_1$  is known in advance, we can construct additional solutions via the following substitution:

$$y_2 = u(t) y_1.$$

### 2.2.1 Obtaining the General Solution in a Repeated Roots Problem

In the previous section, we were unable to find a complete set of homogeneous solutions to our ODE. Therefore let us use this new method to find other solutions that satisfy the homogeneous equation.

$$y_{h,2} = u(t) y_{h,1}.$$

The following example uses a second-order ODE, but the process is applicable in general. Before we can substitute this equation into the ODE, let us find its first and second derivatives:

$$\begin{aligned} y'_{h,2} &= u' y_{h,1} + u y'_{h,1} \\ y''_{h,2} &= u'' y_{h,1} + u' y'_{h,1} + u' y'_{h,1} + u y''_{h,1} = u'' y_{h,1} + 2u' y'_{h,1} + u y''_{h,1} \end{aligned}$$

We can now substitute this equation into the homogeneous ODE:

$$\begin{aligned}
 a_2 y''_{h,2} + a_1 y'_{h,2} + a_0 y_{h,2} &= 0 \\
 a_2 (u'' y_{h,1} + 2u' y'_{h,1} + u y''_{h,1}) + a_1 (u' y_{h,1} + u y'_{h,1}) + a_0 u y_{h,1} &= 0 \\
 a_2 u'' y_{h,1} + 2a_2 u' y'_{h,1} + a_2 u y''_{h,1} + a_1 u' y_{h,1} + a_1 u y'_{h,1} + a_0 u y_{h,1} &= 0 \\
 a_2 u'' e^{rt} + 2a_2 u' r e^{rt} + a_2 u r^2 e^{rt} + a_1 u' e^{rt} + a_1 u r e^{rt} + a_0 u e^{rt} &= 0 \\
 [a_2 u'' + 2a_2 r u' + a_2 r^2 u + a_1 u' + a_1 r u + a_0 u] e^{rt} &= 0 \\
 a_2 u'' + (2a_2 r + a_1) u' + (a_2 r^2 + a_1 r + a_0) u &= 0
 \end{aligned}$$

As  $r = -\frac{a_1}{2a_2}$ , the coefficients of both  $u'$  and  $u$  become zero, resulting in:

$$a_2 u'' = 0$$

so that  $u'' = 0$  because  $a_2 \neq 0$ . Integrating twice gives the solution:

$$u(t) = k_1 t + k_2.$$

Let  $k_2 = 0$  so that:

$$y_{h,2} = t e^{rt}.$$

Finally, combining both solutions yields the general homogeneous solution:

$$y = c_1 e^{rt} + c_2 t e^{rt}.$$

It follows that this result applies to higher-order linear ODEs with repeated roots of any multiplicity.

*Proposition.*

A linear homogeneous ODE of order  $n$  with the root  $r$  of multiplicity  $k \leq n$  has the fundamental form:

$$y = e^{rt} (1 + t + t^2 + \dots + t^k).$$

## 2.3 Fundamental Set of Solutions

When determining the solutions of an  $n$ th order linear homogeneous ODE, we require a **general** solution  $y_h$  which consists of  $n$  linearly independent functions  $y_{h,i}$  (for  $i = 1, 2, \dots, n$ ) such that the set of solutions form a **fundamental set of solutions** over the problem domain.

As was alluded to before, this is done by taking the linear combination of each “basis” function  $y_{h,i}$ , to find the general homogeneous solution:

$$y_h = c_1 y_{h,1} + c_2 y_{h,2} + \dots + c_n y_{h,n}.$$

This is

To show that a set of functions are linearly independent, we must construct a definitive test, therefore, consider the following.

### 2.3.1 Test for Linear Independence

The set of  $(n-1)$ -times differentiable functions  $S = \{y_1, y_2, \dots, y_n\}$  is linearly independent iff

$$c_1 y_1 + c_2 y_2 + \dots + c_n y_n = 0$$

has the trivial solution  $c_1 = \dots = c_n = 0$ , for all  $t$ . Let us then differentiate this equation  $n-1$  times to find the following system of equations:

$$\begin{aligned} c_1 y_1 + c_2 y_2 + \dots + c_n y_n &= 0 \\ c_1 y_1' + c_2 y_2' + \dots + c_n y_n' &= 0 \\ c_1 y_1'' + c_2 y_2'' + \dots + c_n y_n'' &= 0 \\ &\vdots \\ c_1 y_1^{(n-1)} + c_2 y_2^{(n-1)} + \dots + c_n y_n^{(n-1)} &= 0 \end{aligned}$$

which we can express as

$$\begin{bmatrix} y_1 & y_2 & \dots & y_n \\ y_1' & y_2' & \dots & y_n' \\ y_1'' & y_2'' & \dots & y_n'' \\ \vdots & \vdots & \vdots & \vdots \\ y_1^{(n-1)} & y_2^{(n-1)} & \dots & y_n^{(n-1)} \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}.$$

Recall from Linear Algebra that the linear system:

$$\mathbf{A}\mathbf{x} = \mathbf{b}$$

has a unique solution when  $\text{rank}(\mathbf{A}) = \text{rank}([\mathbf{A} \mid \mathbf{b}]) = n$ , where  $\mathbf{b} = \mathbf{0}$ . This is true when the matrix  $\mathbf{A}$  is invertible, so that

$$\begin{aligned} \mathbf{A}\mathbf{c} &= \mathbf{0} \\ \mathbf{A}^{-1}\mathbf{A}\mathbf{c} &= \mathbf{A}^{-1}\mathbf{0} \\ \mathbf{c} &= \mathbf{0} \end{aligned}$$

as required.

Therefore we can conclude that when the determinant of  $\mathbf{A}$  is nonzero, the set  $S$  is linearly independent. Let us summarise these findings in the following section and introduce the Wronskian matrix.

### 2.3.2 The Wronskian

**Definition 2.1** (Wronskian Matrix). Given the set of  $(n-1)$ -times differentiable functions  $S = \{y_1, y_2, \dots, y_n\}$ , the Wronskian matrix  $\mathbf{W}$  is defined as

$$\mathbf{W} = \begin{bmatrix} y_1 & y_2 & \dots & y_n \\ y_1' & y_2' & \dots & y_n' \\ y_1'' & y_2'' & \dots & y_n'' \\ \vdots & \vdots & \vdots & \vdots \\ y_1^{(n-1)} & y_2^{(n-1)} & \dots & y_n^{(n-1)} \end{bmatrix}.$$



The “Wronskian” of these functions is given by

$$W = W[y_1, y_2, \dots, y_n] = \det(\mathbf{W}).$$

**Theorem 2.3.1** (Linear Independence and the Wronskian). *The set of  $(n-1)$ -times differentiable functions  $S = \{y_1, y_2, \dots, y_n\}$  is linearly independent if*

$$W[y_1, y_2, \dots, y_n] \neq 0.$$

*Note that the converse does not apply:  $W[y_1, y_2, \dots, y_n] = 0$  does not imply that  $S$  is linearly dependent.*

**Corollary 2.3.1.1.** *If the set  $S$  is linearly dependent, then*

$$W[y_1, y_2, \dots, y_n] = 0.$$

## 2.4 Particular Solution

As with the homogeneous solution, the method of *Undetermined Coefficients* requires us to guess a solution  $y_p$  based on the form of  $f(t)$ <sup>4</sup>.

### 2.4.1 Choosing a Particular Solution

Consider the vector space of particularly nice functions  $W$  with the basis:

for all  $n \in \mathbb{N}_{>0}$  and  $r \in \mathbb{R}_{>0}$ .

If  $f$  can be written as a linear combination or product of functions in  $W$ , then we can construct a particular solution  $y_p$  that satisfies the nonhomogeneous ODE. The table below shows the choice of  $y_p$  for various  $f$ .

Form of $f$	Particular Solution $y_p$
$P_n(t)$	$\sum_{i=0}^n A_i t^i$
$e^{rt}$	$C e^{rt}$
$T(\omega t)$	$K \cos(\omega t) + M \sin(\omega t)$
$P_n(t) e^{rt}$	$\left( \sum_{i=0}^n A_i t^i \right) e^{rt}$
$e^{rt} T(\omega t)$	$e^{rt} [K \cos(\omega t) + M \sin(\omega t)]$
$P_n(t) e^{rt} T(\omega t)$	$e^{rt} \left[ \left( \sum_{i=0}^n K_i t^i \right) \cos(\omega t) + \left( \sum_{i=0}^n M_i t^i \right) \sin(\omega t) \right]$

Table 1: Choice of particular solution based on the form of the nonhomogeneous term where  $n \in \mathbb{N}_{\geq 0}$ ,  $r, \omega \in \mathbb{R}_{>0}$ , and  $A, C, K, M \in \mathbb{R}$ . Any constant coefficients in  $f$  are to be ignored.

<sup>4</sup>  $f(t)$  is known as the **forcing function** or **input** in the system, and only depends on  $t$ .

In the above table,  $P_n(t)$  is any polynomial function of order  $n$ ,  $T(t)$  is used to denote any of the following trigonometric functions:

$$T(t) = \begin{cases} \cos(t) \\ \sin(t) \\ \cos(t) \pm \sin(t) \end{cases}$$

For more complex forms of  $f$  as shown in the last three rows of the table above, we can use the following facts:

- If  $f$  can be expressed as the linear combination of two forms  $f_1$  and  $f_2$ , then the particular solution is also a linear combination of the particular solutions associated with  $f_1$  and  $f_2$ .
- If  $f$  can be expressed as a product of two forms  $f_1$  and  $f_2$ , then the particular solution will also be the product of the particular solutions associated with  $f_1$  and  $f_2$ . In this situation, it is important to always simplify all arbitrary constants as shown in the table above.

In general, it is sufficient to only memorise the first three rows of the table above.

### 2.4.2 Undetermined Coefficients

Equipped with a particular solution  $y_p$ , we can substitute it into the LHS of the nonhomogeneous ODE to obtain:

$$a_n y_p^{(n)} + \dots + a_2 y_p'' + a_1 y_p' + a_0 y_p = f(t).$$

We then simplify the LHS and factor out any terms that depend on  $t$ . For example:

$$(\dots) f_1 + (\dots) f_2 + \dots + (\dots) f_3 f_4 + \dots = f.$$

The purpose of doing so is to construct a system of equations that allows us to solve for the undetermined coefficients in  $y_p$ .

### 2.4.3 The Resonance Case

In some problems, the forcing term  $f(t)$  may have a form similar to  $y_h$  leading to a linearly dependent solution  $y_p$ . In this case, we can use the *Reduction of Order* method, as we did for repeated roots, and multiply  $y_p$  by  $t$  until it is linearly independent to  $y_h$ .

## 2.5 Initial Value Problems

Real-world systems modelled using differential equations are accompanied by initial conditions that allow us to eliminate any remaining constants in our general solution *after* we have found **both** the homogeneous and particular solutions (where applicable).

For an  $n$ th order ODE, we require  $n$  initial conditions  $y_0$ , corresponding to the first  $n$  derivatives of  $y$ :

$$\begin{aligned} y(t_0) &= y_0 \\ y'(t_0) &= y'_0 \\ &\vdots \\ y^{(n-1)}(t_0) &= y_0^{(n-1)} \end{aligned}$$

note that  $y_0$  is a constant and we are using the dash ( $'$ ) for notational convenience.

### 3 Nonconstant Coefficient ODEs

A nonconstant coefficient ODE (or a linear ODE) has the following form:

$$a_n(t)y^{(n)} + \dots + a_2(t)y'' + a_1(t)y' + a_0(t)y = f(t).$$

where  $a_i$  depends on the independent variable. In the previous section, we found that the method of *Undetermined Coefficients* was insufficient for ODEs with more complex forcing functions. As we will see soon, we can construct a different particular solution where our guess depends on the homogeneous solutions.

#### 3.1 Variation of Parameters

Given a set of homogeneous solutions  $y_1, y_2, \dots$ , and  $y_n$  to the above linear ODE, consider a particular solution of the form

$$y_p = v_1(t)y_1 + v_2(t)y_2 + \dots + v_n(t)y_n$$

where  $v_i$  are now functions of the independent variable.

As before, we want to substitute this function into the linear ODE above, which requires us to compute the derivatives of  $y_p$ . Because  $y_p$  has several products, we will impose additional constraints such that we eliminate any derivatives of  $v$  in our calculations (except for in the final derivative)<sup>5</sup>. This will give:

$$\begin{aligned} y_p^{(k)} &= v_1 y_1^{(k)} + v_2 y_2^{(k)} + \dots + v_n y_n^{(k)} & k = 0, 1, \dots, n-1 \\ y_p^{(n)} &= v_1 y_1^{(n)} + v_2 y_2^{(n)} + \dots + v_n y_n^{(n)} + v_1' y_1^{(n-1)} + v_2' y_2^{(n-1)} + \dots + v_n' y_n^{(n-1)} \end{aligned}$$

where we make the following constraints:

$$0 = v_1' y_1^{(k)} + v_2' y_2^{(k)} + \dots + v_n' y_n^{(k)} \quad k = 0, 1, \dots, n-2.$$

Explicitly this looks like the following:

$$\begin{cases} y_p = v_1 y_1 + v_2 y_2 + \dots + v_n y_n \\ y_p' = v_1 y_1' + v_2 y_2' + \dots + v_n y_n' \\ \vdots \\ y_p^{(n-1)} = v_1 y_1^{(n-1)} + v_2 y_2^{(n-1)} + \dots + v_n y_n^{(n-1)} \\ y_p^{(n)} = v_1 y_1^{(n)} + v_2 y_2^{(n)} + \dots + v_n y_n^{(n)} + v_1' y_1^{(n-1)} + v_2' y_2^{(n-1)} + \dots + v_n' y_n^{(n-1)} \end{cases}$$

with the constraints:

$$\begin{cases} 0 = v_1' y_1 + v_2' y_2 + \dots + v_n' y_n \\ 0 = v_1' y_1' + v_2' y_2' + \dots + v_n' y_n' \\ \vdots \\ 0 = v_1' y_1^{(n-2)} + v_2' y_2^{(n-2)} + \dots + v_n' y_n^{(n-2)} \end{cases}$$

---

<sup>5</sup>See the appendix for an example using a second-order linear ODE.

We can now substitute  $y_p$  into the ODE to find

$$\begin{aligned}
& a_n y_p^{(n)} + a_{n-1} y_p^{(n-1)} + \dots + a_1 y_p' + a_0 y_p = f(t) \\
& a_0 [v_1 y_1 + v_2 y_2 + \dots + v_n y_n] + \\
& a_1 [v_1 y_1' + v_2 y_2' + \dots + v_n y_n'] + \\
& \vdots \\
& a_n [v_1 y_1^{(n)} + v_2 y_2^{(n)} + \dots + v_n y_n^{(n)}] + \\
& a_n [v_1' y_1^{(n-1)} + v_2' y_2^{(n-1)} + \dots + v_n' y_n^{(n-1)}] = f(t) \\
& v_1 [a_n y_1^{(n)} + a_{n-1} y_1^{(n-1)} + \dots + a_1 y_1' + a_0 y_1] + \\
& v_2 [a_n y_2^{(n)} + a_{n-1} y_2^{(n-1)} + \dots + a_1 y_2' + a_0 y_2] + \\
& \vdots \\
& v_n [a_n y_n^{(n)} + a_{n-1} y_n^{(n-1)} + \dots + a_1 y_n' + a_0 y_n] + \\
& a_n [v_1' y_1^{(n-1)} + v_2' y_2^{(n-1)} + \dots + v_n' y_n^{(n-1)}] = f(t)
\end{aligned}$$

This simplifies greatly and results in:

$$v_1' y_1^{(n-1)} + v_2' y_2^{(n-1)} + \dots + v_n' y_n^{(n-1)} = g(t)$$

where  $g(t) = f(t)/a_n$ . Using the constraints stated previously, we reach the following system of equations:

$$\begin{bmatrix} y_1 & y_2 & \dots & y_n \\ y_1' & y_2' & \dots & y_n' \\ \vdots & \vdots & \vdots & \vdots \\ y_1^{(n-2)} & y_2^{(n-2)} & \dots & y_n^{(n-2)} \\ y_1^{(n-1)} & y_2^{(n-1)} & \dots & y_n^{(n-1)} \end{bmatrix} \begin{bmatrix} v_1' \\ v_2' \\ \vdots \\ v_n' \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ g(t) \end{bmatrix}.$$

Using Cramer's rule, we find that the solution to this system is given by

$$v_i' = \frac{W_i}{W}$$

where  $W$  is the Wronskian, and  $W_i$  is the determinant of the Wronskian matrix where the  $i$ th column is replaced with the RHS vector  $\mathbf{b}$ . Integration<sup>6</sup> yields:

$$y_p = y_1 \int \frac{W_1}{W} dt + y_2 \int \frac{W_2}{W} dt + \dots + y_n \int \frac{W_n}{W} dt.$$

In this section, we built our particular solution from solutions to the homogeneous ODE; let us now discuss two methods to determine the homogeneous solution for linear ODEs.

---

<sup>6</sup>Due to the rational form of the integrand, a closed-form solution may not exist.

## 4 Power Series Solutions

Consider the following linear ODE:

$$y'' + P(t)y' + Q(t)y = f(t).$$

let us consider a power series solution of the form

$$y = \sum_{n=0}^{\infty} c_n t^n.$$

This allows us to form the following derivatives:

$$y' = \sum_{n=1}^{\infty} c_n n t^{n-1}, \quad y'' = \sum_{n=2}^{\infty} c_n n(n-1) t^{n-2} \quad \dots$$

As with constant coefficient ODEs, we can determine  $c_n$  through substitution:

$$\sum_{n=2}^{\infty} c_n n(n-1) t^{n-2} + P(t) \sum_{n=1}^{\infty} c_n n t^{n-1} + Q(t) \sum_{n=0}^{\infty} c_n t^n = f(t).$$

Here we must use the following steps to simplify this equation:

1. Shift the index  $n$  of any summation so that the exponent of  $t$  is the same in each summation.
2. Extract lower order terms from summations so that all summations start from the same index.
3. Factor all summations together.

This results in an equation of the form:

$$RR_0 t^p + \sum_{n=n_0}^{\infty} RR t^{n+q} = f(t).$$

where  $p$  and  $q$  are some integer value, and  $RR$  and  $RR_0$  depend on  $n$  and  $c_n$ .

We can solve this expression by equating linearly dependent terms w.r.t.  $t$ . As  $f(t)$  is not guaranteed to have a simple Taylor series expansion, we will typically consider the homogeneous problem first. This forces all coefficients of  $t$  to be equal to 0, and hence allows us to form a **recurrence relationship** using  $RR = 0$ . This relationship will be of the form:

$$c_{n+u} = v(n) c_n$$

where higher order terms depend on lower order terms ( $u > 0$  and  $v$  may depend on  $n$ ). By substituting  $n = n_0, n_0 + 1, n_0 + 2, \dots$ , we can substitute these coefficients back into the original power series, and, if possible, find an explicit form of the coefficients  $c_n$ .

### 4.1 Classification of Points

Values of  $t \in \mathbb{C}$  can be classified into **ordinary points**, at which the ODE's coefficients are analytic, and **singular points** at which some coefficient has a singularity (i.e., division by 0).

A singular point  $t = t_0$  is a **regular singular point** if

$$(t - t_0) P(t) \quad \text{and} \quad (t - t_0)^2 Q(t)$$

are analytic at  $t_0$ . A singular point that is not regular is called an **irregular singular point**.

## 4.2 Frobenius Method

When  $t = t_0$  forms a regular singular point, there exists at least one series solution of the form

$$y = \sum_{n=0}^{\infty} c_n (t - t_0)^{n+r}.$$

where  $r$  is some real number to be determined. This gives us the following derivatives:

$$y' = \sum_{n=0}^{\infty} c_n (n+r) t^{n+r-1}, \quad y' = \sum_{n=0}^{\infty} c_n (n+r)(n+r-1) t^{n+r-2} \quad \dots$$

Using a similar approach, we can substitute these expressions into our ODE to find a second relationship between the roots  $r$ , this is known as the **indicial equation**.

$$\text{IE} t^{r+p} + \sum_{n=n_0}^{\infty} \text{RR} t^{n+r+q} = f(t).$$

where again,  $p$  and  $q$  are some integers, and IE and RR depend on  $n$ ,  $r$ , and  $c_n$ .

### 4.2.1 Cases of Indicial Roots

In the event that the indicial equation yields several **indicial roots**, we must take the following into account. Given real roots  $r_1$  and  $r_2$  with  $r_1 > r_2$ :

1. When  $r_1 - r_2 \notin \mathbb{N}$ :

$$y_1(t) = \sum_{n=0}^{\infty} c_n t^{n+r_1}$$

$$y_2(t) = \sum_{n=0}^{\infty} b_n t^{n+r_2}$$

with  $c_0 \neq 0$  and  $b_0 \neq 0$ .

2. When  $r_1 - r_2 \in \mathbb{N}$ :

$$y_1(t) = \sum_{n=0}^{\infty} c_n t^{n+r_1}$$

$$y_2(t) = c y_1(t) \ln(t) + \sum_{n=0}^{\infty} b_n t^{n+r_2}$$

with  $c_0 \neq 0$  and  $b_0 \neq 0$ , where  $c$  may be 0.

3. When  $r_1 = r_2$ :

$$y_1(t) = \sum_{n=0}^{\infty} c_n t^{n+r_1}$$

$$y_2(t) = y_1(t) \ln(t) + \sum_{n=1}^{\infty} b_n t^{n+r_2}$$

with  $c_0 \neq 0$ .

When roots differ by an integer, notice that solutions may be linearly dependent. In this case, we can strategically choose roots that allow us to form a complete set of solutions. For instance, in the following indicial equation,

$$c_0 r(r+1)t^{r-1} + a_1(r+1)(r+2)t^r = 0$$

we can choose:

- $r = -1$ , where  $c_0$  and  $c_1$  are arbitrary,
- $r = 0$ , where  $c_1 = 0$  but  $c_0$  is arbitrary,
- $r = -2$ , where  $c_0 = 0$  but  $c_1$  is arbitrary.

By choosing  $r = -1$ , we mitigate the risk of reducing higher order terms that may depend on  $c_0$  or  $c_1$  to 0.

## 5 Laplace Transforms

The Laplace transform is a type of integral transform that maps a time-dependent problem in  $t$  to the complex  $s$ -domain, it is defined:

$$F(s) = \mathcal{L}\{f(t)\} = \int_0^{\infty} f(t) e^{-st} dt.$$

In the context of ODEs, this transformation allows us to take difficult nonlinear problems and solve them using the inverse Laplace transformation.

### 5.1 Partial Fraction Decomposition

In this unit, we will use tables to find the inverse Laplace transform, rather than using the explicit definition. To do so, we must make use of partial fraction decomposition (PFD).

Given the fraction

$$\frac{P(x)}{Q(x)}$$

where  $P$  and  $Q$  are polynomials in  $x$  and the degree of  $P$  is less than the degree of  $Q$ , we can decompose a fraction by considering the following cases:

Assume  $Q$  is fully factored.

**Case I:**  $Q$  has distinct polynomial factors:

$$Q(x) = q_1(x) q_2(x) \cdots q_k(x)$$

then,

$$\frac{P(x)}{Q(x)} = \frac{A_1(x)}{q_1(x)} + \frac{A_2(x)}{q_2(x)} + \cdots + \frac{A_k(x)}{q_k(x)}.$$

where  $A_i$  has the form:

$$A_i(x) = a_{i,n-1}x^{n-1} + a_{i,n-2}x^{n-2} + \cdots + a_{i,1}x + a_{i,0}$$

where the order of  $A_i$  will be one less than the order of the denominator  $q_i$ .

**Case II:**  $Q$  has repeated polynomial factors:

$$Q(x) = q(x)^r$$

then,

$$\frac{P(x)}{Q(x)} = \frac{A_1(x)}{q(x)} + \frac{A_2(x)}{q(x)^2} + \cdots + \frac{A_r(x)}{q(x)^r}.$$

where the denominator will have increasing powers of  $q$ .

**Case III:**  $Q$  has both distinct and repeated polynomial factors,  
the fraction will contain a combination of the above two cases.



## 5.2 Heaviside Step Function

The (Heaviside) step function (also unit step function) is defined:

$$u(t) = \begin{cases} 0 & t < 0 \\ 1 & 0 \leq t \end{cases}$$

along with the shifted step function:

$$u_c(t) = u(t - c) = \begin{cases} 0 & t < c \\ 1 & t \geq c. \end{cases}$$

These special functions allow us to express piecewise functions more compactly. For example,

$$f(t) = \begin{cases} f_1(t) & 0 \leq t < a \\ f_2(t) & t \geq a \end{cases}$$

can be expressed as

$$\begin{aligned} f(t) &= f_1(t) u(t - 0) \underbrace{- f_1(t) u(t - a)}_{\text{turn } f_1 \text{ off after } t = a} + f_2(t) u(t - a) \\ &= f_1(t) [u(t) - u(t - a)] + f_2(t) u(t - a). \end{aligned}$$

Additionally,

$$f(t) = \begin{cases} 0 & 0 \leq t < a \\ f_p(t) & a \leq t < b \\ 0 & t \geq b \end{cases}$$

can be expressed as

$$\begin{aligned} f(t) &= f_p(t) u(t - a) - f_p(t) u(t - b) \\ &= f_p(t) [u(t - a) - u(t - b)] \end{aligned}$$

To summarise

$$f_1(t) [u(t) - u(t - a)] + f_2(t) u(t - a) \equiv \begin{cases} f_1(t) & 0 \leq t < a \\ f_2(t) & t \geq a \end{cases}$$

and

$$f(t) [u(t - a) - u(t - b)] \equiv \begin{cases} 0 & 0 \leq t < a \\ f(t) & a \leq t < b \\ 0 & t \geq b \end{cases}$$

## 5.3 Properties of Laplace Transforms

## 6 Linear Systems

For a linear system

$$\mathbf{X}' = A\mathbf{X}$$

solutions are superpositions of the form

$$\mathbf{X}(t) = \mathbf{K}e^{\lambda t}$$

We can find such  $\lambda$  by solving  $|A - \lambda I|$  and substitute to find the corresponding  $\mathbf{K}$  vectors.

### 6.1 Stability

The characteristic polynomial is

$$\lambda^2 - \tau\lambda + \Delta = 0$$

where  $\tau = \text{Tr}(A)$  and  $\Delta = |A|$ .  $\mathbf{K}_i$  is the eigenvector corresponding to  $\lambda_i$ .

Condition		Behaviour	Solution
$\tau^2 - 4\Delta > 0$	$\tau < 0, \Delta > 0$	Stable node	$\mathbf{X}(t) = c_1\mathbf{K}_1e^{\lambda_1 t} + c_2\mathbf{K}_2e^{\lambda_2 t}$
	$\tau > 0, \Delta > 0$	Unstable node	
	$\Delta < 0$	Saddle point	
$\tau^2 = 4\Delta$	$\mathbf{K}_1 \neq a\mathbf{K}_2, \lambda_1 > 0$	Degenerate stable node	$\mathbf{X}(t) = c_1\mathbf{K}_1e^{\lambda_1 t} + c_2\mathbf{K}_2e^{\lambda_1 t}$
	$\mathbf{K}_1 \neq a\mathbf{K}_2, \lambda_1 < 0$	Degenerate unstable node	
	$\mathbf{K}_1 = a\mathbf{K}_2, \lambda_1 < 0$	Degenerate stable node	$\mathbf{X}(t) = c_1\mathbf{K}_1e^{\lambda_1 t} + c_2(\mathbf{K}_2e^{\lambda_1 t} + \mathbf{P}e^{\lambda_1 t})$
	$\mathbf{K}_1 = a\mathbf{K}_2, \lambda_1 > 0$	Degenerate unstable node	
$\tau^2 - 4\Delta < 0$	$\tau = 0$	Centre	$\mathbf{X} = c_1\mathbf{X}_1 + c_2\mathbf{X}_2, \mathbf{K}_1 = \mathbf{B}_1 + i\mathbf{B}_2$
	$\tau \neq 0, \alpha < 0$	Stable spiral node	$\mathbf{X}_1(t) = e^{\alpha t}(\mathbf{B}_1 \cos(\beta t) - \mathbf{B}_2 \sin(\beta t))$
	$\tau \neq 0, \alpha > 0$	Unstable spiral node	$\mathbf{X}_2(t) = e^{\alpha t}(\mathbf{B}_2 \cos(\beta t) + \mathbf{B}_1 \sin(\beta t))$

### 6.2 Linearisation of nonlinear systems

For the nonlinear system  $\frac{dx}{dt} = F(x, y)$ ,  $\frac{dy}{dt} = G(x, y)$ , there exists equilibrium or critical points where  $F(x, y) = G(x, y) = 0$ . If the system has a critical point at  $(x_e, y_e)$ , we can derive an approximate linear system around that point.

$$\begin{aligned} \dot{\mathbf{H}} &= \mathbf{J}|_{(x_e, y_e)} \mathbf{H} \\ \begin{pmatrix} x - x_e \\ y - y_e \end{pmatrix}' &= \begin{pmatrix} \frac{\partial F}{\partial x} & \frac{\partial F}{\partial y} \\ \frac{\partial G}{\partial x} & \frac{\partial G}{\partial y} \end{pmatrix} \bigg|_{(x_e, y_e)} \begin{pmatrix} x - x_e \\ y - y_e \end{pmatrix} \end{aligned}$$

An alternate approach is to use the chain rule directly,  $\frac{dx}{dy} = \frac{dx/dt}{dy/dt} = \frac{F(x, y)}{G(x, y)}$ .

### 6.3 Nullclines

For a linear system of differential equations  $\mathbf{X} = \begin{pmatrix} x \\ y \end{pmatrix}'$ , the curve along which  $x' = 0$  is called the  $x$ -nullcline, and the curve along which  $y' = 0$  is called the  $y$ -nullcline. The points where these nullclines intersect are steady states.

## A Second-Order Variation of Parameters

Consider the following second-order linear ODE

$$a_2(t)y'' + a_1(t)y' + a_0(t)y = f(t).$$

The particular solution is given by

$$y_p = v_1(t)y_1 + v_2(t)y_2$$

where  $y_1$  and  $y_2$  are solutions to the homogeneous problem. Differentiating  $y_p$  once gives us

$$y'_p = v'_1y_1 + v_1y'_1 + v'_2y_2 + v_2y'_2$$

where we will use the assumption

$$v'_1y_1 + v'_2y_2 = 0$$

to arrive at

$$y'_p = v_1y'_1 + v_2y'_2.$$

The second derivative is then

$$y''_p = v'_1y'_1 + v_1y''_1 + v'_2y'_2 + v_2y''_2.$$

If we then substitute these values into our ODE, and factor in terms of  $v$ , we find

$$\begin{aligned} a_2y''_p + a_1y'_p + a_0y_p &= f(t) \\ a_2[v'_1y'_1 + v_1y''_1 + v'_2y'_2 + v_2y''_2] + a_1[v_1y'_1 + v_2y'_2] + a_0[v_1y_1 + v_2y_2] &= f(t) \\ v_1[a_2y''_1 + a_1y'_1 + a_0y_1] + v_2[a_2y''_2 + a_1y'_2 + a_0y_2] + a_2v'_1y'_1 + a_2v'_2y'_2 &= f(t) \\ a_2v'_1y'_1 + a_2v'_2y'_2 &= f(t) \\ v'_1y'_1 + v'_2y'_2 &= g(t) \end{aligned}$$

where  $g(t) = f(t)/a_2(t)$ . This gives us a system of equations where we can solve for  $v'_1$  and  $v'_2$ , using the assumption from earlier:

$$\begin{bmatrix} y_1 & y_2 \\ y'_1 & y'_2 \end{bmatrix} \begin{bmatrix} v'_1 \\ v'_2 \end{bmatrix} = \begin{bmatrix} 0 \\ g(t) \end{bmatrix}.$$

where the coefficient matrix is simply the Wronskian matrix  $\mathbf{W}$ . Using Cramer's rule, we can conclude that

$$v'_1 = \frac{W_1}{W} \quad \text{and} \quad v'_2 = \frac{W_2}{W}$$

where  $W_i$  are the determinants of the Wronskian with the  $i$ th columns replaced with the RHS vector  $\mathbf{b}$ :

$$W_1 = \begin{vmatrix} 0 & y_2 \\ g(t) & y'_2 \end{vmatrix} \quad \text{and} \quad W_2 = \begin{vmatrix} y_1 & 0 \\ y'_1 & g(t) \end{vmatrix}$$

Integrating these results gives

$$v_1 = \int \frac{W_1}{W} dt \quad \text{and} \quad v_2 = \int \frac{W_2}{W} dt$$

so that the particular solution is given by

$$y_p = y_1 \int \frac{W_1}{W} dt + y_2 \int \frac{W_2}{W} dt.$$

Note that it may not be possible to compute these integrals.