

Open high-level data formats and software for gamma-ray astronomy

Christoph Deil^{1,a)}, Catherine Boisson^{3,b)}, Jeremy Perkins¹², Johannes King¹, Peter Eger¹, Michael Mayer⁶, Matthew Wood¹³, Victor Zabalza¹⁵, Jürgen Knödlseider¹¹, Tarek Hassan¹⁰, Lars Mohrmann⁵, Alexander Ziegler⁵, Bruno Khelifi⁴, Daniela Dorner⁵, Gernot Maier⁷, Giovanna Pedalletti⁷, Jaime Rosado¹⁰, José Luis Contreras¹⁰, Julien Lefaucheur³, Kai Brügge⁵, Mathieu Servillat³, Régis Terrier⁴, Roland Walter⁸ and Saverio Lombardi¹⁴

^{a)}Corresponding author: Christoph.Deil@mpi-hd.mpg.de

^{b)}Corresponding author: catherine.boisson@obspm.fr

¹*MPIK, Heidelberg, Germany*

²*NASA/GSFC, USA*

³*LUTH, Observatoire de Paris, Meudon, France*

⁴*APC, University of Paris 7, France*

⁵*FAU, Erlangen, Germany*

⁶*Humboldt University, Berlin, Germany*

⁷*DESY, Zeuthen, Germany*

⁸*Observatoire de Genève, 51 chemin des Maillettes, 1290 Sauverny, Switzerland*

⁹*Universidad Complutense de Madrid*

¹⁰*Institut de Física d'Altes Energies (IFAE), The Barcelona Institute of Science and Technology, Campus UAB, 08193 Bellaterra (Barcelona) Spain*

¹¹*IRAP, Toulouse, France*

¹²*NASA/GSFC*

¹³*SLAC National Accelerator Laboratory*

¹⁴*INAF, Osservatorio Astronomico di Roma, via Frascati 33, 00040 Monte Porzio Catone (Roma), Italy*

¹⁵*University of Leicester, UK*

Abstract. In gamma-ray astronomy, a variety of data formats and proprietary software exist, often developed for one specific mission or experiment. Especially for ground-based imaging atmospheric Cherenkov telescopes (IACTs), data and software have been so far mostly private to the collaborations operating the telescopes. However, there is a general movement in science towards open data and software and the next big IACT array, the Cherenkov Telescope Array (CTA), will be operated as an open observatory.

We have created a Github organisation at <https://github.com/open-gamma-ray-astro> where we are developing high-level data format specifications. A public mailing list was set up at <https://lists.nasa.gov/mailman/listinfo/open-gamma-ray-astro> and a first face-to-face meeting on the IACT high-level data model and formats took place in April 2016 in Meudon (France). The hope is that this open multi-mission effort will help to accelerate the development of open data formats and open-source software for gamma-ray astronomy, leading to synergies in the development of analysis codes and eventually better scientific results (reproducible, multi-mission). This poster will summarize what we have done so far, and has the goal to solicit comments and future contributions from the gamma-ray astronomy community.

TODO

We should probably mention these?




FIGURE 1. The purpose of the `gamma-astro-data-formats` effort is to encourage communication between high-level gamma-ray data producers, science tool developers and data analysts. The goal is to develop and agree on some common data models and formats, to avoid duplication of efforts and confusion by astronomer that want to work with gamma-ray data from different telescopes or try different tools.

- Mention how we work together besides Github: monthly telcons + planned bi-yearly f2f meetings?
- Is there anything we can cite for CTA data challenge 1 or data challenge 2 plans?
- Mention website and some info on how CTA is releasing IRF files at the moment (prod2)?
- The Instrument Response Function Format for the Cherenkov Telescope Array [1] - has now merged with this effort.
- Previous work to define a gamma-ray lightcurve format: [2]

Introduction

Science is aiming towards open and reproducible research. In gamma-ray astronomy, especially for ground-based imaging atmospheric Cherenkov telescopes (IACT), data and software have so far been mostly private to the collaboration operating the telescopes. The Cherenkov Telescope Array (CTA), the next generation of IACT, will be operated as an open observatory, meaning that data and analysis softwares will be public at the end-user, science tools level. Momentum was thus given to the development of open data format and open-source software for gamma-ray astronomy, leading to synergies between experiments, ground based and space missions.

This is illustrated in Figure 1: there are many gamma-ray data producers and science tools to work with gamma-ray data, so agreeing on common data models and formats that cover most common analysis cases makes things simpler for data producers, tool developers and users.

Open source analysis tools for VHE gamma-ray astronomy have emerged. They all meet on common ground of using FITS files for data transfer. The current IACTs (H.E.S.S., MAGIC and VERITAS) all use their own software based on the ROOT library, which makes it impossible to analyze the respective data with anything else than the corresponding software. Current open source analysis tools provide alternative analysis techniques compared to the present standard in VHE astronomy. It is assumed that these techniques (e.g. 3D likelihood analysis such as already implemented for Fermi-LAT) improve the sensitivity of IACTs by roughly 20%. The data model for CTA is currently being developed but still work in progress. A main point of defining the high level data format for CTA is to understand the instrument including its systematics and IRF dependencies. In addition the needs from a user perspective have to be taken into account to create a solution as simple as possible. Having agreed on a common data format for files and on a way how to store and access those files (folder structure), makes mid-level (event energies, positions) and high-level (source position, morphology, spectrum) checks between the different chains, algorithms and open-source tools possible. This will also ease interoperability with other codes (e.g. to check results, combine results in one plot, ...). Currently two open-source science tools packages are being designed for current IACT and CTA data analysis, Gammapy ([3]) and Ctools ([4]). Gammapy is an in-development Astropy-affiliated package. Ctools is based on the GammaLib analysis framework, which is mainly written in C++.




FIGURE 2. *Left:* gamma-astro-data-formats Github issue tracker with ongoing discussions. *Right:* latest version of the gamma-astro-data-formats specifications on Read the Docs (PDF and older tagged versions also available).

TODO: also mention other codes: pointlike [5], Naima [6], 3ML [7], Fermipy¹, Fermi ScienceTools.

We have created a Github organization where we are developing high-level data format specifications, in accordance with astronomy standards.

Resources and Process

The specifications of a given data level format defines the names and semantics of data and header fields. Such specifications are made easy to understand. Specifications of this format are currently written that can form the basis for prototyping for data producers (mainly existing IACTs and simulated CTA data) and consumers (mainly science tool codes). We include example files and some explanations, in addition to the detailed specifications for a given format.

The scope is high-level data, starting with event lists and instrument response functions (IRFs), what is called "data level 3" (DL3) in CTA. The first stable release (archived on Zenodo with a DOI) is coming soon.

If you want to contribute, it is simple:

- Use the existing format and give feedbacks. Propose additions and changes.
- Join the mailing list (see next section). Send an e-mail with an idea or proposal.
- Create a Github account. File an issue with a correction or make a pull request proposing additions.

No formal approval process is in place yet as this is a very recent effort.

- Mailing list for announcements and high level discussions (75 members, including people from all major gamma-ray collaborations :
<https://lists.nasa.gov/mailman/listinfo/open-gamma-ray-astro>
- Github issues and pull requests are used for detailed discussions:
<https://github.com/open-gamma-ray-astro/gamma-astro-data-formats>
- Data format specifications in HTML and PDF format, including example files:
<https://gamma-astro-data-formats.readthedocs.io/>

The specs are written in a markup format called "restructured text" (RST), which gets transformed by Sphinx to HTML or PDF, the latest rendered version is always available online (see Figure 2).

¹<https://github.com/fermiPy/fermipy>

Data models and formats

This section gives an overview of the status and plans for the gamma-ray data model and formats. As mentioned before, this effort was only started very recently and none of the formats should be considered stable. The next two sections will describe the effort to define an event data model and format (DL3), and higher-level formats for images, spectra and lightcurves (DL4) (i.e. content split as already illustrated in Figure 1)

In the data specification document we have created a "general" section that gives precise definitions of common quantities, such as precise definitions of time scales as well as coordinates systems. One example is a precise definition of AZIMUTH and ALTITUDE. We define AZIMUTH to be oriented east of north, and ALTITUDE to be relative to the zenith direction (not the horizon plane or a reference earth ellipsoid) and without applying a refraction correction.

There are some general questions we are discussing about where to be specific or flexible in our format specifications. One example is whether our data format specifications should be tied to FITS (and e.g. say the data type of a column is 1E, which is the data type code for 32-bit float in FITS), or whether it would be better to only say that the data type is "float", implicitly allowing the storage of this column as 32-bit or 64-bit float, and being able to store the data e.g. in text ECSV files in addition to binary FITS files. Another contentious point is whether physical units should be fixed (e.g. "MeV" or "TeV"), or whether the units should be flexible and only serialization formats that support declaration of units (such as FITS or VOTABLE or ECSV) should be supported and science tools are expected to process the unit information correctly.

Data level 3 (DL3) specifications

DL3 format should describe all data released to the end users for analysis with science tools as , EVENT, IRF and TECH (technical data not directly attached to the object observed). Figure 3 illustrates some of the DL3 data content. Prototyping by existing IACTs (H.E.S.S., MAGIC, VERITAS and science tools (Gammapy, Ctools) is a major activity.

Many points are still under discussion/consideration:

- Observations modes, time intervals
- How to link EVENT and IRF
- Pointing and live time information
- IRF axis and validity ranges
- FoV coordinates

Data level 4 & 5 specifications

A format is developed to store spectral analysis results for exchange and publication, not only "flux points" and "upper limits", but also the full likelihood profiles (see examples in Figure 4).

It was first developed in Fermipy² and used for Fermi-LAT (ref). It is now being adopted by ground-based high energy astronomy. Likelihood profiles are important information when those data are to be used in global spectral energy distribution (SED) of a source.

Description of light curves is another topic open for discussion.

Conclusions

- High-level gamma-ray data (DL 3 and up) from different telescopes is very similar, there's always event lists and IRFs, plus some extra info like good time interval and pointing information.
- We have started the first effort to define open data models and data formats for gamma-ray astronomy.
- The motivations for this are the development of open-source tools (Gammapy, ctools, Fermi ST, Fermipy, pointlike, emcee) to analyse gamma-ray data, and that IACTs are starting to produce data mostly in FITS format that these tools shall consume. So "many tools" and "many telescopes" makes a common data model and formats useful.
- We have chosen an open process (Mailing list, Github, monthly telcons, bi-yearly f2f meetings).
- There are useful preliminary specs, we encourage everyone to have a look now and give feedback and contribute. Many important questions are under discussion (refer back to section listing those).
- But there's no stable version or "standard" yet. Especially for CTA the process will have to be more formalised if CTA data is to be released in those formats.

²<https://github.com/fermipy/fermipy>




FIGURE 3. Illustration of major components of IACT (“imaging atmospheric Cherenkov telescope”, here a H.E.S.S. 1 Crab nebula observation) DL3 (“data level 3”) data. The **EVENTS** are stored as a table with the most important parameters shown. To derive spectra and morphology measurements of astrophysical sources, IRFs (“instrument response functions”) are used: the effective area (AEFF), energy dispersion (EDISP) and point spread function (PSF). Sometimes background (BKG) models are also created and released as part of a DL3 data release, and can be treated like an IRF, and sometimes deriving background models is left to the science tools. Note that this picture is not complete, see the “IACT DL3” section.

Acknowledgements

We would like to thank everyone that has contributed to or supported this effort, be it directly via contributions to the format specification, or indirectly via feedback or adopting the existing formats and spending the effort to transform their existing data to the common formats defined here.

We would also like to thank the following services make this form of collaboration possible: Github for collaboration, Sphinx as documentation system and Read the docs for building and hosting the HTML and PDF version of the spec.

REFERENCES

- [1] J. E. Ward, J. Rico, T. Hassan, and f. t. CTA Consortium, ArXiv e-prints August (2015), arXiv:1508.07437 [astro-ph.IM] .
- [2] M. Tluczykont, E. Bernardini, K. Satalecka, R. Clavero, M. Shayduk, and O. Kalekin, AAP **524**, p. A48December (2010), arXiv:1010.5659 [astro-ph.HE] .
- [3] A. Donath, C. Deil, M. P. Arribas, J. King, E. Owen, R. Terrier, I. Reichardt, J. Harris, R. Bühler, and S. Klepser, ArXiv e-prints September (2015), arXiv:1509.07408 [astro-ph.IM] .




FIGURE 4. Gamma-ray “data level 4” examples. *Left:* spectral energy distribution (SED) likelihood profiles (green), with flux points and upper limits as well as a best-model fit overplotted. *Right:* Lightcurve of 3FGL J0349.9-2102 from the third Fermi-LAT catalog.

- [4] J. Knödlseder, M. Mayer, C. Deil, J.-B. Cayrou, E. Owen, N. Kelley-Hoskins, C.-C. Lu, R. Buehler, F. Forest, T. Louge, H. Siejkowski, K. Kosack, L. Gerard, A. Schulz, P. Martin, D. Sanchez, S. Ohm, T. Hassan, and S. Brau-Nogu  , AAP **593**, p. A1August (2016), arXiv:1606.00393 [astro-ph.IM] .
- [5] M. Kerr, “Likelihood methods for the detection and characterization of gamma-ray pulsars with the Fermi large area telescope,” Ph.D. thesis, University of Washington 2010.
- [6] V. Zabalza, ArXiv e-prints September (2015), arXiv:1509.03319 [astro-ph.HE] .
- [7] G. Vianello, R. J. Lauer, P. Younk, L. Tibaldo, J. M. Burgess, H. Ayala, P. Harding, M. Hui, N. Omodei, and H. Zhou, ArXiv e-prints July (2015), arXiv:1507.08343 [astro-ph.HE] .