

Continuous-Time Fourier Transform

Signals and Systems: Experiment 2

Yuzhou Li

yuzhouli@hust.edu.cn

School of Electronic Information and Communications

April 9, 2024

Overview

1. Symbolic Function
2. Loop Calculation
3. Vector Product
4. Matrix Product
5. Experiment Problem

Overview

1. Symbolic Function
2. Loop Calculation
3. Vector Product
4. Matrix Product
5. Experiment Problem

Definitions of CTFT and ICTFT

Continuous-time Fourier transform (CTFT) is given by

$$F(\omega) = \int_{-\infty}^{\infty} f(t) e^{-j\omega t} dt \quad (1)$$

Definitions of CTFT and ICTFT

Continuous-time Fourier transform (CTFT) is given by

$$F(\omega) = \int_{-\infty}^{\infty} f(t) e^{-j\omega t} dt \quad (1)$$

Inverse continuous-time Fourier transform (ICTFT) is defined as

$$f(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} F(\omega) e^{j\omega t} d\omega \quad (2)$$

Example 1: Symbolic Functions for CTFT and ICTFT

Symbolic functions in MATLAB: `fourier()` and `ifourier()`.

```
syms t;    % define a symbol t
FT0 = fourier(cos(t))    % calculate the FT of cos t
f1 = dirac(t);    % calculate the FT of  $\delta(t)$ 
FT1 = fourier(f1)
f2 = heaviside(t);    % calculate the FT of  $u(t)$ 
FT2 = fourier(f2)
syms t0;    % calculate the FT of  $u(t - t_0)$ 
FT3 = fourier(heaviside(t - t0))
```

Overview

1. Symbolic Function
2. Loop Calculation
3. Vector Product
4. Matrix Product
5. Experiment Problem

Loop Calculation for CTFT-1

Consider the main-value interval $[t_1, t_2]$

$$F(w) = \int_{-\infty}^{\infty} f(t) e^{-j\omega t} dt = \int_{t_1}^{t_2} f(t) e^{-j\omega t} dt \quad (3)$$

Loop Calculation for CTFT-1

Consider the main-value interval $[t_1, t_2]$

$$F(w) = \int_{-\infty}^{\infty} f(t) e^{-j\omega t} dt = \int_{t_1}^{t_2} f(t) e^{-j\omega t} dt \quad (3)$$

Define the interval length $T = t_2 - t_1$ and let N be the time-domain sampling number, then the sampling interval $\Delta t = \frac{T}{N}$

$$F(w) = \frac{T}{N} \sum_{n=0}^{N-1} f(t_1 + n\Delta t) e^{-j\omega(t_1 + n\Delta t)} \quad (4)$$

Loop Calculation for CTFT-1

Consider the main-value interval $[t_1, t_2]$

$$F(w) = \int_{-\infty}^{\infty} f(t) e^{-j\omega t} dt = \int_{t_1}^{t_2} f(t) e^{-j\omega t} dt \quad (3)$$

Define the interval length $T = t_2 - t_1$ and let N be the time-domain sampling number, then the sampling interval $\Delta t = \frac{T}{N}$

$$F(w) = \frac{T}{N} \sum_{n=0}^{N-1} f(t_1 + n\Delta t) e^{-j\omega(t_1 + n\Delta t)} \quad (4)$$

Consider $w \in [w_1, w_2]$ and K frequency-domain samples: $\Omega = w_2 - w_1$ and $\Delta w = \frac{\Omega}{K}$

$$F(w_1 + k\Delta w) = \frac{T}{N} \sum_{n=0}^{N-1} f(t_1 + n\Delta t) e^{-j(w_1 + k\Delta w)(t_1 + n\Delta t)} \quad (5)$$

Loop Calculation for CTFT-2

Formula for CTFT calculation in MATLAB

$$F(w_1 + k\Delta w) = \frac{T}{N} \sum_{n=0}^{N-1} f(t_1 + n\Delta t) e^{-j(w_1 + k\Delta w)(t_1 + n\Delta t)}$$

Pseudocode for loop calculation

```
for k = 1, ..., K
    for n = 1, ..., N
        F[k, n] = F[k, n - 1] + \frac{T}{N} f(t_1 + n\Delta t) e^{-j(w_1 + k\Delta w)(t_1 + n\Delta t)}
    end
end
```

Loop Calculation for ICTFT-1

Inverse continuous-time Fourier transform is given by

$$f(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} F(w) e^{j\omega t} dw = \frac{\Omega}{2\pi K} \sum_{k=0}^{K-1} F(w_1 + k\Delta w) e^{j(w_1 + k\Delta w)t} \quad (6)$$

Loop Calculation for ICTFT-1

Inverse continuous-time Fourier transform is given by

$$f(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} F(w) e^{j\omega t} dw = \frac{\Omega}{2\pi K} \sum_{k=0}^{K-1} F(w_1 + k\Delta w) e^{j(w_1 + k\Delta w)t} \quad (6)$$

Discretize the time-domain signal

$$f(t_1 + n\Delta t) = \frac{\Omega}{2\pi K} \sum_{k=0}^{K-1} F(w_1 + k\Delta w) e^{j(w_1 + k\Delta w)(t_1 + n\Delta t)} \quad (7)$$

Loop Calculation for ICTFT-2

Formula for ICTFT calculation in MATLAB

$$f(t_1 + n\Delta t) = \frac{\Omega}{2\pi K} \sum_{k=0}^{K-1} F(w_1 + k\Delta w) e^{j(w_1 + k\Delta w)(t_1 + n\Delta t)}$$

Pseudocode for loop calculation

```
for  $n = 1, \dots, N$ 
    for  $k = 1, \dots, K$ 
         $f[n, k] = f[n, k - 1] + \frac{\Omega}{2\pi K} F(w_1 + k\Delta w) e^{j(w_1 + k\Delta w)(t_1 + n\Delta t)}$ 
    end
end
```

Example 1: Loop calculation for CTFT and ICTFT

Rectangular pulse

$$f(t) = \begin{cases} 1, & |x| < \frac{1}{2} \\ 0, & \text{otherwise} \end{cases} \quad (8)$$

- (1) Plot $f(t)$, $t \in [-1, 1]$;
- (2) Plot $F(w)$, $w \in [-8\pi, 8\pi]$;
- (3) Recover $f(t)$ from $F(w)$.




Figure 2.1: Waveform and spectrum

Overview

1. Symbolic Function
2. Loop Calculation
3. Vector Product
4. Matrix Product
5. Experiment Problem

Vector Product for CTFT

Formula for CTFT calculation in MATLAB

$$F(w_1 + k\Delta w) = \frac{T}{N} \sum_{n=0}^{N-1} f(t_1 + n\Delta t) e^{-j(w_1 + k\Delta w)(t_1 + n\Delta t)}$$

Vector Product for CTFT

Formula for CTFT calculation in MATLAB

$$F(w_1 + k\Delta w) = \frac{T}{N} \sum_{n=0}^{N-1} f(t_1 + n\Delta t) e^{-j(w_1 + k\Delta w)(t_1 + n\Delta t)}$$

Vector form for CTFT calculation

$$F(w_1 + k\Delta w) = \frac{T}{N} \begin{bmatrix} e^{-j(w_1 + k\Delta w)t_1} & e^{-j(w_1 + k\Delta w)(t_1 + \Delta t)} & \dots & e^{-j(w_1 + k\Delta w)(t_2 - \Delta t)} \end{bmatrix} \begin{bmatrix} f(t_1) \\ f(t_1 + \Delta t) \\ \vdots \\ f(t_2 - \Delta t) \end{bmatrix} \quad (9)$$

Vector Product for CTFT

Formula for CTFT calculation in MATLAB

$$F(w_1 + k\Delta w) = \frac{T}{N} \sum_{n=0}^{N-1} f(t_1 + n\Delta t) e^{-j(w_1 + k\Delta w)(t_1 + n\Delta t)}$$

Vector form for CTFT calculation

$$F(w_1 + k\Delta w) = \frac{T}{N} \begin{bmatrix} e^{-j(w_1 + k\Delta w)t_1} & e^{-j(w_1 + k\Delta w)(t_1 + \Delta t)} & \dots & e^{-j(w_1 + k\Delta w)(t_2 - \Delta t)} \end{bmatrix} \begin{bmatrix} f(t_1) \\ f(t_1 + \Delta t) \\ \vdots \\ f(t_2 - \Delta t) \end{bmatrix} \quad (9)$$

Pseudocode for vector product

```
for  $k = 1, \dots, K$   
     $F_k = \frac{T}{N} \mathbf{a}_k^T \mathbf{f}$   
end
```

Vector Product for ICTFT

Formula for ICTFT calculation in MATLAB

$$f(t_1 + n\Delta t) = \frac{\Omega}{2\pi K} \sum_{k=0}^{K-1} F(w_1 + k\Delta w) e^{j(w_1 + k\Delta w)(t_1 + n\Delta t)}$$

Vector Product for ICTFT

Formula for ICTFT calculation in MATLAB

$$f(t_1 + n\Delta t) = \frac{\Omega}{2\pi K} \sum_{k=0}^{K-1} F(w_1 + k\Delta w) e^{j(w_1 + k\Delta w)(t_1 + n\Delta t)}$$

Vector form for ICTFT calculation

$$f(t_1 + n\Delta t) = \frac{\Omega}{2\pi K} \begin{bmatrix} e^{jw_1(t_1 + n\Delta t)} & e^{j(w_1 + \Delta w)(t_1 + n\Delta t)} & \dots & e^{j(w_2 - \Delta w)(t_1 + n\Delta t)} \end{bmatrix} \begin{bmatrix} F(w_1) \\ F(w_1 + \Delta w) \\ \vdots \\ F(w_2 - \Delta w) \end{bmatrix} \quad (10)$$

Vector Product for ICTFT

Formula for ICTFT calculation in MATLAB

$$f(t_1 + n\Delta t) = \frac{\Omega}{2\pi K} \sum_{k=0}^{K-1} F(w_1 + k\Delta w) e^{j(w_1 + k\Delta w)(t_1 + n\Delta t)}$$

Vector form for ICTFT calculation

$$f(t_1 + n\Delta t) = \frac{\Omega}{2\pi K} \begin{bmatrix} e^{jw_1(t_1 + n\Delta t)} & e^{j(w_1 + \Delta w)(t_1 + n\Delta t)} & \dots & e^{j(w_2 - \Delta w)(t_1 + n\Delta t)} \end{bmatrix} \begin{bmatrix} F(w_1) \\ F(w_1 + \Delta w) \\ \vdots \\ f(w_2 - \Delta w) \end{bmatrix} \quad (10)$$

Pseudocode for vector product

```
for  $n = 1, \dots, N$   
     $f_n = \frac{\Omega}{2\pi K} \mathbf{b}_n^\top \mathbf{F}$   
end
```

Example 2: Vector Product for CTFT and ICTFT

Rectangular pulse

$$f(t) = \begin{cases} 1, & |x| < \frac{1}{2} \\ 0, & \text{otherwise} \end{cases} \quad (11)$$

- (1) Plot $f(t)$, $t \in [-1, 1]$;
- (2) Plot $F(w)$, $w \in [-8\pi, 8\pi]$;
- (3) Recover $f(t)$ from $F(w)$.




Figure 3.1: Waveform and spectrum

Overview

1. Symbolic Function
2. Loop Calculation
3. Vector Product
- 4. Matrix Product**
5. Experiment Problem

Matrix Product for CTFT and ICTFT-1

Matrix form for the CTFT

$$\begin{bmatrix} F(w_1) \\ F(w_1 + \Delta w) \\ \vdots \\ f(w_2 - \Delta w) \end{bmatrix} = \frac{T}{N} \begin{bmatrix} e^{-jw_1 t_1} & e^{-jw_1(t_1+\Delta t)} & \dots & e^{-jw_1(t_2-\Delta t)} \\ e^{-j(w_1+\Delta w)t_1} & e^{-j(w_1+\Delta w)(t_1+\Delta t)} & \dots & e^{-j(w_1+\Delta w)(t_2-\Delta t)} \\ \vdots & \vdots & & \vdots \\ e^{-j(w_2-\Delta w)t_1} & e^{-j(w_2-\Delta w)(t_1+\Delta t)} & \dots & e^{-j(w_2-\Delta w)(t_2-\Delta t)} \end{bmatrix} \begin{bmatrix} f(t_1) \\ f(t_1 + \Delta t) \\ \vdots \\ f(t_2 - \Delta t) \end{bmatrix} \quad (12)$$

Matrix Product for CTFT and ICTFT-1

Matrix form for the CTFT

$$\begin{bmatrix} F(w_1) \\ F(w_1 + \Delta w) \\ \vdots \\ f(w_2 - \Delta w) \end{bmatrix} = \frac{T}{N} \begin{bmatrix} e^{-jw_1 t_1} & e^{-jw_1(t_1+\Delta t)} & \dots & e^{-jw_1(t_2-\Delta t)} \\ e^{-j(w_1+\Delta w)t_1} & e^{-j(w_1+\Delta w)(t_1+\Delta t)} & \dots & e^{-j(w_1+\Delta w)(t_2-\Delta t)} \\ \vdots & \vdots & & \vdots \\ e^{-j(w_2-\Delta w)t_1} & e^{-j(w_2-\Delta w)(t_1+\Delta t)} & \dots & e^{-j(w_2-\Delta w)(t_2-\Delta t)} \end{bmatrix} \begin{bmatrix} f(t_1) \\ f(t_1 + \Delta t) \\ \vdots \\ f(t_2 - \Delta t) \end{bmatrix} \quad (12)$$

Matrix form for the ICTFT

$$\begin{bmatrix} f(t_1) \\ f(t_1 + \Delta t) \\ \vdots \\ f(t_2 - \Delta t) \end{bmatrix} = \frac{\Omega}{2\pi K} \begin{bmatrix} e^{jw_1 t_1} & e^{j(w_1+\Delta w)t_1} & \dots & e^{j(w_2-\Delta w)t_1} \\ e^{jw_1(t_1+\Delta t)} & e^{j(w_1+\Delta w)(t_1+\Delta t)} & \dots & e^{j(w_2-\Delta w)(t_1+\Delta t)} \\ \vdots & \vdots & & \vdots \\ e^{jw_1(t_2-\Delta t)} & e^{j(w_1+\Delta w)(t_2-\Delta t)} & \dots & e^{j(w_2-\Delta w)(t_2-\Delta t)} \end{bmatrix} \begin{bmatrix} F(w_1) \\ F(w_1 + \Delta w) \\ \vdots \\ f(w_2 - \Delta w) \end{bmatrix} \quad (13)$$

Matrix Product for CTFT and ICTFT-1

Matrix form for the CTFT

$$\begin{bmatrix} F(w_1) \\ F(w_1 + \Delta w) \\ \vdots \\ f(w_2 - \Delta w) \end{bmatrix} = \frac{T}{N} \begin{bmatrix} e^{-jw_1 t_1} & e^{-jw_1(t_1+\Delta t)} & \dots & e^{-jw_1(t_2-\Delta t)} \\ e^{-j(w_1+\Delta w)t_1} & e^{-j(w_1+\Delta w)(t_1+\Delta t)} & \dots & e^{-j(w_1+\Delta w)(t_2-\Delta t)} \\ \vdots & \vdots & & \vdots \\ e^{-j(w_2-\Delta w)t_1} & e^{-j(w_2-\Delta w)(t_1+\Delta t)} & \dots & e^{-j(w_2-\Delta w)(t_2-\Delta t)} \end{bmatrix} \begin{bmatrix} f(t_1) \\ f(t_1 + \Delta t) \\ \vdots \\ f(t_2 - \Delta t) \end{bmatrix} \quad (12)$$

Matrix form for the ICTFT

$$\begin{bmatrix} f(t_1) \\ f(t_1 + \Delta t) \\ \vdots \\ f(t_2 - \Delta t) \end{bmatrix} = \frac{\Omega}{2\pi K} \begin{bmatrix} e^{jw_1 t_1} & e^{j(w_1+\Delta w)t_1} & \dots & e^{j(w_2-\Delta w)t_1} \\ e^{jw_1(t_1+\Delta t)} & e^{j(w_1+\Delta w)(t_1+\Delta t)} & \dots & e^{j(w_2-\Delta w)(t_1+\Delta t)} \\ \vdots & \vdots & & \vdots \\ e^{jw_1(t_2-\Delta t)} & e^{j(w_1+\Delta w)(t_2-\Delta t)} & \dots & e^{j(w_2-\Delta w)(t_2-\Delta t)} \end{bmatrix} \begin{bmatrix} F(w_1) \\ F(w_1 + \Delta w) \\ \vdots \\ f(w_2 - \Delta w) \end{bmatrix} \quad (13)$$

Simplified matrix form

$$\mathbf{F} = \frac{T}{N} \mathbf{U} \mathbf{f}, \quad \mathbf{f} = \frac{\Omega}{2\pi K} \mathbf{V} \mathbf{F} \quad (14)$$

Matrix Product for CTFT and ICTFT-2

How to obtain \mathbf{U} and \mathbf{V} : Kronecker product

$$\begin{bmatrix} w_1 \\ w_1 + \Delta w \\ \vdots \\ w_2 - \Delta w \end{bmatrix} \otimes [t_1 \quad t_1 + \Delta t \quad \cdots \quad t_2 - \Delta t] = \begin{bmatrix} w_1 t_1 & w_1(t_1 + \Delta t) & \cdots & w_1(t_2 - \Delta t) \\ (w_1 + \Delta w)t_1 & (w_1 + \Delta w)(t_1 + \Delta t) & \cdots & (w_1 + \Delta w)(t_2 - \Delta t) \\ \vdots & \vdots & \ddots & \vdots \\ (w_2 - \Delta w)t_1 & (w_2 - \Delta w)(t_1 + \Delta t) & \cdots & (w_2 - \Delta w)(t_2 - \Delta t) \end{bmatrix} \quad (15)$$

\otimes denotes Kronecker tensor product: `kron()` in MATLAB.

Example 3: Matrix product for CTFT and ICTFT

Rectangular pulse

$$f(t) = \begin{cases} 1, & |x| < \frac{1}{2} \\ 0, & \text{otherwise} \end{cases} \quad (16)$$

- (1) Plot $f(t)$, $t \in [-1, 1]$;
- (2) Plot $F(w)$, $w \in [-8\pi, 8\pi]$;
- (3) Recover $f(t)$ from $F(w)$.




Figure 4.1: Waveform and spectrum

Overview

1. Symbolic Function
2. Loop Calculation
3. Vector Product
4. Matrix Product
5. Experiment Problem

Problem 1: CTFT and ICTFT

Triangular pulse:

$$f(t) = \begin{cases} E \left(1 - \frac{2|t|}{\tau}\right), & t \leq \frac{\tau}{2} \\ 0, & t > \frac{\tau}{2} \end{cases} \quad (17)$$

with $E = 1$ and $\tau = 1$. For $t \in [-1, 1]$ and $w \in [-50, 50]$:

- (1) Determine $g(t)$ that satisfies $f(t) = g(t) * g(t)$ and plot $f(t)$ and $g(t)$;
- (2) Plot $F(w)$, $G(w)$, and $G_e(w) = G(w) \cdot G(w)$;
- (3) Plot $F_e(w) = \text{CTFT}\{g(t) * g(t)\}$;
- (4) Compare $F(w)$, $G_e(w)$, and $F_e(w)$;
- (5) Compare the time costs of 3 methods in a bar form.




Figure 5.1: Waveform and spectrum.

Thank You!