

# Duality Formulaton of Max-Margin Classifier\*

Arash Rouhani

March 1, 2011

## Abstract

This paper presents a short mathematical explenation of how to formulate the Max-Margin Classifier problem in it's dual form. I go through the derivation step by step by formulating the lagrangian function and then performing sensible steps finally reach the desired dual formulation.

## 1 Primal formulation

To be written ...

## 2 Lagrangian function

There will be as always 3 kinds of terms, coming from either the *the objective function*, *a constraint* or *a variable constraint*.

$$L(w, b, \xi, \mu, \nu) = \frac{1}{2} \|w\|_2^2 + \sum_{i=1}^n C\xi_i + \mu_i(y_i b + 1 - y_i w^T x_i - \xi_i) - \nu_i \xi_i \quad (1)$$

We now derivativs for each variable-type and get an equation. There are 3 'types' of variables here, so we will get 3 equations.

### 2.1 Derivative with respect to w

$$\frac{dL}{dw} = 0 = w - \sum_{i=1}^n \mu_i y_i x_i \quad (2)$$

### 2.2 Derivative with respect to b

$$\frac{dL}{db} = 0 = \sum_{i=1}^n \mu_i y_i \quad (3)$$

---

\*This is the first time I use latex

### 2.3 Derivative with respect to $\xi$

$$\frac{dL}{db} = 0 = C - \mu_i - \nu_i \quad (4)$$

## 3 Reducing the lagrangian by substitution

First we substitute (4) which is clearly eliminating the  $\xi$  terms.

$$L(w, b, \mu) = \frac{1}{2} \|w\|_2^2 + \sum_{i=1}^n \mu_i (y_i b + 1 - y_i w^T x_i) \quad (5)$$

Let's rewrite the function using  $\|w\|_2^2 = w^T w$  and other simple rewritings.

$$L(w, b, \mu) = w^T \left( \frac{1}{2} w - \sum_{i=1}^n \mu_i y_i x_i \right) + b \sum_{i=1}^n \mu_i y_i + \sum_{i=1}^n \mu_i \quad (6)$$

Now we use (2) and (3) in order to reduce once more.

$$L(w, \mu) = \sum_{i=1}^n \mu_i - \frac{1}{2} w^T \sum_{i=1}^n \mu_i y_i x_i \quad (7)$$

Now we use (2) again to rewrite  $w^T$ , finally eliminating all free variables we had in the primal.

$$L(\mu) = \sum_{i=1}^n \mu_i - \frac{1}{2} \sum_{j=1}^n \mu_j y_j x_j^T \sum_{i=1}^n \mu_i y_i x_i \quad (8)$$

Now we finally rewrite the lagrangian to it's well-recognized form.

$$L(\mu) = \sum_{i=1}^n \mu_i - \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \mu_i \mu_j y_i y_j x_i^T x_j \quad (9)$$

## 4 Finding the constraints

So far we've only retrieved the objective function out of the Lagrangian, now the constraints remain. Firstly we must notice that we've always known that  $\mu, \nu > 0$ , dual variables are always  $\geq 0$  (otherwise it would be a *gain* to violate the constraints). The rest of the information we are going to work with is the equalities we got after the derivations in chapter 2.

The first constraint we take is simply equation (3). The second constraint I claim is

$$\begin{aligned} 0 &\leq \mu_i \leq C \\ 1 &\leq i \leq n \end{aligned}$$

$\mu_i \geq 0$  Holds by definition.  $C \geq \mu_i$  holds as of (4) by regulating  $\nu \geq 0$  in  $\mu_i = C - \nu_i$ . It's clear that we get the constraint claimed above.