

Continuity in the Zariski Topology

A Bridge to Algebraic Geometry

Timothy Tarter

Department of Mathematics and Statistics
James Madison University

Math 435 Final Project
November 27, 2025

Motivating Questions

Motivating Questions

- 1 For a fixed $f(x)$, how can we prove or disprove that $f(x)$ is continuous in the Zariski topology?
- 2 How do varieties X, Y relate to each other, and what can $\mathcal{O}(X)$ tell us about $\mathcal{O}(Y)$ tell us about Y ? How about $K(X)$ and $K(Y)$?
- 3 Furthermore, for a fixed variety X , what is the relationship between $\mathcal{O}(X)$, $A(X)$, and $K(X)$?
- 4 How do birational maps induce isomorphisms on open subsets, and how do local / global glueing theorems help us tie things back to topology?

Motivating Questions

- 1 For a fixed $f(x)$, how can we prove or disprove that $f(x)$ is continuous in the Zariski topology?
- 2 How do varieties X, Y relate to each other, and what can $\mathcal{O}(X)$ tell us about $\mathcal{O}(Y)$ tell us about Y ? How about $K(X)$ and $K(Y)$?
- 3 Furthermore, for a fixed variety X , what is the relationship between $\mathcal{O}(X)$, $A(X)$, and $K(X)$?
- 4 How do birational maps induce isomorphisms on open subsets, and how do local / global glueing theorems help us tie things back to topology?
- 5 Can we use restriction maps on open subsets to generalize continuity?

Motivating Questions

- 1 For a fixed $f(x)$, how can we prove or disprove that $f(x)$ is continuous in the Zariski topology?
- 2 How do varieties X, Y relate to each other, and what can $\mathcal{O}(X)$ tell us about $\mathcal{O}(Y)$ tell us about Y ? How about $K(X)$ and $K(Y)$?
- 3 Furthermore, for a fixed variety X , what is the relationship between $\mathcal{O}(X)$, $A(X)$, and $K(X)$?
- 4 How do birational maps induce isomorphisms on open subsets, and how do local / global glueing theorems help us tie things back to topology?
- 5 Can we use restriction maps on open subsets to generalize continuity?
- 6 How can we measure failures of glueability with the cohomology of sheaves? What is the topological intuition for this?

Motivating Questions

- ① For a fixed $f(x)$, how can we prove or disprove that $f(x)$ is continuous in the Zariski topology?
- ② How do varieties X, Y relate to each other, and what can $\mathcal{O}(X)$ tell us about $\mathcal{O}(Y)$ tell us about Y ? How about $K(X)$ and $K(Y)$?
- ③ Furthermore, for a fixed variety X , what is the relationship between $\mathcal{O}(X)$, $A(X)$, and $K(X)$?
- ④ How do birational maps induce isomorphisms on open subsets, and how do local / global glueing theorems help us tie things back to topology?
- ⑤ Can we use restriction maps on open subsets to generalize continuity?
- ⑥ How can we measure failures of glueability with the cohomology of sheaves? What is the topological intuition for this?

*Red entries are answered in the accompanied paper.

Getting a Grip on Affine n -Space

Intro to Affine Space

Definition

We define the **affine n-space**, \mathbb{A}^n over an algebraically closed field k to be the set of n -tuples with coordinates in k^n .

Intro to Affine Space

Definition

We define the **affine n-space**, \mathbb{A}^n over an algebraically closed field k to be the set of n -tuples with coordinates in k^n .

It's worth noting that \mathbb{A}^n is very similar to a vector space over k - but they are not the same structure.

Intro to Affine Space

Definition

We define the **affine n-space**, \mathbb{A}^n over an algebraically closed field k to be the set of n -tuples with coordinates in k^n .

It's worth noting that \mathbb{A}^n is very similar to a vector space over k - but they are not the same structure.

The key difference: \mathbb{A}^n doesn't preserve 0. In fact, affine space only preserves things invariant under linear transformation.

What Makes "Sense" in \mathbb{A}^n ?

What Makes Sense	What Doesn't Make Sense
Groups of rigid motions around a point	The origin - or any fixed point in \mathbb{A}^n
Ellipses	Circles
Parallelograms	Squares
Invariants of geometric symmetries	Angles between vectors and magnitudes of vectors

Functions and Algebraic Sets in Affine Space

Functions and Algebraic Sets in Affine Space

Definition

We define a function f by $f : \mathbb{A}^n \rightarrow k$.

Functions and Algebraic Sets in Affine Space

Definition

We define a function f by $f : \mathbb{A}^n \rightarrow k$.

Definition

We define the zero set of a set of functions $\{f_1, \dots, f_s\} = T \subseteq k[x_1, \dots, x_n]$ by

$$Z(T) = \{p \in \mathbb{A}^n ; f_i(p) = 0 \text{ for all } f_i \in T\}$$

Functions and Algebraic Sets in Affine Space

Definition

We define a function f by $f : \mathbb{A}^n \rightarrow k$.

Definition

We define the zero set of a set of functions $\{f_1, \dots, f_s\} = T \subseteq k[x_1, \dots, x_n]$ by

$$Z(T) = \{p \in \mathbb{A}^n ; f_i(p) = 0 \text{ for all } f_i \in T\}$$

Definition

$Y \subseteq \mathbb{A}^n$ is called an **algebraic set** if there exists a set of functions $T \subseteq k[x_1, \dots, x_n]$ with $Y = Z(T)$.

Introducing the Zariski Topology

Introducing the Zariski Topology

Definition

We define the Zariski topology on affine space by the following rules:

- Closed sets are algebraic sets, and,
- Open sets are the complements of algebraic sets.

Introducing the Zariski Topology

Definition

We define the Zariski topology on affine space by the following rules:

- Closed sets are algebraic sets, and,
- Open sets are the complements of algebraic sets.

It is clear that this is a topology, since, \emptyset, \mathbb{A}^n are open by fixing the constant function at zero, then at some non-zero value.

Introducing the Zariski Topology

Definition

We define the Zariski topology on affine space by the following rules:

- Closed sets are algebraic sets, and,
- Open sets are the complements of algebraic sets.

It is clear that this is a topology, since, \emptyset, \mathbb{A}^n are open by fixing the constant function at zero, then at some non-zero value.

Additionally, it is somewhat intuitive that arbitrary unions of complements of algebraic sets are still complements of algebraic sets, and that finite intersections of complements of algebraic sets are also complements of algebraic sets.

Affine Varieties

Definition

A set $Y \subseteq \mathbb{A}^n$ which is closed in the Zariski topology is called **irreducible** iff there does not exist two closed subsets of Y , Y_1 and Y_2 respectively such that $Y = Y_1 \cup Y_2$.

Affine Varieties

Definition

A set $Y \subseteq \mathbb{A}^n$ which is closed in the Zariski topology is called **irreducible** iff there does not exist two closed subsets of Y , Y_1 and Y_2 respectively such that $Y = Y_1 \cup Y_2$.

Definition

A set $Y \subseteq \mathbb{A}^n$ is called an **affine variety** if Y is an irreducible algebraic set. Open subsets of affine varieties are called **quasi-affine varieties**.

Affine Varieties

Definition

A set $Y \subseteq \mathbb{A}^n$ which is closed in the Zariski topology is called **irreducible** iff there does not exist two closed subsets of Y , Y_1 and Y_2 respectively such that $Y = Y_1 \cup Y_2$.

Definition

A set $Y \subseteq \mathbb{A}^n$ is called an **affine variety** if Y is an irreducible algebraic set. Open subsets of affine varieties are called **quasi-affine varieties**.

Algebraic geometry is (largely) the study and classification of varieties of many kinds (affine, quasi-affine, projective, and quasi-projective).

Affine Varieties

Definition

A set $Y \subseteq \mathbb{A}^n$ which is closed in the Zariski topology is called **irreducible** iff there does not exist two closed subsets of Y , Y_1 and Y_2 respectively such that $Y = Y_1 \cup Y_2$.

Definition

A set $Y \subseteq \mathbb{A}^n$ is called an **affine variety** if Y is an irreducible algebraic set. Open subsets of affine varieties are called **quasi-affine varieties**.

Algebraic geometry is (largely) the study and classification of varieties of many kinds (affine, quasi-affine, projective, and quasi-projective). We will develop certain tools (structural invariants) in this lecture and accompanied paper which can be used to classify varieties up to what is called birationality. The popular modern AG tools are called numerical invariants - usually cohomology - which (sort of) measure the genus of a simplicial complex. With topology as a frame of reference, this is a step up from the notion of homotopy classes of a space.

Examining Continuous Functions in \mathbb{A}^1

Remarks About Closed Sets in \mathbb{A}^1

Consider a function $f : \mathbb{A}^1 \rightarrow \mathbb{A}^1$. We know that $Z(f)$ *must* be a finite set by the Fundamental Theorem of Algebra.

Remarks About Closed Sets in \mathbb{A}^1

Consider a function $f : \mathbb{A}^1 \rightarrow \mathbb{A}^1$. We know that $Z(f)$ *must* be a finite set by the Fundamental Theorem of Algebra.

Conclusion: Closed sets are finite (or all of \mathbb{A}^1 or \emptyset).

Remarks About Closed Sets in \mathbb{A}^1

Consider a function $f : \mathbb{A}^1 \rightarrow \mathbb{A}^1$. We know that $Z(f)$ *must* be a finite set by the Fundamental Theorem of Algebra.

Conclusion: Closed sets are finite (or all of \mathbb{A}^1 or \emptyset).

As a corollary, we will also note that varieties in \mathbb{A}^1 are singleton sets.

An Equivalent Statement to Topological Continuity

An Equivalent Statement to Topological Continuity

Definition

$f : \mathbb{A}^n \rightarrow \mathbb{A}^m$ is continuous iff for every open set $V \subseteq \mathbb{A}^m$, $f^{-1}(V)$ is open in \mathbb{A}^n .

An Equivalent Statement to Topological Continuity

Definition

$f : \mathbb{A}^n \rightarrow \mathbb{A}^m$ is continuous iff for every open set $V \subseteq \mathbb{A}^m$, $f^{-1}(V)$ is open in \mathbb{A}^n .

Theorem (An Equivalent Statement)

$f : \mathbb{A}^n \rightarrow \mathbb{A}^m$ is continuous iff for every closed set $V \subseteq \mathbb{A}^m$, $f^{-1}(V)$ is closed in \mathbb{A}^n .

An Equivalent Statement to Topological Continuity

Definition

$f : \mathbb{A}^n \rightarrow \mathbb{A}^m$ is continuous iff for every open set $V \subseteq \mathbb{A}^m$, $f^{-1}(V)$ is open in \mathbb{A}^n .

Theorem (An Equivalent Statement)

$f : \mathbb{A}^n \rightarrow \mathbb{A}^m$ is continuous iff for every closed set $V \subseteq \mathbb{A}^m$, $f^{-1}(V)$ is closed in \mathbb{A}^n .

Proof

Let $f : \mathbb{A}^n \rightarrow \mathbb{A}^m$ be continuous. Then for every open $V \subseteq \mathbb{A}^m$, $f^{-1}(V)$ is open in \mathbb{A}^n .

An Equivalent Statement to Topological Continuity

Definition

$f : \mathbb{A}^n \rightarrow \mathbb{A}^m$ is continuous iff for every open set $V \subseteq \mathbb{A}^m$, $f^{-1}(V)$ is open in \mathbb{A}^n .

Theorem (An Equivalent Statement)

$f : \mathbb{A}^n \rightarrow \mathbb{A}^m$ is continuous iff for every closed set $V \subseteq \mathbb{A}^m$, $f^{-1}(V)$ is closed in \mathbb{A}^n .

Proof

Let $f : \mathbb{A}^n \rightarrow \mathbb{A}^m$ be continuous. Then for every open $V \subseteq \mathbb{A}^m$, $f^{-1}(V)$ is open in \mathbb{A}^n .

But $\mathbb{A}^m \setminus V = V^C$ is closed in \mathbb{A}^m .

An Equivalent Statement to Topological Continuity

Definition

$f : \mathbb{A}^n \rightarrow \mathbb{A}^m$ is continuous iff for every open set $V \subseteq \mathbb{A}^m$, $f^{-1}(V)$ is open in \mathbb{A}^n .

Theorem (An Equivalent Statement)

$f : \mathbb{A}^n \rightarrow \mathbb{A}^m$ is continuous iff for every closed set $V \subseteq \mathbb{A}^m$, $f^{-1}(V)$ is closed in \mathbb{A}^n .

Proof

Let $f : \mathbb{A}^n \rightarrow \mathbb{A}^m$ be continuous. Then for every open $V \subseteq \mathbb{A}^m$, $f^{-1}(V)$ is open in \mathbb{A}^n .

But $\mathbb{A}^m \setminus V = V^C$ is closed in \mathbb{A}^m . And $f^{-1}(V^C) = f^{-1}(\mathbb{A}^m \setminus V) = \mathbb{A}^n \setminus f^{-1}(V)$ is closed in \mathbb{A}^n .

An Equivalent Statement to Topological Continuity

Definition

$f : \mathbb{A}^n \rightarrow \mathbb{A}^m$ is continuous iff for every open set $V \subseteq \mathbb{A}^m$, $f^{-1}(V)$ is open in \mathbb{A}^n .

Theorem (An Equivalent Statement)

$f : \mathbb{A}^n \rightarrow \mathbb{A}^m$ is continuous iff for every closed set $V \subseteq \mathbb{A}^m$, $f^{-1}(V)$ is closed in \mathbb{A}^n .

Proof

Let $f : \mathbb{A}^n \rightarrow \mathbb{A}^m$ be continuous. Then for every open $V \subseteq \mathbb{A}^m$, $f^{-1}(V)$ is open in \mathbb{A}^n .

But $\mathbb{A}^m \setminus V = V^C$ is closed in \mathbb{A}^m . And $f^{-1}(V^C) = f^{-1}(\mathbb{A}^m \setminus V) = \mathbb{A}^n \setminus f^{-1}(V)$ is closed in \mathbb{A}^n . So inverse images of closed sets being closed imply topological continuity.



Putting Things Together in \mathbb{A}^1

Putting Things Together in \mathbb{A}^1

Theorem

$f : \mathbb{A}^1 \rightarrow \mathbb{A}^1$ is continuous iff inverse images of finite sets are finite.

A Strict Classification of \mathbb{A}^1

The Big Theorem

Theorem

The only functions $f : \mathbb{A}^1 \rightarrow \mathbb{A}^1$ which are continuous are non-constant polynomials.

The Big Theorem

Theorem

The only functions $f : \mathbb{A}^1 \rightarrow \mathbb{A}^1$ which are continuous are non-constant polynomials.

This really breaks into three statements:

The Big Theorem

Theorem

The only functions $f : \mathbb{A}^1 \rightarrow \mathbb{A}^1$ which are continuous are non-constant polynomials.

This really breaks into three statements:

- 1 Non-constant polynomial functions are continuous on the affine line.

The Big Theorem

Theorem

The only functions $f : \mathbb{A}^1 \rightarrow \mathbb{A}^1$ which are continuous are non-constant polynomials.

This really breaks into three statements:

- 1 Non-constant polynomial functions are continuous on the affine line.
- 2 Constant polynomial functions are not continuous on the affine line.

The Big Theorem

Theorem

The only functions $f : \mathbb{A}^1 \rightarrow \mathbb{A}^1$ which are continuous are non-constant polynomials.

This really breaks into three statements:

- 1 Non-constant polynomial functions are continuous on the affine line.
- 2 Constant polynomial functions are not continuous on the affine line.
- 3 Arbitrary functions which are not polynomial functions are not continuous on the affine line.

The Big Theorem

Theorem

The only functions $f : \mathbb{A}^1 \rightarrow \mathbb{A}^1$ which are continuous are non-constant polynomials.

This really breaks into three statements:

- 1 Non-constant polynomial functions are continuous on the affine line.
- 2 Constant polynomial functions are not continuous on the affine line.
- 3 Arbitrary functions which are not polynomial functions are not continuous on the affine line.

The Bigger Picture

The Bigger Picture

The Bigger Picture

- The notion of continuity in the Zariski topology generalizes to continuity on dense, open subsets of varieties.

The Bigger Picture

- The notion of continuity in the Zariski topology generalizes to continuity on dense, open subsets of varieties.
- In turn, this generalizes to a notion of glueability of continuous functions on intersection patches of open subsets.

The Bigger Picture

- The notion of continuity in the Zariski topology generalizes to continuity on dense, open subsets of varieties.
- In turn, this generalizes to a notion of glueability of continuous functions on intersection patches of open subsets.
- We can measure the failure of glueability of continuous data with sheaf cohomology.

The Bigger Picture

- The notion of continuity in the Zariski topology generalizes to continuity on dense, open subsets of varieties.
- In turn, this generalizes to a notion of glueability of continuous functions on intersection patches of open subsets.
- We can measure the failure of glueability of continuous data with sheaf cohomology.




marigold-22



You'll ask an algebraic geometer what a topological space is and they'll go on a tangent about how really a topological space is just something you can put a sheaf on and that a sheaf is really just something that gives you a good cohomology so really what you should be asking is what is cohomology and then not answer the question.

Github QR Code



References

- Artin, M. (2018). [AR] Algebra (1st ed.). Pearson.
- Hartshorne, R. (2010). [HS1] Algebraic geometry (1st ed.). Springer.
- John Mackintosh Howie. (2006). [HW] Fields and Galois theory. Springer.
- Vakil, R. (2025). [VK] The Rising Sea. Princeton University Press.
- Tsai, Lauren (2024). The Riemann-Roch Theorem. University of Chicago REU

Additional resources used:

<https://lekili.freemath.xyz/teaching/topics/Blowups.pdf>

<https://www.youtube.com/@HarpreetBedimath>