



ICS141: Discrete Mathematics for Computer Science I

Dept. Information & Computer Sci., University of Hawaii

Originals slides by Dr. Baek and Dr. Still, adapted by J. Stelovsky

Based on slides Dr. M. P. Frank and Dr. J.L. Gross

Provided by McGraw-Hill



Lecture 5

Chapter 1. The Foundations

1.4 Nested Quantifiers

1.5 Rules of Inference



Previously...

© The McGraw-Hill Companies, Inc. all rights reserved.

TABLE 1 Quantifiers.

<i>Statement</i>	<i>When True?</i>	<i>When False?</i>
$\forall x P(x)$	$P(x)$ is true for every x .	There is an x for which $P(x)$ is false.
$\exists x P(x)$	There is an x for which $P(x)$ is true.	$P(x)$ is false for every x .

© The McGraw-Hill Companies, Inc. all rights reserved.

TABLE 2 De Morgan's Laws for Quantifiers.

<i>Negation</i>	<i>Equivalent Statement</i>	<i>When Is Negation True?</i>	<i>When False?</i>
$\neg \exists x P(x)$	$\forall x \neg P(x)$	For every x , $P(x)$ is false.	There is an x for which $P(x)$ is true.
$\neg \forall x P(x)$	$\exists x \neg P(x)$	There is an x for which $P(x)$ is false.	$P(x)$ is true for every x .

Nesting of Quantifiers

- Example:

Let the domain of x and y be people.

Let $L(x,y)$ = “ x likes y ” (A statement with 2 free variables – not a proposition)

- Then $\exists y L(x,y)$ = “There is someone whom x likes.” (A statement with 1 free variable x – not a proposition)

- Then $\forall x (\exists y L(x,y))$ =

“Everyone has someone whom they like.”

(A Proposition with 0 free variables.)

Nested Quantifiers

- Nested quantifiers are quantifiers that occur within the scope of other quantifiers.
- The order of the quantifiers is important, unless all the quantifiers are universal quantifiers or all are existential quantifiers.

© The McGraw-Hill Companies, Inc. all rights reserved.

TABLE 1 Quantifications of Two Variables.

<i>Statement</i>	<i>When True?</i>	<i>When False?</i>
$\forall x \forall y P(x, y)$ $\forall y \forall x P(x, y)$	$P(x, y)$ is true for every pair x, y .	There is a pair x, y for which $P(x, y)$ is false.
$\forall x \exists y P(x, y)$	For every x there is a y for which $P(x, y)$ is true.	There is an x such that $P(x, y)$ is false for every y .
$\exists x \forall y P(x, y)$	There is an x for which $P(x, y)$ is true for every y .	For every x there is a y for which $P(x, y)$ is false.
$\exists x \exists y P(x, y)$ $\exists y \exists x P(x, y)$	There is a pair x, y for which $P(x, y)$ is true.	$P(x, y)$ is false for every pair x, y .

Nested Quantifiers

- Let the domain of x and y is \mathbf{R} , and $P(x,y): xy = 0$. Find the truth value of the following propositions.
 - $\forall x \forall y P(x, y)$ (F)
 - $\forall x \exists y P(x, y)$ (T)
 - $\exists x \forall y P(x, y)$ (T)
 - $\exists x \exists y P(x, y)$ (T)
- $\forall x \exists y P(x,y) \neq \exists y \forall x P(x,y)$
 - For every x , there exists y such that $x + y = 0$. (T)
 - There exists y such that, for every x , $x + y = 0$. (F)

\mathbf{R} : set of real numbers

Nested Quantifiers: Example

- Let the domain = $\{1, 2, 3\}$. Find an expression equivalent to $\forall x \exists y P(x,y)$ where the variables are bound by substitution instead:

- Expand from inside out or outside in.

- Outside in:

$$\forall x \exists y P(x,y)$$

$$\equiv \exists y P(1,y) \wedge \exists y P(2,y) \wedge \exists y P(3,y)$$

$$\equiv [P(1,1) \vee P(1,2) \vee P(1,3)] \wedge$$

$$[P(2,1) \vee P(2,2) \vee P(2,3)] \wedge$$

$$[P(3,1) \vee P(3,2) \vee P(3,3)]$$

Quantifier Exercise

- If $R(x,y)$ = “ x relies upon y ,” express the following in unambiguous English when the domain is all people

$$\forall x(\exists y R(x,y)) =$$

Everyone has *someone* to rely on.

$$\exists y(\forall x R(x,y)) =$$

There’s a poor overburdened soul whom *everyone* relies upon (including himself)!

$$\exists x(\forall y R(x,y)) =$$

There’s some needy person who relies upon *everybody* (including himself).

$$\forall y(\exists x R(x,y)) =$$

Everyone has *someone* who relies upon them.

$$\forall x(\forall y R(x,y)) =$$

Everyone relies upon *everybody*, (including themselves)!

Negating Nested Quantifiers

- Successively apply the rules for negating statements involving a single quantifier
- Example: Express the negation of the statement $\forall x \exists y (P(x,y) \wedge \exists z R(x,y,z))$ so that all negation symbols immediately precede predicates.

$$\begin{aligned} & \neg \forall x \exists y (P(x,y) \wedge \exists z R(x,y,z)) \\ & \equiv \exists x \neg \exists y (P(x,y) \wedge \exists z R(x,y,z)) \\ & \equiv \exists x \forall y \neg (P(x,y) \wedge \exists z R(x,y,z)) \\ & \equiv \exists x \forall y (\neg P(x,y) \vee \neg \exists z R(x,y,z)) \\ & \equiv \exists x \forall y (\neg P(x,y) \vee \forall z \neg R(x,y,z)) \end{aligned}$$

Equivalence Laws

- $\forall x \forall y P(x,y) \equiv \forall y \forall x P(x,y)$

$$\exists x \exists y P(x,y) \equiv \exists y \exists x P(x,y)$$

- $\forall x (P(x) \wedge Q(x)) \equiv (\forall x P(x)) \wedge (\forall x Q(x))$

$$\exists x (P(x) \vee Q(x)) \equiv (\exists x P(x)) \vee (\exists x Q(x))$$

- Exercise:

See if you can prove these yourself.



Notational Conventions

- Quantifiers have higher precedence than all logical operators from propositional logic:

$$(\forall x P(x)) \wedge Q(x)$$

- Consecutive quantifiers of the same type can be combined:

$$\forall x \forall y \forall z P(x,y,z) \equiv \forall x,y,z P(x,y,z)$$

$$\text{or even } \forall xyz P(x,y,z)$$

1.5 Rules of Inference

- ***An argument***: a sequence of statements that end with a conclusion
- Some forms of argument (“valid”) never lead from correct statements to an incorrect conclusion. Some other forms of argument (“fallacies”) can lead from true statements to an incorrect conclusion.
- ***A logical argument*** consists of a list of (possibly compound) propositions called premises/hypotheses and a single proposition called the conclusion.
- ***Logical rules of inference***: methods that depend on logic alone for deriving a new statement from a set of other statements. (Templates for constructing valid arguments.)

Valid Arguments (I)

- Example: A logical argument

If I dance all night, then I get tired.

I danced all night.

Therefore I got tired.

- Logical representation of underlying variables:

p : I dance all night. q : I get tired.

- Logical analysis of argument:

$p \rightarrow q$ premise 1

p premise 2

$\therefore q$ conclusion



Valid Arguments (II)

- A form of logical argument is ***valid*** if whenever every premise is true, the conclusion is also true. A form of argument that is not valid is called a ***fallacy***.



Inference Rules: General Form

- An *Inference Rule* is
 - A pattern establishing that if we know that a set of *premise* statements of certain forms are all true, then we can validly deduce that a certain related *conclusion* statement is true.

$$\begin{array}{l} \textit{premise 1} \\ \textit{premise 2} \\ \dots \\ \hline \therefore \textit{conclusion} \end{array}$$

“ \therefore ” means “therefore”

Inference Rules & Implications

- Each valid logical inference rule corresponds to an implication that is a tautology.

$\begin{array}{l} \textit{premise 1} \\ \textit{premise 2} \\ \dots \\ \hline \therefore \textit{conclusion} \end{array}$	Inference rule
---	-----------------------

- Corresponding tautology:
$$((\textit{premise 1}) \wedge (\textit{premise 2}) \wedge \dots) \rightarrow \textit{conclusion}$$

Modus Ponens



$$\begin{array}{l} p \\ p \rightarrow q \\ \hline \therefore q \end{array}$$

Rule of ***Modus ponens***
(a.k.a. *law of detachment*)

“the mode of
affirming”

- $(p \wedge (p \rightarrow q)) \rightarrow q$ is a tautology

p	q	$p \rightarrow q$	$p \wedge (p \rightarrow q)$	$(p \wedge (p \rightarrow q)) \rightarrow q$
T	T	T	T	T
T	F	F	F	T
F	T	T	F	T
F	F	T	F	T

- Notice that the first row is the only one where premises are all true

Modus Ponens: Example

If $\left\{ \begin{array}{l} p \rightarrow q : \text{"If it snows today} \\ \text{then we will go skiing"} \\ p : \text{"It is snowing today"} \end{array} \right\}$ assumed TRUE

Then $\therefore q$: "We will go skiing" is TRUE

If $\left\{ \begin{array}{l} p \rightarrow q : \text{"If } n \text{ is divisible by 3} \\ \text{then } n^2 \text{ is divisible by 3"} \\ p : \text{"} n \text{ is divisible by 3"} \end{array} \right\}$ assumed TRUE

Then $\therefore q$: " n^2 is divisible by 3" is TRUE

Modus Tollens



$$\frac{\neg q \quad p \rightarrow q}{\therefore \neg p}$$

Rule of *Modus tollens*

“the mode of denying”

- $(\neg q \wedge (p \rightarrow q)) \rightarrow \neg p$ is a tautology

- Example

If $\left\{ \begin{array}{l} p \rightarrow q : \text{“If this jewel is really a diamond} \\ \text{then it will scratch glass”} \\ \neg q : \text{“The jewel doesn’t scratch glass”} \end{array} \right\}$ assumed TRUE

Then $\therefore \neg p$: “The jewel is not a diamond” is TRUE

More Inference Rules

- $$\frac{p}{\therefore p \vee q}$$

Rule of **Addition**

Tautology: $p \rightarrow (p \vee q)$

- $$\frac{p \wedge q}{\therefore p}$$

Rule of **Simplification**

Tautology: $(p \wedge q) \rightarrow p$

- $$\frac{p}{q} \quad \frac{q}{\therefore p \wedge q}$$

Rule of **Conjunction**

Tautology: $[(p) \wedge (q)] \rightarrow p \wedge q$



Examples

- State which rule of inference is the basis of the following arguments:
 - It is below freezing now. Therefore, it is either below freezing or raining now.
 - It is below freezing and raining now. Therefore, it is below freezing now.
- p : It is below freezing now.
 q : It is raining now.
 - $p \rightarrow (p \vee q)$ (rule of addition)
 - $(p \wedge q) \rightarrow p$ (rule of simplification)

Hypothetical Syllogism

- $$\begin{array}{l} p \rightarrow q \\ q \rightarrow r \\ \hline \therefore p \rightarrow r \end{array}$$

Rule of **Hypothetical syllogism**

Tautology:

$$[(p \rightarrow q) \wedge (q \rightarrow r)] \rightarrow (p \rightarrow r)$$

- Example: State the rule of inference used in the argument:

“If it rains today, ^{*p*} then ^{*q*} we will not have a ^{*q*} barbecue today. If ^{*q*} we do not have a barbecue today, then ^{*r*} we will have a barbecue tomorrow. Therefore, if it rains today, then ^{*r*} we will have a barbecue tomorrow.” ^{*p*}

Disjunctive Syllogism

- $$\begin{array}{c} p \vee q \\ \neg p \\ \hline \therefore q \end{array}$$

Rule of ***Disjunctive syllogism***

Tautology: $[(p \vee q) \wedge (\neg p)] \rightarrow q$

- Example

- Ed's wallet is in his back pocket or it is on his desk. $(p \vee q)$ p q
- Ed's wallet is not in his back pocket. $(\neg p)$
- Therefore, Ed's wallet is on his desk. (q)