



Meiosis and genetic variation



Meiosis and genetic variation

Genome: Complete sequence of an organism's DNA.




Includes **genes** (control traits) and non-coding DNA organized in **chromosomes**.



Mary E. Cline (CCHN)

Genes

- Eukaryotic DNA is organized in chromosomes.
 - DNA sequence that produces a particular proteins
 - Genes have specific places on chromosomes.



Heredity


- Heredity - way of transferring genetic information to offspring
- Chromosome theory of heredity: chromosomes carry genes.
- Gene - "unit of heredity".



Reproduction

■ Asexual

- ❑ Many single-celled organisms reproduce by splitting, budding, parthenogenesis.
- ❑ Some multicellular organisms can reproduce asexually, produce **clones** (*offspring genetically identical to parent*).




Sexual reproduction

- Fusion of two gametes to produce a single zygote.
- Introduces greater genetic variation, allows genetic recombination.
- With exception of self-fertilizing organisms (e.g. some plants), zygote has gametes from two different parents.

Chromosomes


- Karyotype:
 - ordered display of an individual's chromosomes.
 - Collection of chromosomes from mitotic cells.
 - Staining can reveal visible band patterns, gross anomalies.

Karyotyping



Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.


Human male
G-bands



Male


Female

Human Female
G-bands



Homologues

- Chromosomes exist in homologous pairs in diploid cells.









Exception: **Sex chromosomes** (X, Y).

Other chromosomes are known as **autosomes**, they have homologues.

In humans ...


- 23 chromosomes donated by each parent (total = 46 or 23 pairs).
- Gametes (sperm/ova):
 - Contain 22 autosomes and 1 sex chromosome.
 - Are haploid (haploid number " n " = 23 in humans).
- Fertilization/syngamy results in zygote with 2 haploid sets of chromosomes - now **diploid**.
 - Diploid cell; $2n = 46$. ($n=23$ in humans)
- Most cells in the body produced by mitosis.
- Only gametes are produced by meiosis.

Chromosome numbers

	Organism	Number of chromosomes
	pea plant	14
	sun flower	34
	cat	38
	puffer fish	42
	human	46
	dog	78

All are **even** numbers -
diploid ($2n$) sets of
homologous
chromosomes!

Ploidy = number of
copies of each
chromosome.




MEIOSIS I: Separates homologous chromosomes

INTERPHASE

PROPHASE I

METAPHASE I

ANAPHASE I



MEIOSIS II: Separates sister chromatids


**TELOPHASE I
AND CYTOKINESIS**

PROPHASE II


METAPHASE II

ANAPHASE II

**TELOPHASE II
AND CYTOKINESIS**



Meiosis



Meiosis 1

First division of meiosis

- **Prophase 1:** Each chromosome duplicates and remains closely associated. These are called sister chromatids. Crossing-over can occur during the latter part of this stage.
- **Metaphase 1:** Homologous chromosomes align at the equatorial plate.
- **Anaphase 1:** Homologous pairs separate with sister chromatids remaining together.
- **Telophase 1:** Two daughter cells are formed with each daughter containing only one chromosome of the homologous pair.

Meiosis II


Second division of meiosis: *Gamete formation*

- **Prophase 2:** DNA does not replicate.
- **Metaphase 2:** Chromosomes align at the equatorial plate.
- **Anaphase 2:** Centromeres divide and sister chromatids migrate separately to each pole.
- **Telophase 2:** Cell division is complete. Four haploid daughter cells are obtained.

Meiosis – key differences from mitosis

- Meiosis reduces the number of chromosomes by half.
- Daughter cells differ from parent, and each other.
- Meiosis involves two divisions, Mitosis only one.
- Meiosis I involves:
 - **Synapsis** - homologous chromosomes pair up. **Chiasmata** form (crossing over of non-sister chromatids).
 - In Metaphase I, **homologous pairs line up** at metaphase plate.
 - In Anaphase I, *sister chromatids do NOT separate*.
 - Overall, **separation of homologous pairs of chromosomes**, rather than sister chromatids of individual chromosome.

Mitosis vs. meiosis



SUMMARY

Event	Mitosis	Meiosis
DNA replication	Occurs during interphase before nuclear division begins	Occurs once, during the interphase before meiosis I begins
Number of divisions	One, including prophase, metaphase, anaphase, and telophase	Two, each including prophase, metaphase, anaphase, and telophase
Synapsis of homologous chromosomes	Does not occur	Synapsis is unique to meiosis: During prophase I, the homologous chromosomes join along their length, forming tetrads (groups of four chromatids); synapsis is associated with crossing over between nonsister chromatids
Number of daughter cells and genetic composition	Two, each diploid ($2n$) and genetically identical to the parent cell	Four, each haploid (n), containing half as many chromosomes as the parent cell; genetically nonidentical to the parent cell and to each other
Role in the animal body	Enables multicellular adult to arise from zygote; produces cells for growth and tissue repair	Produces gametes; reduces chromosome number by half and introduces genetic variability among the gametes


Meiosis creates genetic variation

- During normal cell growth, ***mitosis* produces daughter cells identical to parent cell** ($2n$ to $2n$)
- *Meiosis* results in **genetic variation** by **shuffling** of maternal and paternal chromosomes and **crossing over**.

No daughter cells formed during meiosis are genetically identical to either mother or father


During sexual reproduction, fusion of the unique haploid gametes produces truly unique offspring.

Independent assortment



Independent assortment

Number of combinations: 2^n




e.g. 2 chromosomes in haploid

$$2n = 4; n = 2$$

$2^n = 2^2 = 4$ possible combinations

In humans




e.g. 23 chromosomes in haploid

$2n = 46$; $n = 23$

$2^n = 2^{23} = \sim 8$ million possible combinations!

Crossing over



Chiasmata – sites of crossing over, occur in **synapsis**.
Exchange of genetic material between non-sister chromatids.



Crossing over produces **recombinant chromosomes**.



Random fertilization

At least 8 million combinations from Mom,
and another 8 million from Dad ...


>64 trillion combinations for a diploid
zygote!!!



Copyright © Pearson Education, Inc., publishing as Benjamin Cummings

Meiosis & sexual life cycles

- **Life cycle** = sequence of stages in organisms reproductive history; conception to reproduction.
- **Somatic** cells = any cell other than gametes, most of the cells in the body.
- **Gametes** produced by meiosis.



Generalized animal life cycle