

# MA204: Mathematics IV

## Complex Analysis: Some Elementary Functions

Gautam Kalita  
IIIT Guwahati

# Exponential function

From the Euler's formula, we have

$$e^{iy} = \cos y + i \sin y.$$

If  $z = x + iy$ , then the exponential function is defined as

$$e^z = e^{x+iy} = e^x e^{iy} = e^x (\cos y + i \sin y).$$

As a result,

$$|e^z| = e^x \text{ and } \arg(e^z) = y + 2n\pi \text{ for } n \in \mathbb{Z}.$$

# Exponential function

From the Euler's formula, we have

$$e^{iy} = \cos y + i \sin y.$$

If  $z = x + iy$ , then the exponential function is defined as

$$e^z = e^{x+iy} = e^x e^{iy} = e^x (\cos y + i \sin y).$$

As a result,

$$|e^z| = e^x \text{ and } \arg(e^z) = y + 2n\pi \text{ for } n \in \mathbb{Z}.$$

## Properties:

- (a)  $e^{z_1+z_2} = e^{z_1} e^{z_2}$ .
- (b)  $e^z \neq 0$  for all  $z \in \mathbb{C}$ .
- (c)  $e^z$  is an entire function and  $\frac{d}{dz} e^z = e^z$ .

## Exponential function

From the Euler's formula, we have

$$e^{iy} = \cos y + i \sin y.$$

If  $z = x + iy$ , then the exponential function is defined as

$$e^z = e^{x+iy} = e^x e^{iy} = e^x (\cos y + i \sin y).$$

As a result,

$$|e^z| = e^x \text{ and } \arg(e^z) = y + 2n\pi \text{ for } n \in \mathbb{Z}.$$

### Properties:

- (a)  $e^{z_1+z_2} = e^{z_1} e^{z_2}$ .
- (b)  $e^z \neq 0$  for all  $z \in \mathbb{C}$ .
- (c)  $e^z$  is an entire function and  $\frac{d}{dz} e^z = e^z$ .

**Question:** Is  $e^z$  a bijection?

## Exponential function

**Problem:** Find complex values of  $w$  for which  $e^{z+w} = e^z$ . Can you say something about period of the exponential function?

# Exponential function

**Problem:** Find complex values of  $w$  for which  $e^{z+w} = e^z$ . Can you say something about period of the exponential function?

**Problem:** Solve the equation  $e^z = -1$ .

## Exponential function

**Problem:** Find complex values of  $w$  for which  $e^{z+w} = e^z$ . Can you say something about period of the exponential function?

**Problem:** Solve the equation  $e^z = -1$ .

**Problem:** If  $z = x + iy$ , then show that  $|e^{2z+i} + e^{iz^2}| \leq e^{2x} + e^{-2xy}$ .

## Exponential function

**Problem:** Find complex values of  $w$  for which  $e^{z+w} = e^z$ . Can you say something about period of the exponential function?

**Problem:** Solve the equation  $e^z = -1$ .

**Problem:** If  $z = x + iy$ , then show that  $|e^{2z+i} + e^{iz^2}| \leq e^{2x} + e^{-2xy}$ .

**Problem:** Give conditions on  $z$  such that  $e^z$  is (a) real (b) purely imaginary.



## Complex Logarithm

It is easy to see that the exponential function  $e^z : \mathbb{C} \rightarrow \mathbb{C}$  is neither one-one as  $e^{z+2ni\pi} = e^z$  nor onto. Thus the inverse of the function does not exist in the complex plane  $\mathbb{C}$ .

## Complex Logarithm

It is easy to see that the exponential function  $e^z : \mathbb{C} \rightarrow \mathbb{C}$  is neither one-one as  $e^{z+2ni\pi} = e^z$  nor onto. Thus the inverse of the function does not exist in the complex plane  $\mathbb{C}$ .

However, the exponential function  $e^z : \mathbb{C} \rightarrow \mathbb{C} - \{0\}$  is on-to. For any  $z \in \mathbb{C} - \{0\}$ , we have  $e^w = z$ . As a result,  $w := \log z = \ln |z| + i \arg(z)$ .

## Complex Logarithm

It is easy to see that the exponential function  $e^z : \mathbb{C} \rightarrow \mathbb{C}$  is neither one-one as  $e^{z+2ni\pi} = e^z$  nor onto. Thus the inverse of the function does not exist in the complex plane  $\mathbb{C}$ .

However, the exponential function  $e^z : \mathbb{C} \rightarrow \mathbb{C} - \{0\}$  is on-to. For any  $z \in \mathbb{C} - \{0\}$ , we have  $e^w = z$ . As a result,  $w := \log z = \ln |z| + i \arg(z)$ .

Thus we can define the **complex logarithm** as

$$\log z = \ln |z| + i \arg(z).$$

## Complex Logarithm

It is easy to see that the exponential function  $e^z : \mathbb{C} \rightarrow \mathbb{C}$  is neither one-one as  $e^{z+2ni\pi} = e^z$  nor onto. Thus the inverse of the function does not exist in the complex plane  $\mathbb{C}$ .

However, the exponential function  $e^z : \mathbb{C} \rightarrow \mathbb{C} - \{0\}$  is on-to. For any  $z \in \mathbb{C} - \{0\}$ , we have  $e^w = z$ . As a result,  $w := \log z = \ln |z| + i \arg(z)$ .

Thus we can define the **complex logarithm** as

$$\log z = \ln |z| + i \arg(z).$$

Clearly, the complex logarithm  $\log z$  is not well defined. It is multi-valued assignment.

The **principal value** of the logarithm is defined as

$$\text{Log } z = \ln |z| + i \text{Arg}(z).$$

## Complex Logarithm

It is easy to see that the exponential function  $e^z : \mathbb{C} \rightarrow \mathbb{C}$  is neither one-one as  $e^{z+2ni\pi} = e^z$  nor onto. Thus the inverse of the function does not exist in the complex plane  $\mathbb{C}$ .

However, the exponential function  $e^z : \mathbb{C} \rightarrow \mathbb{C} - \{0\}$  is on-to. For any  $z \in \mathbb{C} - \{0\}$ , we have  $e^w = z$ . As a result,  $w := \log z = \ln |z| + i \arg(z)$ .

Thus we can define the **complex logarithm** as

$$\log z = \ln |z| + i \arg(z).$$

Clearly, the complex logarithm  $\log z$  is not well defined. It is multi-valued assignment.

The **principal value** of the logarithm is defined as

$$\text{Log } z = \ln |z| + i \text{Arg}(z).$$

Note that  $\text{Log } z$  is well defined single valued function.

# Complex logarithm

**Note:**

- (a) If  $z \neq 0$ , then  $e^{\log z} = z$ . What about  $\log(e^z)$ ?
- (b) The function  $\text{Log } z$  is not continuous on the negative real axis.
- (c) The function  $\text{Log } z$  is analytic everywhere except on the negative real axis and at zero.
- (d)  $\text{Log}(z_1 z_2) = \text{Log } z_1 + \text{Log } z_2$ . What if  $\text{Log}$  is replaced by  $\log$ ?
- (e) For  $z_2 \neq 0$ , we have  $\text{Log}\left(\frac{z_1}{z_2}\right) = \text{Log } z_1 - \text{Log } z_2$ . What if  $\text{Log}$  is replaced by  $\log$ ?

## Complex logarithm

**Note:**

- (a) If  $z \neq 0$ , then  $e^{\log z} = z$ . What about  $\log(e^z)$ ?
- (b) The function  $\text{Log } z$  is not continuous on the negative real axis.
- (c) The function  $\text{Log } z$  is analytic everywhere except on the negative real axis and at zero.
- (d)  $\text{Log}(z_1 z_2) = \text{Log } z_1 + \text{Log } z_2$ . What if  $\text{Log}$  is replaced by  $\log$ ?
- (e) For  $z_2 \neq 0$ , we have  $\text{Log}\left(\frac{z_1}{z_2}\right) = \text{Log } z_1 - \text{Log } z_2$ . What if  $\text{Log}$  is replaced by  $\log$ ?

**Question:** What is the derivative of  $\log z$ ?

# Complex logarithm

**Note:**

- (a) If  $z \neq 0$ , then  $e^{\log z} = z$ . What about of  $\log(e^z)$ ?
- (b) The function  $\text{Log } z$  is not continuous on the negative real axis.
- (c) The function  $\text{Log } z$  is analytic everywhere except on the negative real axis and at zero.
- (d)  $\text{Log}(z_1 z_2) = \text{Log } z_1 + \text{Log } z_2$ . What if  $\text{Log}$  is replaced by  $\log$ ?
- (e) For  $z_2 \neq 0$ , we have  $\text{Log}\left(\frac{z_1}{z_2}\right) = \text{Log } z_1 - \text{Log } z_2$ . What if  $\text{Log}$  is replaced by  $\log$ ?

**Question:** What is the derivative of  $\log z$ ?

**Question:** Find valued of  $\log(1 + i)$ ,  $\text{Log}(-i)$ .



## Branch, Branch cut, and Branch point

**Branch of a multiple valued function:** Let  $F$  be a multiple valued function defined on a domain  $D$ . A function  $f$  is said to be a branch of the multiple valued function  $F$  if in a domain  $D_0 \subset D$  if  $f(z)$  is single valued and analytic in  $D_0$ .

## Branch, Branch cut, and Branch point

**Branch of a multiple valued function:** Let  $F$  be a multiple valued function defined on a domain  $D$ . A function  $f$  is said to be a branch of the multiple valued function  $F$  if in a domain  $D_0 \subset D$  if  $f(z)$  is single valued and analytic in  $D_0$ .

**Branch Cut:** The portion of a line or a curve introduced in order to define a branch of a multiple valued function is called branch cut.

## Branch, Branch cut, and Branch point

**Branch of a multiple valued function:** Let  $F$  be a multiple valued function defined on a domain  $D$ . A function  $f$  is said to be a branch of the multiple valued function  $F$  if in a domain  $D_0 \subset D$  if  $f(z)$  is single valued and analytic in  $D_0$ .

**Branch Cut:** The portion of a line or a curve introduced in order to define a branch of a multiple valued function is called branch cut.

**Branch Point:** Any point that is common to all branch cuts is called a branch point.

## Complex exponent

For complex numbers  $z \neq 0$  and  $w$ , the exponent  $w$  to  $z$  is defined as

$$z^w = e^{w \log z}.$$

## Complex exponent

For complex numbers  $z \neq 0$  and  $w$ , the exponent  $w$  to  $z$  is defined as

$$z^w = e^{w \log z}.$$

The complex exponent  $z^w$  is a multi-valued function.

## Complex exponent

For complex numbers  $z \neq 0$  and  $w$ , the exponent  $w$  to  $z$  is defined as

$$z^w = e^{w \log z}.$$

The complex exponent  $z^w$  is a multi-valued function.

The **principal branch** or the **principal value** of the complex exponent  $z^w$  is given by

$$\text{P.V.}(z^w) = e^{w \text{Log } z}.$$

## Complex exponent

**Problem:** Find values of  $i^{-i}$ ,  $(1+i)^i$ , and  $i^{(1-i)}$ .

## Complex exponent

**Problem:** Find values of  $i^{-i}$ ,  $(1+i)^i$ , and  $i^{(1-i)}$ .

**Problem:** Discuss analyticity for the function  $f(w) = z^w$  and  $g(z) = z^w$ . Find derivative of the functions.



## Trigonometric function

Note that  $e^{ix} = \cos x + i \sin x$  and  $e^{-ix} = \cos x - i \sin x$ . Thus, we have

$$\sin x = \frac{e^{ix} - e^{-ix}}{2i} \text{ and } \cos x = \frac{e^{ix} + e^{-ix}}{2}.$$

As a result, it is natural to extend the definition as follows:

$$\sin z = \frac{e^{iz} - e^{-iz}}{2i} \text{ and } \cos z = \frac{e^{iz} + e^{-iz}}{2}.$$

## Trigonometric function

Note that  $e^{ix} = \cos x + i \sin x$  and  $e^{-ix} = \cos x - i \sin x$ . Thus, we have

$$\sin x = \frac{e^{ix} - e^{-ix}}{2i} \text{ and } \cos x = \frac{e^{ix} + e^{-ix}}{2}.$$

As a result, it is natural to extend the definition as follows:

$$\sin z = \frac{e^{iz} - e^{-iz}}{2i} \text{ and } \cos z = \frac{e^{iz} + e^{-iz}}{2}.$$

### Properties:

- (a) Both  $\cos z$  and  $\sin z$  are entire functions.
- (b) Except few, all elementary properties of real trigonometric functions are also satisfied by complex trigonometric functions.

## Trigonometric function

Note that  $e^{ix} = \cos x + i \sin x$  and  $e^{-ix} = \cos x - i \sin x$ . Thus, we have

$$\sin x = \frac{e^{ix} - e^{-ix}}{2i} \text{ and } \cos x = \frac{e^{ix} + e^{-ix}}{2}.$$

As a result, it is natural to extend the definition as follows:

$$\sin z = \frac{e^{iz} - e^{-iz}}{2i} \text{ and } \cos z = \frac{e^{iz} + e^{-iz}}{2}.$$

### Properties:

- (a) Both  $\cos z$  and  $\sin z$  are entire functions.
- (b) Except few, all elementary properties of real trigonometric functions are also satisfied by complex trigonometric functions.

**Question:** Is  $\cos z$  or  $\sin z$  bounded?

## Trigonometric function

Note that  $e^{ix} = \cos x + i \sin x$  and  $e^{-ix} = \cos x - i \sin x$ . Thus, we have

$$\sin x = \frac{e^{ix} - e^{-ix}}{2i} \text{ and } \cos x = \frac{e^{ix} + e^{-ix}}{2}.$$

As a result, it is natural to extend the definition as follows:

$$\sin z = \frac{e^{iz} - e^{-iz}}{2i} \text{ and } \cos z = \frac{e^{iz} + e^{-iz}}{2}.$$

### Properties:

- (a) Both  $\cos z$  and  $\sin z$  are entire functions.
- (b) Except few, all elementary properties of real trigonometric functions are also satisfied by complex trigonometric functions.

**Question:** Is  $\cos z$  or  $\sin z$  bounded?

Thus we can define  $\tan$ ,  $\cot$ ,  $\sec$ , and  $\csc$  function in the usual way.

# Trigonometric function

**Problem:** Show that  $1 + \tan^2 z = \sec^2 z$ .

## Trigonometric function

**Problem:** Show that  $1 + \tan^2 z = \sec^2 z$ .

**Problem:** Find values of  $z$  for which (a)  $\sin z = -2$  (b)  $\cos z = k$

## Hyperbolic trigonometric function

One can define the hyperbolic sine and cosine functions as

$$\sinh z = \frac{e^z - e^{-z}}{2} \text{ and } \cosh z = \frac{e^z + e^{-z}}{2}.$$

Thus it is easy to see that

$$\sinh z = i \sin iz \text{ and } \cosh z = \cos iz.$$

## Hyperbolic trigonometric function

One can define the hyperbolic sine and cosine functions as

$$\sinh z = \frac{e^z - e^{-z}}{2} \text{ and } \cosh z = \frac{e^z + e^{-z}}{2}.$$

Thus it is easy to see that

$$\sinh z = i \sin iz \text{ and } \cosh z = \cos iz.$$

(a)  $\cosh z$  and  $\sinh z$  are entire functions.



## Hyperbolic trigonometric function

One can define the hyperbolic sine and cosine functions as

$$\sinh z = \frac{e^z - e^{-z}}{2} \text{ and } \cosh z = \frac{e^z + e^{-z}}{2}.$$

Thus it is easy to see that

$$\sinh z = i \sin iz \text{ and } \cosh z = \cos iz.$$

(a)  $\cosh z$  and  $\sinh z$  are entire functions.

**Can you state some properties of these hyperbolic trigonometric functions?**

## Hyperbolic trigonometric function

One can define the hyperbolic sine and cosine functions as

$$\sinh z = \frac{e^z - e^{-z}}{2} \quad \text{and} \quad \cosh z = \frac{e^z + e^{-z}}{2}.$$

Thus it is easy to see that

$$\sinh z = i \sin iz \quad \text{and} \quad \cosh z = \cos iz.$$

(a)  $\cosh z$  and  $\sinh z$  are entire functions.

**Can you state some properties of these hyperbolic trigonometric functions?**

$\cos^2 z + \sin^2 z = 1$	$\cosh^2 z - \sinh^2 z = 1$
$\sin(-z) = -\sin z, \cos(-z) = \cos z$	
$\sin(z + 2k\pi) = \sin z, \cos(z + 2k\pi) = \cos z$	
$\sin z = 0$ iff $z = n\pi$	
$\cos z = 0$ iff $z = (2n + 1)\frac{\pi}{2}$	
$\frac{d}{dz} \sin z = \cos z, \frac{d}{dz} \cos z = -\sin z$	

## Hyperbolic trigonometric function

One can define the hyperbolic sine and cosine functions as

$$\sinh z = \frac{e^z - e^{-z}}{2} \quad \text{and} \quad \cosh z = \frac{e^z + e^{-z}}{2}.$$

Thus it is easy to see that

$$\sinh z = i \sin iz \quad \text{and} \quad \cosh z = \cos iz.$$

(a)  $\cosh z$  and  $\sinh z$  are entire functions.

**Can you state some properties of these hyperbolic trigonometric functions?**

$\begin{aligned} \cos^2 z + \sin^2 z &= 1 \\ \sin(-z) &= -\sin z, \cos(-z) = \cos z \\ \sin(z + 2k\pi) &= \sin z, \cos(z + 2k\pi) = \cos z \\ \sin z &= 0 \text{ iff } z = n\pi \\ \cos z &= 0 \text{ iff } z = (2n + 1)\frac{\pi}{2} \\ \frac{d}{dz} \sin z &= \cos z, \frac{d}{dz} \cos z = -\sin z \end{aligned}$	$\cosh^2 z - \sinh^2 z = 1$
--	-----------------------------

$$\cos z = \cos(x + iy) = \cos x \cosh y - i \sin x \sinh y$$

$$\sin z = \sin(x + iy) = \sin x \cosh y + i \cos x \sinh y$$

## Hyperbolic trigonometric function

**Problem:** Solve (a)  $\sin z = \cosh 4$  (b)  $\sinh z = -i$  (c)  $\cosh z = -2$ .

**Problem:** Find values of (a)  $\tan^{-1}(1 + i)$  (b)  $\sinh^{-1}(-i)$ .

# Thank You

## Any Question!!!