

Logarithms Questions for CAT

All rights reserved. No part of this publication may be reproduced, distributed, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, or stored in any retrieval system of any nature without the permission of cracku.in, application for which shall be made to support@cracku.in

Questions

Instructions

For the following questions answer them individually

Question 1

If $log_32, log_3(2^x-5), log_3(2^x-7/2)$ are in arithmetic progression, then the value of x is equal to

- **A** 5
- **B** 4
- **C** 2
- **D** 3

Answer: D

Explanation:

$$2log(2^x - 5) = log2 + log(2^x - 7/2)$$

Let
$$2^x = 1$$

$$=> (t-5)^2 = 2(t-7/2)$$

$$\Rightarrow t^2 + 25 - 10t = 2t - 7$$

$$\Rightarrow t^2 - 12t + 32 = 0$$

Therefore, x = 2 or 3, but $2^x > 5$, so x = 3

Question 2

Let $u=(\log_2 x)^2-6\log_2 x+12$ where x is a real number. Then the equation $x^u=256$, has

- A no solution for x
- **B** exactly one solution for x
- **C** exactly two distinct solutions for x
- **D** exactly three distinct solutions for x

Answer: B

Explanation:

$$x^u=256$$

Taking log to the base 2 on both the sides,

$$u * \log_2 x = \log_2 256$$

$$=>[(\log_2 x)^2 - 6\log_2 x + 12] * \log_2 x = 8$$

$$(\log_2 x)^3 - 6(\log_2 x)^2 + 12\log_2 x = 8$$

Let $log_2x = t$

$$t^3 - 6t^2 + 12t - 8 = 0$$

$$(t-2)^3 = 0$$

Therefore, $log_2x=2$

 $\Rightarrow x = 4$ is the only solution

Hence, option B is the correct answer.

Question 3

If $logyx = (a * log_zy) = (b * log_xz) = ab$, then which of the following pairs of values for (a, b) is not possible?

- (-2, 1/2)
- (1,1)
- (0.4, 2.5)
- $(\pi, 1/\pi)$
- (2,2)

Answer: E

Explanation:

$$logyx = ab$$

$$a * log_z y = ab \Rightarrow log_z y = b$$

$$b * log_x z = ab \Rightarrow log_x z = a$$

$$logyx = logzy * logxz \Rightarrow logx/logy = logy/logz * logz/logx$$

$$=> log x = log y$$

$$\Rightarrow logy = logx$$

$$\Rightarrow (log x)^2 = (log y)^2$$

=>
$$logx = logy$$
 or $logx = -logy$

So,
$$x = y \text{ or } x = 1/y$$

So,
$$ab = 1 \text{ or } -1$$

Option 5) is not possible



CAT 2023 STUDY PLAN



Question 4

If x >= y and y > 1, then the value of the expression $log_x(x/y) + log_y(y/x)$ can never be

- A -1
- -0.5
- 0
- D 1

Answer: D

Explanation:

$$log_x(x/y) + log_y(y/x) = 1 - log_x(y) + 1 - log_y(x)$$

$$= 2 - (log_x y + 1/log_x y) \le 0$$
 (Since $log_x y + 1/log_x y \ge 2$)

So, the value of the expression cannot be 1

Question 5

If $\log_2\log_7{(x^2-x+37)}$ = 1, then what could be the value of 'x'?

- B
- None of these

Answer: C

Explanation:

$$\log_2 \log_7 (x^2 - x + 37) =$$

$$\log_7(x^2 - x + 37) = 2$$

$$(x^2 - x + 37) = 7^2$$

Given eq. can be reduced to $x^2 - x + 37 = 49$

So x can be either -3 or 4.

Question 6

Suppose, $\log_3 x = \log_{12} y = a$, where x,y are positive numbers. If G is the geometric mean of x and y, and $\log_6 G$ is equal to

- A \sqrt{a}
- **B** 2a
- **C** a/2
- **D** a

Answer: D

Explanation:

We know that $\log_3 x = a$ and $\log_{12} y = a$

Hence,
$$x=3^a$$
 and $y=12^a$

Therefore, the geometric mean of x and y equals \sqrt{x}

This equals
$$\sqrt{3^a imes 12^a} = 6^a$$

Hence, $G=6^a$ Or, $\log_6 G=a$

Take Free CAT Mock (with video Solutions)

Question 7

The value of $\log_{0.008}\sqrt{5}+\log\sqrt{3}\,81-7$ is equal to

- **A** 1/3
- **B** 2/3
- **C** 5/6
- **D** 7/6

Answer: C

Explanation:

$$\log_{0.008} \sqrt{5} + \log_{\sqrt{3}} 81 - 7$$

$$81 = 3^4$$
 and $0.008 = {}^{8}_{1000} = {}^{2^3}_{10^3} = {}^{1}_{5^3} = 5^{-3}$

Hence,

$$\log_{0.008} \sqrt{5} + 8 - 7$$

$$\log_{5^{-3}} 5^{\frac{1}{2}} + 8 - 7$$

$$^{log5^{0.5}}_{log5^{-3}}+1$$

$$-\frac{1}{6} + 1$$



If x is a real number such that $\log_3 5 = \log_5 (2+x)$, then which of the following is true?

- 0 < x < 3
- 23 < x < 30
- x > 30
- 3 < x < 23

Answer: D

Explanation:

$$\begin{aligned} &1 < \log_3 5 < 2 \\ &\Rightarrow 1 < \log_5 (2+x) < 2 \\ &\Rightarrow 5 < 2+x < 25 \\ &\Rightarrow 3 < x < 23 \end{aligned}$$

Question 9

If $log(2^a \times 3^b \times 5^c)$ is the arithmetic mean of $log(2^2 \times 3^3 \times 5)$, $log(2^6 \times 3 \times 5^7)$, and $log(2 \times 3^2 \times 5^4)$, then a equals

Answer:3

Explanation:

Explanation:
$$log(2^a \times 3^b \times 5^c) = \frac{log(2^2 \times 3^3 \times 5) + log(2^6 \times 3 \times 5^7) + log(2 \times 3^2 \times 5^4)}{3}$$

$$log(2^a \times 3^b \times 5^c) = \frac{log(2^{2+6+1} \times 3^{3+1+2} \times 5^{1+7+4})}{3}$$

$$log(2^a \times 3^b \times 5^c) = \frac{log(2^9 \times 3^6 \times 5^{12})}{3}$$

$$3log(2^a \times 3^b \times 5^c)$$
 = $log(2^9 \times 3^6 \times 5^{12})$

Hence, 3a = 9 or a = 3

SCHOLARSHIP WAIVER UPTO ₹ 7000 OFF

CAT 2023 COACHING BY IIM ALUMNI

Question 10

If x is a positive quantity such that $2^x = 3^{\log_5 2}$, then x is equal to

- $\log_5 8$
- $1 + \log_3(\frac{5}{3})$
- $\log_5 9$
- $1 + \log_5(\frac{3}{5})$

Answer: D

Explanation:

Givne that: $2^x = 3^{\log_5 2}$

$$\Rightarrow 2^x = 2^{\log_5 3}$$

$$\Rightarrow x = \log_5 3$$

$$3 * 5$$

$$\Rightarrow x = \log_5$$
 5





Take Free CAT Mock (with video Solutions)

SCHOLARSHIP WAIVER
UPTO ₹ 7000 OFF

CAT 2023 COACHING
BY IIM ALUMNI

CAT 2023 Coaching By IIM Alumni

CAT Previous Papers PDF



Free CAT Exam Preparation App

CAT Formulas PDF [Download Now]

Top-500 Free CAT Questions (With Solutions)


Free CAT Study Material

Free CAT Preparation Video Lectures

Daily Free CAT Practice Tests



Free Videos for CAT Preparation



Important Verbal Ability Questions for CAT (Download PDF)

Data Interpretation for CAT Questions (download pdf)

Logical Reasoning for CAT Questions (download pdf)

Quantitative Aptitude for CAT Questions (download pdf)

Know the CAT Percentile Required for IIM Calls

Join MBA Telegram Group

Enroll for Excellent CAT/MBA courses

Cracku CAT Success Stories

Subscribe to MBA Exams Youtube Channel